

Electronic Supplementary Information

to

Orange peel derived nanobiochar for targeted cancer therapy

Daniela Iannazzo,^{1*} Consuelo Celesti,^{1,2*} Claudia Espro,¹ Angelo Ferlazzo,¹ Salvatore V. Giofrè,³ Mario Scuderi,⁴ Silvia Scalese,⁴ Bartolo Gabriele,⁵ Raffaella Mancuso,⁵ Ida Zicarelli,⁵ Giuseppa Visalli,⁶ Angela Di Pietro⁶

¹ *Department of Engineering, University of Messina, Contrada Di Dio, I-98166 Messina, Italy;*

² *Department of Clinical and Experimental Medicine, University of Messina, Via Consolare Valeria, I-98125 Messina, Italy*

³ *Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, Viale F. Stagno d'Alcontres, 98166 Messina, Italy*

⁴ *Institute for Microelectronics and Microsystems, National Research Council (CNR-IMM), Ottava Strada n.5, I-95121 Catania, Italy*

⁵ *Laboratory of Industrial and Synthetic Organic Chemistry (LISOC), Department of Chemistry and Chemical Technologies, University of Calabria, Via Pietro Bucci 12/C, 87036 Arcavacata di Rende (CS), Italy*

⁶ *Department of Biomedical and Dental Sciences and Morphological and Functional Images, University Hospital of Messina, Via Consolare Valeria, 1, 98100 Messina, Italy*

**Correspondence: diannazzo@unime.it; ccelesti@unime.it*

Contents

Figure S1. Representative CLSM image of A549 cells treated for 24 h with NBC-B.....S2




Figure S1. Representative CLSM image of A549 cells treated for 24 h with NBC-B to assess the cell uptake.