

Oleic Acid-Based Self Micro-Emulsifying Delivery System for Enhancing Antifungal Activities of Clotrimazole

Ting-Lun Yang ¹, Chien-Ming Hsieh ², Ling-Jei Meng ¹, Tsuimin Tsai ³ and Chin-Tin Chen ^{1,*}


¹ Department of Biochemical Science and Technology, College of Life Science, National Taiwan University, Taipei 106, Taiwan; f05b22054@ntu.edu.tw (T.-L.Y.); r08b22021@ntu.edu.tw (L.-J.M.)

² School of Pharmacy, College of Pharmacy, Taipei Medical University, Taipei 110, Taiwan; cmhsieh@tmu.edu.tw

³ FormuRx Pharmaceuticals Co. Ltd., Taipei 10617, Taiwan; tmtsai00@gmail.com

* Correspondence: chintin@ntu.edu.tw; Tel.: +886-2-3366-9487; Fax: +886-2-3366-2271

C. albicans



Fluconazole-resistant *C. albicans*




Figure S1. Representative plate of disk diffusion assay showing inhibition zones against wild-type and fluconazole resistant *C. albicans* by F35 loading with or without CLT and commercial CLT cream, Mycosten®. After inoculating 2×10^6 CFU/mL of wild-type and fluconazole-resistant *C. albicans* on RPMI 1640 medium, each group was treated with F35 and Mycosten® containing 0.1 mg or 0.5 mg CLT, then incubate at 37 °C for 48 h.




Figure S2. Representative plate of disk diffusion assay showing inhibition zones against wild-type and fluconazole resistant *C. tropicalis* by F35 loading with or without CLT and commercial CLT cream, Mycosten®. After inoculating 2×10^6 CFU/mL of wild-type and fluconazole-resistant *C. tropicalis* on RPMI 1640 medium, each group was treated with F35 and Mycosten® containing 0.1 mg or 0.5 mg CLT, then incubate at 37°C for 48 h.




Figure S3. Representative plate of disk diffusion assay showing inhibition zones against wild-type and fluconazole resistant *C. albicans* and *C. tropicalis* by F35 gel loading with or without 0.1 mg CLT and commercial CLT cream, Mycosten®. After inoculating 2×10^6 CFU/mL of *C. albicans*, fluconazole-resistant *C. albicans*, *C. tropicalis* and fluconazole-resistant *C. tropicalis* on RPMI 1640 medium, each group was treated with F35 gel and Mycosten® containing 0.1 mg CLT, then incubate at 37 °C for 48 h.