

Supplementary Material

Fish Scale Gelatin Nanofibers with *Helichrysum Italicum* and *Lavandula Latifolia* Essential Oils for Bioactive Wound-Healing Dressings

Carmen Gaidau ¹, Maria Râpă ^{2,*}, Maria Stanca ^{1,*}, Mariana-Luiza Tanase ³, Laura Olariu ³, Rodica Roxana Constantinescu ¹, Andrada Lazea-Stoyanova ⁴, Cosmin-Andrei Alexe ¹ and Madalina Tudorache ⁵

- ¹ The National Research & Development Institute for Textiles and Leather-Division Leather and Footwear Research Institute, Bucharest 31251, Romania; carmen.gaidau@icpi.ro (C.G.); rodica.roxana@yahoo.com (R.R.C.); cosminandrei.alex@icpi.ro (C.-A.A.)
 - ² Faculty of Materials Science and Engineering, National University of Science and Technology Politehnica Bucharest, Bucharest 060042, Romania; maria.rapa@upb.ro
 - ³ SC Biotehnos SA, 3–5 Gorunului Street, Otopeni 075100, Romania; luiza.craciun@biotehnos.com (M.-L.T.); lolariu@biotehnos.com (L.O.)
 - ⁴ National Institute for Laser, Plasma and Radiation Physics, 409 Atomistilor Street, Magurele, Ilfov 077125, Romania; andrada@infim.ro
 - ⁵ Laboratory for Quality Control and Process Monitoring, Faculty of Chemistry, University of Bucharest, 4-12 Regina Elisabeta Boulevard, Bucharest 030018, Romania; madalina.sandulescu@g.unibuc.ro
- * Correspondence: maria.rapa@upb.ro (M.R.); maria.stanca@icpi.ro (M.S.)




Figure S1. Chemical structure for: (a) Neryl acetate; (b) Linalool.



Figure S2. Emulsions of lavender and immortelle essential oils.



Figure S3. Electrospinning of fish scale gelatin dispersions on polypropylene sheet.






Figure S4. GP chromatograms for: (a) F-EDTA, and (b) F+EDTA.



Relative front (Rf)

(a)



Relative front (Rf)

(b)

Figure S5. SDS-PAGE electrophoresis gels band analyses and the molecular weight distribution of fish scale gelatins. (a) F-EDTA; (b) F+EDTA.