

Supplementary Material

Phenolic Fingerprint, Bioactivity and Nanoformulation of *Prunus spinosa* L. Fruit Extract for Skin Delivery

Maria De Luca ^{1,2}, Carlo Ignazio Giovanni Tuberioso ³, Ramon Pons ⁴, María Teresa García ⁴,
Maria del Carmen Morán ^{5,6}, Giulio Ferino ⁷, Antonio Vassallo ^{1,8}, Giuseppe Martelli ¹, Carla Caddeo ^{3,*}

¹ Department of Science, University of Basilicata, Viale dell'Ateneo Lucano 10, 85100 Potenza, Italy

² KAMABIO Srl, Via Al Boschetto 4/B, 39100 Bolzano, Italy

³ Department of Life and Environmental Sciences, University of Cagliari, SS 554 – bivio per Sestu, Monserrato, 09042 Cagliari, Italy

⁴ Department of Surfactants and Nanobiotechnology, Institute for Advanced Chemistry of Catalonia (IQAC-CSIC), c/Jordi Girona, 18-26, 08034 Barcelona, Spain

⁵ Department of Biochemistry and Physiology, Physiology section, Faculty of Pharmacy and Food Science, University of Barcelona, Avda. Joan XXIII 27-31, 08028 Barcelona, Spain

⁶ Institute of Nanoscience and Nanotechnology - IN2UB, University of Barcelona, Avda. Diagonal, 645, 08028 Barcelona, Spain

⁷ CeSAR, University of Cagliari, SS 554 – bivio per Sestu, Monserrato, 09042 Cagliari, Italy

⁸ Spinoff TNcKILLERS s.r.l., Viale dell'Ateneo Lucano 10, 85100 Potenza, Italy

* Correspondence: caddeoc@unica.it




Figure S1. LC-DAD chromatogram of *P. spinosa* extract at $\lambda = 520$ nm. Peaks identification: A1: cyanidin-3-*O*-glucoside, A2: cyanidin-3-*O*-rutinoside, A3: peonidin-3-*O*-glucoside, and A4: peonidin-3-*O*-rutinoside. Chromatographic conditions are reported in the main text.