

# Response to Skinner: Risk stratification in hypertrophic cardiomyopathy: Time to think about the elec- trocardiogram

Juan Pablo Kaski

University College London and Great  
Ormond Street Hospital, UK

We thank Prof Skinner for his thoughtful comments on our review article in this Journal.<sup>1</sup> We welcome his recognition of the need for prospective collaborative studies in this important field; the ability to identify which children with hypertrophic cardiomyopathy (HCM) would benefit most from implantable cardioverter-defibrillator (ICD) therapy would represent a significant advance in paediatric heart muscle disease clinical care. In a rare disease with a relatively low prevalence of events, large-scale multicentre collaborations are essential to answer this question and many others. The Australian,<sup>2</sup> North American<sup>3</sup> and, most recently, UK<sup>4</sup> cohorts are testament to this. We agree that the electrocardiogram (ECG) may be of use in risk stratification in paediatric HCM, and acknowledge not just the work from Sweden<sup>5,6</sup> referenced in Skinner's commentary, but also innovative work from the Bologna group.<sup>7</sup> As highlighted in our previous meta-analysis, however, the difficulty lies in the fact different studies have investigated different ECG parameters, resulting in inconsistent data.<sup>8</sup> Furthermore, it is unclear whether the addition of ECG parameters to existing risk

algorithms would improve their performance. This hypothesis, of course, needs to be evaluated in large collaborative studies. We have recently established an international paediatric HCM consortium, involving 40 expert centres and with over 1600 patients recruited to date, with the aim of investigating established and novel risk factors for childhood HCM, including the 12-lead ECG - we would be delighted if Prof Skinner would join us in this endeavour.

## References

1. Norrish G, Kaski JP. Risk of sudden cardiac death in childhood hypertrophic cardiomyopathy: Time to solve the mystery. *Cardiogenetics* 2018;8:7201.
2. Nugent AW, Daubeney PE, Chondros P, et al. Clinical features and outcomes of childhood hypertrophic cardiomyopathy: results from a national population-based study. *Circulation* 2005;112:1332-8.
3. Colan SD, Lipshultz SE, Lowe AM, et al. Epidemiology and cause-specific outcome of hypertrophic cardiomyopathy in children: findings from the Pediatric Cardiomyopathy Registry. *Circulation* 2007;115:773-81.
4. Norrish G, Field E, McLeod K, et al. Clinical presentation and survival of childhood hypertrophic cardiomyopathy: a retrospective study in United Kingdom. *Eur Heart J* 2018 [Epub ahead of print].
5. Ostman-Smith I, Sjöberg G, Rydberg A, et al. Predictors of risk for sudden death in childhood hypertrophic cardiomy-

Correspondence: Juan Pablo Kaski,  
University College London and Great Ormond  
Street Hospital, UK.  
E-mail: j.kaski@ucl.ac.uk

Key words: Hypertrophic cardiomyopathy;  
electrocardiogram; risk stratification.

Received for publication: 28 December 2018.  
Accepted for publication: 22 January 2019.

This work is licensed under a Creative  
Commons Attribution NonCommercial 4.0  
License (CC BY-NC 4.0).

©Copyright J.P. Kaski, 2019  
Licensee PAGEPress, Italy  
*Cardiogenetics* 2019; 9:8048  
doi:10.4081/cardiogenetics.2019.8048

- opathy: the importance of the ECG risk score. *Open Heart* 2017;4:e000658.
6. Ostman-Smith I, Wisten A, Nylander E, et al. Electrocardiographic amplitudes: a new risk factor for sudden death in hypertrophic cardiomyopathy. *Eur Heart J* 2010;31:439-49.
  7. Biagini E, Pazzi C, Olivetto I, et al. Usefulness of electrocardiographic patterns at presentation to predict long-term risk of cardiac death in patients with hypertrophic cardiomyopathy. *Am J Cardiol* 2016;118:432-9.
  8. Norrish G, Cantarutti N, Pissaridou E, et al. Risk factors for sudden cardiac death in childhood hypertrophic cardiomyopathy: a systematic review and meta-analysis. *Eur J Prev Cardiol* 2017; 24:1220-30.