

Supplementary file

Enhancement of Methanogenic Activity in Volumetrically Undersized Reactor by Mesophilic Co-Digestion of Sewage Sludge and Aqueous Residue

Vincenzo Torretta ¹, Athanasia K. Tolkou ², Ioannis A. Katsoyiannis ², Francesca Maria Caccamo ³, Marco Carnevale Miino ³, Marco Baldi ⁴ and Maria Cristina Collivignarelli ^{3,5}

¹ Department of Theoretical and Applied Sciences, Insubria University of Varese, via G.B. Vico, 46, 21100 Varese, Italy; vincenzo.torretta@uninsubria.it (V.T.)

² Department of Chemistry, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece; tolka-tha@chem.auth.gr (A.K.T.); katsogia@chem.auth.gr (I.A.K.)

³ Department of Civil Engineering and Architecture, University of Pavia, via Ferrata 3, 27100 Pavia, Italy; francescamaria.caccamo01@universitadipavia.it (F.M.C.); marco.carnevalemiino01@universitadipavia.it (M.C.M.); mcristina.collivignarelli@unipv.it (M.C.C.)

⁴ Department of Chemistry, University of Pavia, viale Taramelli 12, 27100 Pavia, Italy; marco.baldi@unipv.it (M.B.)

⁵ Interdepartmental Centre for Water Research, University of Pavia, via Ferrata 3, 27100, Pavia, Italy

* Correspondence: tolkatha@chem.auth.gr (A.K.T.); Tel.: (+30)-2310-997997

SUPPLEMENTARY FILE S1




Figure S1. Conductivity in the anaerobic digester during the experimentation.