Supplementary Information

The Sewage Sludge Biochar and Its Composts Influence the Phosphate Sorption in an Alkaline–Calcareous Soil

Rabia Abdur Rehman ¹, Muhammad Farooq Qayyum ^{1,*}, Ghulam Haider ², Kate Schofield ³, Muhammad Abid ¹, Muhammad Rizwan ⁴ and Shafaqat Ali ^{4,5,*}

- Department of Soil Science, Faculty of Agricultural Sciences & Technology, Bahauddin Zakariya University, Multan 60800, Pakistan; rabima-lik075@gmail.com (R.A.R.); muhammadabid@bzu.edu.pk (M.A.)
- ² Department of Plant Biotechnology, Atta-ur-Rahman School Applied Biosciences, National University of Sciences and Technology (NUST), Islamabad 44000, Pakistan; ghulam.haider@asab.nust.edu.pk
- ³ Biogeochemistry Research Centre, B504 Portland Square, University of Plymouth, Drake Circus, Plymouth PL4 8AA, UK; kate.schofield@plymouth.ac.uk
- ⁴ Department of Environmental Sciences and Engineering, Government College University, Faisalabad 38000, Pakistan; mrazi1532@yahoo.com
- ⁵ Department of Biological Sciences and Technology, China Medical University, Taichung 40402, Taiwan
- * Correspondence: farooq.qayyum@bzu.edu.pk (M.F.Q.); shafaqataligill@gcuf.edu.pk (S.A.)




Figure S1. X-ray diffraction pattern of different treatment mixtures.




Figure S2. Sorption equation fitted using Langmuir (5 d incubation samples).




Figure S3. Sorption equation fitted using Langmuir (45 d incubation samples).




Figure S4. Sorption equation fitted using Freundlich (5 d incubation samples).




Figure S5. Sorption equation fitted using Freundlich (45 d incubation samples).