



Review

# Particularities of Fungicides and Factors Affecting Their Fate and Removal Efficacy: A Review

Georgios D. Gikas <sup>1</sup>, Paraskevas Parlakidis <sup>2</sup>, Theodoros Mavropoulos <sup>1</sup> and Zisis Vryzas <sup>2,\*</sup>

<sup>1</sup> Laboratory of Ecological Engineering and Technology, Department of Environmental Engineering, School of Engineering, Democritus University of Thrace, 67100 Xanthi, Greece; ggkikas@env.duth.gr (G.D.G.); theodoremaupoulos@gmail.com (T.M.)

<sup>2</sup> Laboratory of Agricultural Pharmacology and Ecotoxicology, Department of Agricultural Development, Democritus University of Thrace, 68200 Orestias, Greece; pparlaki@agro.duth.gr

\* Correspondence: zvryzas@agro.duth.gr

**Abstract:** Systemic fungicide use has increased over the last decades, despite the susceptibility of resistance development and the side effects to human health and the environment. Although herbicides and insecticides are detected more frequently in environmental samples, there are many fungicides that have the ability to enter water bodies due to their physicochemical properties and their increasing use. Key factors affecting fungicide fate in the environment have been discussed, including the non-target effects of fungicides. For instance, fungicides are associated with the steep decline in bumblebee populations. Secondary actions of certain fungicides on plants have also been reported recently. In addition, the use of alternative eco-friendly disease management approaches has been described. Constructed Wetlands (CWs) comprise an environmentally friendly, low cost, and efficient fungicide remediation technique. Fungicide removal within CWs is dependent on plant uptake and metabolism, absorption in porous media and soil, hydrolysis, photodegradation, and biodegradation. Factors related to the efficacy of CWs on the removal of fungicides, such as the type of CW, plant species, and the physicochemical parameters of fungicides, are also discussed in this paper. There are low-environmental-risk fungicides, phytohormones and other compounds, which could improve the removal performance of CW vegetation. In addition, specific parameters such as the multiple modes of action of fungicides, side effects on substrate microbial communities and endophytes, and plant physiological response were also studied. Prospects and challenges for future research are suggested under the prism of reducing the risk related to fungicides and enhancing CW performance.

**Keywords:** constructed wetlands; fungicides; water contamination; plant physiological responses; microbiome



**Citation:** Gikas, G.D.; Parlakidis, P.; Mavropoulos, T.; Vryzas, Z. Particularities of Fungicides and Factors Affecting Their Fate and Removal Efficacy: A Review. *Sustainability* **2022**, *14*, 4056. <https://doi.org/10.3390/su14074056>

Academic Editor: Zhihua Xiao

Received: 4 March 2022

Accepted: 27 March 2022

Published: 29 March 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Fungi pathogens can cause significant crop losses and consequently an essential reduction in the global food supply. Grains, fruits, and vegetables are amongst the crops demanding extensive use of fungicides during the growing season in order to be protected from diseases [1]. Furthermore, their application is very common in postharvest packaging plants, parks located in urban areas, and protected forest areas. Fungicide use has increased rapidly over the last decade. Approximately four hundred thousand tons of fungicides are applied globally, which represents 17.5% of global pesticide applications [2]. However, fungicide sales exceed 40% of the total pesticide sales in the European Union; the sales of inorganic fungicides comprise 54%, and organic ones 46%, of which 14.1% are (dithio)carbamates, 6.7% imidazoles and triazoles, 1.3% benzimidazoles, 0.8% morpholines, and 23.1% other fungicides and bactericides [3].

Fungicides can enter natural waters through agricultural wastewater from washing or loading of spraying equipment, leaks occurring due to improper maintenance of empty

containers (point source pollution), and diffusion through run-off, spraying drift, leaching, and subsurface drainage [1].

Fungicides include a broad range of compounds with different modes of action belonging to various chemical classes. Acute and/or chronic toxicity of organic fungicides to aquatic or terrestrial organisms have been reported in many studies. The extensive use of chemical pesticides in agriculture also raises concerns for public health. Exposure experiments in rats showed endocrine-disrupting, biochemical, histopathological, and hematological effects [4]. Several fungicides are characterized as hazardous chemicals by the World Health Organization (WHO) and are banned in the European Union.

Several factors, such as intensive farming, resistance management, market needs, new diseases, phytotoxicity, and safety to humans and other organisms, drive companies to invest heavily in innovative fungicide formulations. Therefore, the registration of new fungicides is constant, and the addition of innovative characteristics such as systemicity, persistent activity, low resistance development risk, and public health and environmental safety, is necessary. Some fungicides can also act against fungi, indirectly influencing positively the host plant physiological responses.

The introduction of alternative organic fungicide formulations promises various advantages, such as higher efficacy and lower risk for pathogens to develop resistance to fungicides. Disease management has improved due to the recent introduction of nanoscience and the development of nanofungicides. The main advantages of these nanometer particle sized formulations are the lower application dose of active ingredients, higher effectiveness, pathogen target specificity, and lower distribution to the environment [5]. However, nanofungicides can cause side effects on non-target organisms, and remediation techniques have not been developed, as nanofungicides' environmental fate has not been studied at a large scale [6]. Another recent advance in disease management is the introduction of chiral fungicides; however, both the European Commission and the United States Environmental Protection Agency (US EPA) point out that crucial information gaps exist in their environmental risk assessment [7]. The more recent use of chemical plant defense or resistance activators is another plant antifungal approach based on plant immunity activation rather than targeting the fungi directly [8].

Biofungicides are proposed as another alternative environmentally friendly, non-chemical approach to plant disease control, avoiding many of the associated side effects [9]. Biofungicides consist of two groups: microorganism-based pesticides (microbial fungicides) and plant extracts or plant-based-product pesticides (biochemical fungicides). Both of these groups can successfully manage various pathogen species, such as *Botrytis* spp., *Sclerotinia sclerotiorum*, *Monilinia* spp., and *Phytophthora* spp. [10,11]. Biofungicides are reported as eco-friendly compounds that do not present phytotoxic and other adverse effects. Their antifungal effectiveness depends on interactions with the host plants [12]. Another low environmental risk strategy is the use of chelating agents, which are known as fertilizer products; they can increase the bioavailability of fungicides to pathogens. Recent studies indicate that chelation is a promising tool for disease management [13,14].

The development of novel fungicides is dynamic. As a result, new prospects and challenges have arisen regarding their environmental fate and removal from contaminated systems by remediation techniques such as constructed wetlands (CWs). Various researchers have reported that CWs can effectively manage pesticide contamination, including fungicides [15–17]. Pesticide fate in the CW environment is a result of different natural processes such as biodegradation, hydrolysis, photolysis, adsorption, plant uptake, and sedimentation. The efficiency of CWs on fungicide removal depends on the physicochemical properties of the target compound, such as the half-life in water and soil (DT50), the affinity of fungicides to organic carbon  $K_{oc}$ , octanol/water partition coefficient ( $K_{ow}$ ), and water solubility. Plant species, hydraulic retention time, vegetation mass, hydraulic loading rate, microbial biofilm, and porous composition can be classified as the key CW characteristics that can determine their efficiency [18–20]. However, CW system performance can be limited by low plant uptake and the inhibition of microbial activity. There is recent evidence

that favors the induction of microorganisms to a CW system. The bioaugmentation of CWs is a recent regulation strategy for enriching or protecting microbial activity and enhancing plant metabolism functions against xenobiotics [17,19].

Studies on fungicides' environmental fate and ecological risk assessments for the non-target organisms are not as extensive as for herbicides and insecticides. As a result, researchers underestimate the remediation strategies related to fungicide removal. In the last decade, several review studies have described the environmental fate and the side-effects on non-target organisms caused by conventional fungicide use, while novel fungicides, such as nanofungicides and chiral fungicides, have not been thoroughly investigated. Furthermore, the recent introduction of plant metabolism boosters in the agricultural sector, for instance biofungicides and exogenous phytohormones, gives insights into their possible utilization in CWs to improve the removal performance and their interactions between them and CW biological parameters. Consequently, the aim of this study is to point out possible research gaps or underreported issues in environmental risk assessments and the respective mitigation attempts using CWs. Additionally, this study aims to highlight potential CW strategies for fungicide removal, focusing on interactions between CW plants and microorganisms and fungicide exposure.

## 2. Environmental Fate and Risk of Chemical Fungicides

### 2.1. Inorganic Fungicides

The most studied and used inorganic fungicides are copper and sulfur. Copper and sulfur are presented in numerous forms and are applied in various crops to control important foliar fruit diseases. Various copper and sulfur formulations for organic farming have been authorized by the European Union. Following their biogeochemical cycle, inorganic fungicides can run off to surface water and be adsorbed in the soil and sediments. Copper and sulfur are important trace metals for organisms' fundamental functions, yet large concentrations can be harmful [21].

When copper oxide is dissolved in water, the dominant and biocidal oxidation ion is  $\text{Cu}^{2+}$  [22]. Acidity and salinity play an important role in aquatic biota toxicity. High pH values result in a reduction of available hydrogen ions, which leads to copper toxicity. Therefore, copper ions can be attached at aquatic organisms' cells. Other physicochemical factors that influence the toxicity levels are the dissolved organic matter and dissolved water organic carbon content [21,23]. As Beck and Saundo-Wilhelmy [24] have reported, the tendency of sediments to adsorb heavy metals is high, and thus sediments can facilitate the availability of toxic chemicals in the water and aquatic organisms. Some species have a high level of sensitivity to copper, whereas others can efficiently overcome it. Copper is bioaccumulated in fish, decapod crustaceans, and algae and stored in bivalves, barnacles, and aquatic insects. The most sensitive species to copper exposure are cyanobacteria, while coccolithophores and dinoflagellates have a lower sensitivity to copper, and diatoms present resistance to copper [25].

Copper cannot be degraded in soil but can be accumulated through copper-based degradation compounds occurring in different forms. Copper's mobility in the soil profile is characterized as medium to low. A recent study reported that the high concentrations of copper in vineyards soils and groundwater was caused mainly by copper-based fungicide use, negatively affecting water quality and food safety [26]. Copper residues in soil could cause toxic effects on macro- and microorganisms, adversely influencing the various beneficial interactions in soil, such as pesticide biodegradation, soil structure, nutrition availability for plants, and pathogen resistance [27]. Element copper is able to cause toxicity to beneficial bacteria and fungi in the environment [28]. Diaz-Ravina et al. [29] reported that the microbiocidal activity in vineyard soil can be significantly reduced by high application rates and prolonged use of copper-based pesticides. For instance, high-dose application of Cu-based pesticides can have negative effects on arbuscular mycorrhiza fungi (AMF) [30]. Schoffer et al. [31] reported that copper soil pollution is more common in countries or

regions that have not enacted regulations for copper-based pesticides applications, which consequently follow only commercial formulation guidelines.

Sulfur can be found in various forms in the environment, such as gas (i.e.,  $\text{SO}_3$ ) and salt (i.e.,  $\text{MgSO}_4$ ), which are created through bacterial physiological processes. In sediments and soils, sulfur can be found as a trace element or in an inorganic form. Sulfur can cause toxicity to bacteria and fungi that are beneficial to the environment, which are not considered as crop pests. In addition, studies indicate that sulfur can be phytotoxic to some plants, such as cucurbits, apricots, and raspberries [32,33]. Kuklinska et al. [28] also reported that *Vibrio fischeri* is sensitive to sulfur exposure. The available information regarding sulfur interaction with organisms, its toxicity threshold, and its environmental fate is limited compared to copper.

## 2.2. Organic Fungicides

The environmental fate of organic fungicides (Table 1) depends on various physico-chemical parameters, such as ionization ( $\text{pK}_a$ ), water solubility, volatility,  $K_{ow}$ , and half-life in soil and water ( $\text{DT}_{50}$ ). Soil texture, organic carbon content, pH, clay mineral type, dissolved organic matter, and cation exchange capacity also play an essential role in the environmental fate of fungicides, defining processes such as run-off to surface water, adsorption, or leaching. In addition, rainfall, irrigation, microbiological degradation, hydrolysis, photolysis, and application rate could affect fungicide fate [34,35].

**Table 1.** List of common chemical fungicides [36,37].

Chemical Family	Group Name	Active Substances	Target Site	Mode of Action
Acylalanines	Phenyl Amides (PA)	metalaxyl	RNA polymerase I	Disruption of nucleic acid synthesis-RNA polymerase 1
Thiophanates	Methyl Benzimidazole Carbamates (MBC)	thiophanate	protein B1: $\beta$ -tubuline assembly in mitosis	Inhibition of mitosis and cell division (Beta-tubulin assembly in mitosis)
Pyridine-carboxamides	Succinate-dehydrogenase inhibitors (SDHI)	boscalid	complex II: succinate-dehydrogenase	Inhibition of mitochondrial ATP production in fungal cells
Pyridinyl-ethyl-benzamides	Succinate-dehydrogenase inhibitors (SDHI)	fluopyram	complex II: succinate-dehydrogenase	Succinate dehydrogenase inhibition within mitochondria blocking electron transport
Methoxy-carbamates	Quinone outside Inhibitors (QoI)	pyraclostrobin	complex III: cytochrome bc1	Respiration inhibitor of QoI
Dicarboximides	dicarboximides	iprodione	MAP/Histidine-Kinase in osmotic signal transduction	Signal transduction inhibitor
Triazoles	De-Methylation Inhibitors (DMI)	tebuconazole	C14- demethylase in sterol biosynthesis	Sterol 14-demethylase enzyme inhibition in membranes
Ethyl phosphonates	phosphonates	fosetyl-Al	phosphonates	Mycelial growth and spore production—Plant's defense elicitor
Dithio-carbamates and relatives	dithio-carbamates and relatives	mancozeb	multi-site contact activity	Chemicals with multi-site activity

Fungicide residues in surface water (e.g., streams, lakes, rivers) and groundwater have been detected by many monitoring studies worldwide. The majority of these studies were focused on a few fungicides of local importance. The extensive (multiple applications and high doses) use of fungicides in specific crops (e.g., vineyards, horticulture, orchards, etc.) can lead to pollution of nearby natural waters from fungicide residues. Hence, the spatial and temporal distribution of fungicide residues in surface waters varies throughout the year and amongst agroecosystem compartments. Usually, the highest concentrations of curative fungicides are detected during growing or preharvest seasons, whereas preventative fungicides are found at early plant growth stages and during the winter period. Regions planted with grape and tree crops have received high application doses of fungicides, resulting in high detection frequency and high concentrations of fungicides in the ecosystems [1,38].

The presence of various pesticides has been investigated in vineyard groundwater bodies in northern Italy. The environmental quality standard set by the EU (0.1 µg/L) was exceeded by five fungicides (metalaxyl-M, fluopicolide, penconazole, tetraconazole, and dimetomorph), presenting significantly high concentrations. The maximum concentrations of metalaxyl-M and penconazole were 8.015 µg/L and 18.72 µg/L, respectively [35]. In addition, a similar monitoring study was conducted in Spanish vineyards, where the detection frequency of metalaxyl, dimethomorph, and penconazole reached 50%. Moreover, the highest concentration was observed for same fungicides (metalaxyl and penconazole) [39]. These results indicate that the extensive use of fungicides in vineyards can cause an essential surface and groundwater pollution.

Papadakis et al. [34] conducted a pesticide monitoring study in two river basins in North Greece, with corn, cotton and cereals as the main crops, over a two-year period. Twenty-nine fungicides were detected at least once, while multiple detections (7 to 10 times) of boscalid, diphenylamine, etridiazole, and hexachlorobenzene were also observed. Extremely high concentrations for seven fungicides (azoxystrobin, diphenylamine, etridiazole, propiconazole, tebuconazole, quintozene, and difenoconazole), ranging from 0.153 to 0.819 µg/L, were identified. In the worst-case scenario, the risk quotient index was higher than one for four fungicides. The results showed that fungicides contribute to ecotoxicological risk for river basins.

The presence of 24 fungicides was investigated in the surface water and sediments of a horticulture area in Australia. The agricultural activity of the studied area included tree fruits, bulbs, vineyards, vegetables, and herbs. Although the authors reported that the individual fungicide residues did not pose environmental risks, due to low ecotoxicological endpoints, several fungicides were detected in concentrations above 0.2 µg/L (iprodione, myclobutanil, pyrimethanil, cyproconazole, trifloxystrobin, and fenarimol) and others had a detection frequency ranging between 18 and 36% (myclobutanil, trifloxystrobin, pyrimethanil, difenoconazole, and metalaxyl). The temporal distribution of residues was affected by the chemical class of the fungicides. Preventing fungicides were detected across the whole season, and curative fungicides mostly in March or October [40]. Although agricultural activity is the main source of pollution, urban and industrial activities can pollute the environment as well. Merel et al. [41] confirmed that the presence of carbendazim in the Rhine river (west Germany) originated from industrial wastewaters.

Recently, new ecotoxicological endpoints have been introduced for many fungicides due to their secondary side effects. The majority of studies for toxicological effects on non-target organisms have been conducted on a laboratory scale, using model organisms such as *Lemna* spp., *Daphnia* spp., and *Dario* spp. [1]. *Dario rerio* is an essential organism for toxicological studies, as *Dario* is sensitive to the exposure of toxic compounds. Endocrine dysfunction, oxidative stress, and immune system disorders were observed when zebrafish were exposed to carbendazim during larval and fetal stages at concentrations above 4000 ng/L [42]. Apart from carbendazim, tebuconazole caused adverse effects on the congenital system of Zebrafish, limiting locomotion at concentrations 4 and 6 mg/L [43]. The acute and chronic toxicity of strobilurins kresoxim-methyl, pyraclostrobin, and trifloxystrobin were investigated in *D. magna* neonates and embryos by Cui et al. [44].



The results showed that *Daphnia* embryos are more sensitive to fungicide exposure than neonates, presenting 157.3 µg/L, 3.9 µg/L, and 1.7 µg/L 48-h EC<sub>50</sub> for kresoxim-methyl, pyraclostrobin, and trifloxystrobin, respectively. In addition, the lowest-observed-effect concentrations were similar to the environmental concentrations, and thus the authors reported that the studied fungicides were very toxic for *D. magna*.

There are available studies that indicate *Lemna* spp. sensitivity to fungicide exposure [45]. However, many researchers demonstrate that *Lemna* spp. can efficiently remediate contaminated natural waters and thus investigate improvement approaches. Walsh et al. [46] reported that the density over which the plants can grow on the surface of the wastewater, i.e., how much of the surface of the medium they cover, is a crucial parameter for duckweed-mediated remediation. As density increases, *L. minor* growth and TN and TP removal rates decline. To understand the remediation rate of duckweed in wetlands, total petroleum hydrocarbons in wetlands and tissues of duckweed were studied, and the experimental data were applied to the first-order kinetic rate model. After 120 days, *L. paucicostata* had successfully removed 97.91% percent of hydrocarbons from wetlands [47]. Regarding fungicides, a study showed that *L. minor* cultures could remove up to 76% of copper and 40% of dimethomorph after 96 h, suggesting *L. minor* is a suitable macrophyte plant for utilization in phytoremediation systems [48]. Many researchers have investigated the fungicides' side-effect toxicity, which can be increased when used in pesticide mixes due to synergistic interactions with other pesticides [49]. Synergy raises serious concerns for public health and environmental safety globally. Interactions between fungicides and insecticides, as well as how honeybees are affected, are of particular interest to researchers [50]. Individually, each fungicide may cause low acute toxicity to honeybees. However, a fungicide's toxicity effect could be increased when mixed with insecticides. Several studies have shown that the toxicity effects of specific pyrethroid, neonicotinoid, and organophosphate insecticides can be enhanced by combining sterol biosynthesis-inhibiting (SBI) fungicides. The combined exposure of honey bees to the fungicide propiconazole and the neonicotinoid insecticide clothianidin and a mixture containing acetamiprid and propiconazole resulted in increased mortality rates compared to exposure only to insecticides [51]. Based on their mode of action (fungicides and insecticides), similar interactions are expected to other non-target organisms. In addition, the toxicological effects of common fungicides and insecticides on honeybees were investigated during almond bloom. The insecticides chlorantraniliprole, diflubenzuron, and methoxyfenozide were tested on worker bee adults, and the fungicides iprodione and propiconazole, and a mixture of two fungicides pyraclostrobin and boscalid, were applied individually or in fungicide-insecticide mixtures to larvae. Increased larvae mortality was observed when chlorantraniliprole was applied with iprodione or propiconazole. The chlorantraniliprole-propiconazole combination was also highly toxic to adult workers. Besides the combination of chlorantraniliprole-fungicides, no synergistic effects were reported [52].

Moreover, a recent study showed that copper hydroxide is non-toxic to *Clarias gariepinus*, but mild toxicity effects were observed when it was combined with glyphosate [53]. In addition, synergistic interactions between active substances and adjuvants or inert ingredients have been reported in the literature. Pesticide commercial formulations are often thought to be more hazardous to bees than the active ingredients alone. For instance, the oral toxicity to bees of Bravo Weather Stik® (a chlorothalonil commercial formulation) showed a four-fold higher toxicity in comparison with active ingredient chlorothalonil alone [54].

Several researchers suggest that AMF symbiosis with plants in root systems has beneficial impacts on increased nutrient uptake, drought resistance, and resistance to pathogens. Similarly, plant growth-promoting rhizobacteria (PGPR) can affect plants beneficially, improving root physiology and enhancing plant growth, since they colonize the rhizosphere of plants. AMF and PGPR can be exposed (directly or indirectly) to fungicides through soil application, seed treatment, foliar spray, and wash-off from plant leaves and then drift to the soil. Thus, the physiological interactions between AMF and PGPR and host plants can be affected. Therefore, researchers focus on the combination of the advan-

tages of AMF and PGPR with a suitable chemical protection, friendly to AMF and PGPR colonies [55,56]. The impact of azoxystrobin and flutolanil (Amistar and Monarch) on the AMF *Rhizophagus irregularis* MUCL 41,833 in potatoes was studied. The applied recommended doses of azoxystrobin and flutolanil to control *Rhizoctonia solani* in potato crops and their effects against the spore germination, root colonization, extraradical mycelium development, and spore production of AMF *R. irregularis* MUCL 41,833 were assessed. Flutolanil did not cause any impact on spore germination or extra-radical development, whereas azoxystrobin significantly reduced spore germination and extra-radical development. Further, root colonization and arbuscule formation was negatively affected only by flutolanil [57]. In another study, strains of the PGPR *Pseudomonas* spp. showed tolerance when exposed to high concentrations of carbendazim and hexaconazole. The results also showed that fungicides can adversely affect the germination efficiency, growth, and physiological development of *Raphanus sativus*, but the combined application of PGPR controlled the adverse effects [56].

### 2.3. Chiral Fungicides

Many fungicides have an asymmetric center, which can provide two types of stereoisomers: enantiomers and diastereomers. Enantiomers have identical physicochemical properties but behave differently in asymmetric environments, such as in their biochemical processes. Enantiomers also show different biological activity, environmental fate, and toxicological profile. Diastereomers may not have identical physicochemical properties, and their biological activity usually varies [58].

Stereoselective fungicides differ in terms of toxicity, bioaccumulation, and bioavailability on non-target organisms [58]. Deng et al. [59] investigated the toxicity of four stereoisomers of metconazole to the aquatic algae *Chorella pyrenoidosa*. The results showed that the 96 h EC<sub>50</sub> values were different, following the pattern cis-1S,5R-Z > trans-1S,5S > trans-1R,5R > cis-1R. In addition, the photosynthesis dysfunction, the generation of reactive oxygen species (ROS), and the antioxidant response were induced more drastically by 1S,5S. In a similar study, three enantiomers of epoxiconazole were tested for their toxicological impact on the green alga *Scenedesmus obliquus*. The EC<sub>50</sub> values followed the order (+)-epoxiconazole > (−)-epoxiconazole > rac-epoxiconazole, whilst different effects on the determined chlorophyll contents, malondialdehyde contents, and antioxidant enzyme activities of algae cells were observed [60].

The environmental behavior of chiral fungicides has mainly been studied by evaluating their half-lives in crops and soils. The half-life values can provide interesting information about the potential biodegradation and the persistence of the studied compounds [54]. While fungicide enantiomers may present different half-life values from the racemic mixture, similar values were observed in other cases. The half-life of penconazole enantiomers in plant tissues and soil was determined in a field experiment. The results showed that the penconazole enantiomer (−) was degraded significantly faster than its (+) isomer in grapes and soil [61]. On the contrary, propiconazole stereoisomers were studied under aerobic, anaerobic, and sterile conditions by incubating the stereoisomers in three different types of soil, with the study investigating the dissipation process. The results showed no significant stereoselectivity under anaerobic and sterile conditions in all tested soil after 200 days of incubation, which is in contrast with the aerobic conditions where significant stereoselectivity was identified [62].

### 2.4. Nanofungicides

A fungicide is classed as nanofungicide if the size of the active ingredients ranges between 10 and 100 nm. The use of nanoparticles in fungi disease management can be divided in two categories: nanoparticles as protectants (alone) and as carriers for organic fungicides [63]. The main advantages of nanoparticles as carriers for organic fungicides are the improvement of persistence and activity of the active ingredient, the increased ability

for translocation within plants, the confrontation of the low-water-solubility problems, and the achievement of slow release.

Preventative nanofungicides can be applied directly to roots, leaves or seeds. Copper, silver, zinc oxide, and titanium dioxide are amongst the most studied nanofungicides. The effectiveness of ZnO, Ag, CuO, and Cu nanoparticles was compared with a commercial formulation containing  $\text{Cu}(\text{OH})_2$  against common fungi strains such as *Bacillus cinerea*, *Alternaria alternata*, etc. The comparison evaluated mycelial growth, colony formation, seed germination, and the hyphal and spore morphology of the fungi. Mycelial growth of fungi strains was inhibited in vitro by all the nanoparticles, but the most effective were Cu and ZnO. In addition, the nanoparticles were more lethal at the spore germination stage [64]. Shyla et al. [65] investigated the antifungal activity of zinc oxide, silver, and titanium dioxide particles against *Macrophomina phaseolina*, which can infest oilseed and pulse crops. At lower concentrations, Ag nanoparticles were more effective than ZnO and  $\text{TiO}_2$  against target fungi. In general, silver presented the highest antifungal activity from the other metals. Silver ions can cause dysfunction in thiol groups of fungal cell walls. As a result, the electron transport chain, energy metabolism, and transmembrane function are disrupted. Furthermore, fungal DNA can be mutated, respiratory chain dissociated, membrane permeability decreased, and cell lysis affected by silver-based fungicides.

Another popular nanoparticle fungicide with low toxicity risk to human health and the environment is chitosan. Chitosan can block nutrient supply, prevent the biosynthesis of mRNA and proteins, disrupt the cell membrane, and inhibit  $\text{H}^+$ -ATPase of fungi. Some of the fungi that can be managed by chitosan are Fusarium crown rot, root rot in tomato, gray mold of grapes, and rice blast disease in rice [66]. Chitosan–lactide copolymer nanoparticles were used as carriers for pyraclostrobin, a low-water-soluble fungicide. The carrier was tested against *Colletotrichum gossypii* at different concentrations and compared to a commercial formulation of pyraclostrobin. The results showed that nanofungicide effectiveness did not exceed the commercial one at three and five days post-application. Nevertheless, the nanofungicide antifungal activity was increased at day 7 post-application [67]. Janatova et al. [68] achieved a successful *Aspergillus niger* control by formulating five individual essential oils with silica material MCM-41 in nanocapsules. Their effectiveness was reported to be higher than commercial oils at 14 days post-*Aspergillus niger* infection.

Despite the advantages of nanofungicides, these compounds can enter natural waters through leaching, run-off, or spray-drift. Soil properties such as surface charge, cation species, and the type of soil can define the mobility of nanoparticles in the soil. In addition, the nanoparticles can modify the soil sorption capacity of pesticides, resulting in the fluctuations of their toxicity severity. As a result, toxicity effects have been observed in humans, plants, microorganisms, and vertebrates due to their exposure to nanofungicides [69]. According to Ameen et al. [70], the exposure of nanopesticides can have a different impact on plant growth depending on application conditions such as application rate and size and type of nanomaterial. Nanoparticles can be taken up and cross the plasma membrane through various processes, such as endocytosis, pore formation, and carrier proteins [71]. Plants can activate defense systems and overcome stress parameters (including nanoparticles). However, there is the possibility for plants to fail to overcome toxicity effects, showing symptoms such as damaged DNA, reduced rate of transpiration, and others [72]. For instance, decreased content of leaf photosynthetic pigment (chlorophyll a and b) and reduced biomass (17–20%) have been observed in maize after application of the nanofungicide  $\text{Cu}(\text{OH})_2$  [73]. In addition, nanofungicides have the potential to harm beneficial soil bacteria and fungus. Abd-alla et al. [74] reported that high concentrations of Ag-nanoparticles reduced mycorrhizal colonization, glomalin content, and mycorrhizal responsiveness of AMF *Glomus aggregatum*. Hence, the nitrogen-fixing *Azotobacter vinelandii* presented various toxicity symptoms under Ag-nanoparticle presence, such as deduced cell number, apoptosis structural damage, inhibition of biological nitrogen fixation, and ROS generation [75].



Furthermore, various aquatic organisms have been investigated for their responses to nanoparticle exposure. Mortality, hatching delays, and various developmental malformations were shown when zebrafish embryos were exposed to nanoparticles [76].

### 2.5. Chemical Plant Defense Activators

Chemical plant defense activators constitute another group of fungicides with novel mode of action. Acibenzolar s-methyl (ASM) is a compound that can induce plants' defense mechanisms, such as systemic acquired resistance, with salicylic acid taking part in the process. ASM can be used as an alternative solution to common bactericides and fungicides. Many researchers report that ASM is able to effectively manage various diseases in different crops, e.g., onion and tomato [77]. In addition, ASM induces the production of enzymes and phytoalexins when the plant is chemically, biologically, or physically stressed. However, ASM has been identified as phytotoxic, has been linked to production losses, and in some situations, may exacerbate other pest assaults [78]. Potassium phosphate (PP) is a salt that is applied in cultures as a foliar fertilizer. Plant defense function is also stimulated by PP use. PP is characterized by high mobility and solubility. As a result, harmful oomycetes are successfully controlled in different cultures, such as papaya, tomato, and potato [78,79]. Another salt, fosetyl-Al, is a worldwide broad-spectrum fungicide that is commonly used in horticulture. Its action is based on preventing the development of fungi spores and mycelium, inhibiting the pathogen penetrating into the plant. It was also documented that fosetyl-Al plays a role in plant defense mechanism activation [80].

The environmental fate and toxicological effects of chemical plant activators to non-target organisms have not been extensively studied. The effects of fosetyl-Al on model species *Danio rerio* in water and *Enchytraeus crypticus* in soil were evaluated at a laboratory scale. The ecotoxicological assessment showed that fosetyl-Al is not considered safe for *D. rerio* and *E. crypticus* for concentrations higher than the PECs, which are 1.067 mg/kg for soil and 0.06496 mg/L for surface water [81]. In the case of ASM, Guziejewski et al. [82] reported that it is moderately to highly toxic to fish, moderately toxic to invertebrates, and highly toxic to aquatic plants.

## 3. The Role of Vegetation and Microbial Communities on Fungicide Removal in CWs and Other Phytoremediation Systems

Phytoremediation is a technology that uses plants and microorganisms located in the rhizosphere in order to remove, mitigate, break down, and retain pollutants such as pesticides in soils, surface water, and groundwater. However, phytoremediation processes can be adversely affected by pesticides, which can cause phytotoxicity, as plants are exposed to stressful conditions. CWs are a type of phytoremediation system based on wetland plant species, which usually grow in soil or gravel substrate [83].

Among other factors, the vegetation and its related functions are crucial for pesticide removal in CW systems. Vegetation provides pollutant uptake, phytoaccumulation, degradation, and sorption through rhizosphere, flocculation, sedimentation, and suitable conditions for enhanced microbial activity [84]. Therefore, an efficient phytoremediation process in CWs depends on two main factors: the tolerance of plants to pollutants and the presence of favorable conditions for microorganism growth in the rhizosphere, contributing to contaminant degradation [19]. In most studies, phytoaccumulation and plant uptake are associated with  $K_{ow}$  of each compound. High or low  $\text{Log}K_{ow}$  values may facilitate uptake or translocation, respectively. However, the optimum  $\text{Log}K_{ow}$  values for uptake, translocation, and accumulation range between 3.0 and 4.0 [85]. Furthermore, researchers [18] have reported that macrophytes such as *Typha latifolia*, *Phragmites australis*, and others have high potential to absorb various pesticides, accumulating them in roots, stems, and leaves. In addition, the phytoavailability of a pesticide is determined by its molecular size (weight). A tiny molecule pesticide (Mr 500 Da) is often absorbed passively through diffusion, whereas a pesticide with a molecular weight more than 500 usually requires ATP hydrolysis to drive absorption [86].

Plants have established complex mechanisms for degrading xenobiotics like pesticides into detoxified compounds. Glycosyltransferase in plants catalyze the conjugation of sugars with endogenous metabolites or exogenous compounds. The purpose of glycosylation is to make substrates more water soluble, making them easier to degrade [86]. Many GTs have been reported to protect plants from the toxicity of fungicides by O-glycosylation. The O-glycosylation products of their hydroxylated metabolites have been found in grape and strawberry for the fungicides thiabendazole, pyrimethanil, and cyprodinil [87].

Microbial diversity and richness have a dominant role in the stability and maintenance of CW treatment efficacy. Fungicide-degrading microorganisms (beneficial for plant growth) can be found either in filter bed material (porous media) or in plant roots. In addition, epiphytic or endophytic microorganisms can colonize wetland plants. The pesticide biotransformation, through microbial communities, is mainly conducted at the rhizosphere, which is favored by the release of oxygen and organic exudates of plants [88]. The combination of phytoremediation with bioaugmentation improves treatment efficiency compared to individual approaches. Therefore, researchers have recently focused on microcosmos functions and the potential amplification of microbiota activity [89].

The contribution of vegetation and the associated microorganisms to fungicide removal through CWs and other plant-based remediation systems is described below, presenting various related studies (Table 2).

**Table 2.** The use of constructed wetlands for fungicide removal.

Target Fungicide	Vegetation	Microbial Changes/Contribution	Plant Removal	Wastewater Type	CW Type	Reference
Fluopyram	<i>Phragmites australis</i> , <i>Typha latifolia</i>	bioaugmentation with <i>Pseudomonas spp</i>	efficiency	synthetic	pilot scale HSF	[19]
Tebuconazole	<i>Juncus effuses</i> , <i>Berula erecta</i> , <i>Iris pseudacorus</i> , <i>Phragmites australis</i> , <i>Typha latifolia</i>	not specified	<i>P. australis</i> absorbed higher amount of fluopyram than <i>T.</i> <i>latifolia</i>	synthetic	pilot scale HSF, FWS	[90]
Boscalid	<i>Phragmites australis</i>	not specified	<i>B. erecta</i> achieved significantly higher removal efficiency than the other plant species	synthetic	pilot scale HSF	[15]
Chlorothalonil	<i>Phragmites australis</i>	response to fungicide exposure: non-optimal bacteria group growth	the presence of vegetation greatly enhances boscalid removal	synthetic	pilot scale HSF	[91]
Tebuconazole	<i>Juncus effuses</i> , <i>Berula erecta</i> , <i>Iris pseudacorus</i> , <i>Phragmites australis</i> , <i>Typha latifolia</i>	plants promoted higher microbial activity than unplanted CWs	non-specified	synthetic	pilot scale HSF, FWS, VFS	[92]
Tebuconazole, imazamil	<i>Juncus effuses</i> , <i>Berula erecta</i> , <i>Iris pseudacorus</i> , <i>Phragmites australis</i> , <i>Typha latifolia</i>	nitrifying bacteria may play an active role in biodegradation	<i>B. erecta</i> achieved significantly higher removal efficiency than the other plant species	synthetic	pilot scale HSF, FWS	[93]

HSF: horizontal subsurface flow; FWS: free water-surface system; VFS: vertical flow system.

### 3.1. Strobilurins

The action of strobilurins against fungal mitochondrial respiration is based on the electron transport block in the cytochrome  $bc_1$  complex (complex III), between cytochrome b and cytochrome  $c_1$ , at the  $Q_o$  site. Therefore, ATP synthesis (energy supply) is blocked, causing oxidative stress and fungus cell death [94]. Pyraclostrobin ( $\text{LogK}_{ow} = 3.99$ ) and azoxystrobin ( $\text{LogK}_{ow} = 2.50$ ) are the best sellers among strobilurin fungicides and are mainly applied to soybean, grape, wheat, and corn crops. Other commonly used strobilurin fungicides are trifloxystrobin, picoxystrobin, kresoxim-methyl, and fluaxastrobin, with  $\log K_{ow}$  values of 4.5, 3.6, 3.4, and 2.86, respectively, and their water solubility inversely proportional to  $\text{LogK}_{ow}$  values [87,95].

Maillar et al. [96] indicated that a stormwater wetland achieved 100% removal of kresoxim methyl and 93% of azoxystrobin. *P. Australis* (70–80%) was the main wetland vegetation during the monitoring period. Other plants tested were *T. latifolia* and *Choenoplectus lacustris*. Although plant uptake was not quantified, vegetation was assumed to contribute to fungicide removal due to the optimum  $\text{Log K}_{ow}$  values of the tested compounds.

In addition, the phytoaccumulation ability of three plants (*Juncus effusus*, *Pontederia cordata* and *Sagittaria latifolia*) against azoxystrobin was assessed in order to be used in wetland remediation systems. Plants were planted in sandy soil, with azoxystrobin being applied to every two months. Researchers quantified the azoxystrobin residues in plants and soil during the experiment. *P. cordata* achieved the highest removal rate (51.7%). Particularly, the root of *P. cordata* contained up to 39.8% of the total absorbed azoxystrobin [97]. Furthermore, Elsaesser et al. [98] supported that the presence of the submerged macrophyte *Elodea nuttallii* (Planch) increased the removal efficiency of a vegetated flow-through stream system in comparison with an unplanted one, achieving more than 90% removal of the initial concentration of trifloxystrobin.

The complex between cytochrome b and cytochrome  $c_1$ , which is unsettled by strobilurins, appears in all eukaryotic organisms, including plants. Thus, the exposure of CWs plants to strobilurins may cause reduced plant growth and thus decreased phytoaccumulation ability. Pedersen et al. [99] tested the response of isolated mitochondria from wheat and showed that the electron transport in the cytochrome  $bc_1$  was blocked.

### 3.2. Demethylation Inhibitors (DMI)

The DMI fungicide group consists of various chemical classes, such as triazoles and imidazoles, which have a common mode of action. DMIs inhibit the cytochrome P450, which is responsible for ergosterol production and the synthesis of the cell wall of fungi [100]. Difenconazole, tebuconazole, imazalil, epoxiconazole, and cyproconazole are considered as commonly used members of the DMI group, and their respective  $\text{LogK}_{ow}$  values are 4.36, 4.1, 3.89, 3.3, and 3.09 [84].

Lv et al. [92] studied the removal efficiency of five different CWs planted with *T. latifolia* (cattail), *P. australis* (common reed), *I. pseudacorus* (yellow flag), *J. effusus* (soft rush), and *B. erecta* (water parsnip) against imazalil and tebuconazole. Planted CWs presented much higher removal efficiency than unplanted ones during the summer period. Despite the low phytoaccumulation of fungicides, the plants contributed to fungicide biodegradation, though the active microbial community hosted in their rhizosphere. Both fungicides were mainly phytoaccumulated by *B. erecta*, while *J. effusus* has shown significantly greater plant uptake in the winter. During the experiment, the CW vegetation accumulated only 5–6% of the initial applied fungicides (nominal concentrations). Another study indicated that the tebuconazole uptake by plants was quite low, ranging between 2.1 and 12.5%. However, the role of plant metabolism in tebuconazole dissipation was remarkable [89]. Furthermore, the role of three macrophytes in difenconazole removal was investigated. Researchers suggested that the two main factors contributing to difenconazole removal were the physicochemical properties of the tested compound and the biomass of the macrophytes [101].

### 3.3. Other Chemical Fungicides

Boscalid is a relatively new succinate dehydrogenase inhibitor (SDHI), with high effectiveness and a broad spectrum. Its mode of action is based on the inhibition of fungal respiration by blocking the ubiquinone binding site [102]. Papaenvagelou et al. [15] demonstrated that the moderately lipophilic boscalid was bioaccumulated by *T. latifolia* and *P. australis*, increasing the total removal efficacy of CWs. In addition, the lower uptake ability of plants during the winter period was attributed to a lower activity of plants when compared with the summer period.

The environmental fate studies of chloronitriles are focused on the fungicide chlorothalonil. Chlorothalonil can bind and deplete cellular glutathione (GSH) and can inhibit glycolysis by binding with glyceraldehyde 3-phosphate dehydrogenase (GAPDH), causing cell death [103]. A study related to chlorothalonil removal using CWs showed that the fungicide removal was correlated with the presence of bacteria *Pseudomonas* spp., which were favored by *P. australis* [90].

As mentioned before, high copper residues in vineyard soil could have toxicological side effects. Hyperaccumulators can absorb high amounts of copper, especially in their shoots. The *Brassicaceae* family contains several plants able to accumulate significant amounts of copper. However, hyperaccumulators present low plant growth and thus low plant biomass, which is strongly correlated with copper plant uptake. Therefore, the choice of an appropriate plant is mainly based on the shoots' ability to absorb high copper amounts [104].

Anilinopyrimidine fungicides might inhibit methionine biosynthesis. In addition, several researchers have shown that anilinopyrimidine fungicides can prevent the secretion of fungal hydrolytic enzymes such as laccases, lipases, and proteases [105]. Dosnon-Olette et al. [106] tested the potential toxicity effects of pyrimethanil on five macrophytes, with their removal performance also investigated. Among the tested species, *L. minor* achieved the higher phytoaccumulation of pyrimethanil (17.1%). Moreover, *L. minor* presented high tolerance to pyrimethanil exposure. A similar study was conducted using the macrophyte species *L. minor* and *S. polyrrhiza*. The target compound was the morpholine fungicide dimethomorph. Both plants were not affected by fungicide exposure, with their removal rate reaching up to 18% [107].

Ag nanoparticles are possible carriers of fungicides, and thus their fate was investigated in CW systems. The effects of Ag nanoparticles on microbial communities in the CW substrates and the contribution to nanoparticle removal of CW vegetation were examined. The results showed that Ag nanoparticles caused changes in microbial community and also in the abundance of the key functional bacteria, decreasing the removal efficiency of the system. In addition, the plants of CW (*Iris wilsonii*) absorbed low amounts of Ag nanoparticles. However, the removal efficiency of Ag nanoparticles was evaluated at 95.72%, which indicates that the CW could effectively remove Ag nanoparticles [108]. Similarly, Cao et al. [109] investigated the removal performance of Ag nanoparticles through CWs. *I. pseudacorus* was planted in CWs and took part in the removal of Ag nanoparticles. Their residues were detected in the roots and leaves of plants. Roots could accumulate significantly higher amounts than leaves, with an overall removal performance higher than 98%.

## 4. Tools, Technologies, and Methodologies for Fungicide Phytoaccumulation Improvement

### 4.1. Plant-Growth Regulators and Phytohormones

The recent introduction of plant metabolism amplifiers, such as biofungicides and exogenous plant hormones, in the agricultural sector to control plant pathogens has given rise to speculation about their potential application in CWs to improve removal performance, as well as their interactions with CW vegetation and microbial communities. PGPR (Plant Growth-Promoting Rhizobacteria), phytohormones, chelators, and other biotechnological tools are considered as Low Risk Pesticides (LRPs) or eco-friendly disease management

approaches. In addition, under field conditions, or in various remediation techniques, PGPR, phytohormones, and chelators interact with chemical fungicides and affect their fate and removal efficacy.

Phytohormones are known as small organic molecules, which are involved in various growth and development processes of plants and also in plant immunity, supporting biotic or abiotic stress control [110]. Among the identified nine categories of phytohormones, ethylene (ET), salicylates (SA), jasmonates (JA), and cytokinins (CK) act as defense regulators in plant immunity against various pathogens, including fungi [111]. Researchers report that the exogenous application of plant hormones can enhance plant resistance against fungi pathogens [112].

Melatonin was first discovered in plants in 1995, prompting a surge of research on the hormone's role and activities in plants. Nowadays, researchers conduct studies in order to examine the impacts of melatonin in agriculture. Increase of the biomass production, improvement of seed germination, and photosynthesis activity and tolerance to biotic stress was also reported as one of the benefits of melatonin presence in plants. As a result, the exogenous application of melatonin against fungi was studied [113]. Zhang et al. [114] reported that melatonin could successfully control the infection of *Phytophthora infestans* in potato cultures. The melatonin treatment caused ultrastructure cell changes, lower pressure resistance of *P. infestans*, and an underdeveloped mycelium. Interestingly, melatonin acted synergistically with a commercial fungicide containing fluopicolide and propamocarb hydrochloride, indicating that the reduction of chemical fungicide use is possible. The combined application of classical fungicides, nanofungicides, plant growth regulators, chemical plant activators, microbial fungicides, and nutrients obfuscates the study of their environmental fate and ecotoxicity issues. The use of melatonin could enhance the fungicide removal performance of CW vegetation. Melatonin influences numerous gene expressions in different plant growth stages and physiological stages. Melatonin can act as a regulator for these plant hormones, increasing or inducing their beneficial effects on plants. Furthermore, the regulation of primary or secondary metabolism and osmosis has been reported after melatonin application. In addition, melatonin interferes with the redox network within plants, balancing both reactive nitrogen and reactive oxygen species (ROS), such as hydrogen peroxide, hypochlorous acid, and nitric oxide, which occasionally can be harmful to plants [115]. Yan et al. [116] reported that exogenous melatonin limited the phytotoxicity effects of carbendazim on tomatoes and increased plant resistance to fungicide exposure. In particular, ROS levels were decreased, antioxidant enzymatic activity was enhanced, and the induced production of glutathione contributed to fungicide detoxification.

Local and systemic acquired resistance was achieved by SA exogenous application against various fungi, such as *Fusarium oxysporum*, *A. alternata*, *Magnaporthe grisea*, and *Colletotrichum gloeosporoides* [117]. The application of 2 mM of SA was enough to reduce *B. cinerea* severity in tomato by 67% [118]. The JA application can induce plant defense, inhibit the death of plants cells, and prevent the expansion of fungi infection [119]. According to Zalewski et al. [120], a treatment of Winter Triticale with 1 mM of methyl jasmonate in specific growth stages controlled a further expansion of various triticale grain fungi pathogens, such as *A. alternata* and *Cladosporium cladosporioides*. However, a higher application dose of methyl jasmonate was correlated with higher disease severity, caused by mycotoxin-producing fungi.

Auxins are organic compounds with low molecular weight. Their presence within plants can enhance cellular and plant growth. Among them, indole acetic acid (IAA) is the most essential [121]. Auxins are demonstrated as successful contributors to phytoremediation performance. Rostami et al. [122] reported that the auxin treatments of phytoaccumulators enhances the biomass of the roots and stems. The root system of treated plants presented higher growth than other plant parts and simultaneously an increased secretion of exudates from plant root to rhizosphere. In addition, a well-developed root system provides favorable conditions for microorganism growth, as well as increased microbial



population and activity. IAA can be produced with the involvement of various bacterial species, such as *Bacillus* spp., *Acinetobacter* spp., *Pseudomonas* spp., *Azospirillum* spp., and *Azotobacter* spp. The amount of produced auxins depend on the genus and species of the bacteria and their growth conditions [123].

Gibberellins, such as gibberellic acid, are crucial for various growth and development processes of plants. Many plant species enhance the production of amine proline acid and protectant compounds. Proline can protect vital macromolecules such as proteins and increase enzymatic activity under stressful conditions. The application of exogenous gibberellins promotes proline synthesis in plant tissues, resulting in increased plant defense and a positive plant growth regulation. Apart from proline production, gibberellins can enhance the chlorophyll content and plant biomass when plants are exposed to toxic compounds [124,125].

Salicylic acid is a phenol compound that promotes plant defense response against biological or non-biological stresses and reduces the effects of toxic compounds, as a curative. It is also considered a preventive compound for the toxic effects of xenobiotics on plants. Plant physiological processes such as bud growth, membrane permeability, mitochondrial respiration, stomatal closure, material transfer, photosynthesis, growth rate, and ion absorption are affected by salicylic acid presence [126]. Salicylic acid causes metabolic reactions and thus physiological changes to plants, which are adopted by plants. These changes strengthen plant tolerance to various stressful environmental factors [127]. The exogenous salicylic acid application was tested for phytoremediation systems targeting copper. The results showed increased plant resistance to copper and a reduction of the adverse effects of copper [128].

A major function of cytokinins (CK) is to synthesize chlorophyll. The continuous chlorophyll production through CK keeps the leaves “young”, collecting free radicals and transforming etioplasts to chloroplasts. As a result, plants develop adaptations and tolerance to environmental conditions. In addition, studies showed that CK interact with plant defense systems, increasing glutathione production and eventually plant tolerance to contaminants. Moreover, CK application can enhance the plant biomass [120].

The favorable effects of plant growth regulator (PGR) treatments in the phytoremediation of heavy-metal-contaminated soil were reported by Rostami and Azhdarpoor [120]. However, the potential positive effects of PGR as an eco-friendly treatment for pesticides have not been investigated in CW vegetation. Moreover, many fungicides have a pivotal role in plant physiology and the production/regulation of phytohormones.

#### 4.2. Chelation

Many researchers report that the use of polysaccharides as chelating agents for vital elements can enhance nutrient bioavailability for plants, contributing to their uptake, translocation, and utilization within the plants. In addition, amino acid complexes with micronutrients can increase plant growth and yields through photosynthesis process improvement [129]. Nowadays, similar chelates are examined for their ability to stimulate plant immunity or for their direct effects against fungi. Buzón-Durán et al. [130] investigated the antifungal ability of chitosan oligomers (COS) and four amino acids—cysteine, glycine, proline, and tyrosine—against *Fusarium culmorum* in Spelt using field trials and in vitro seeding assays. The results showed that the COS–tyrosine mixture was the most effective in fungal growth inhibition. In field studies, the COS–tyrosine mixture reduced the disease severity by 83.5%. In addition, a mixture with chitosan and riboflavin achieved satisfactory control of *Penicillium digitatum* in in vitro experiments, promoting a potentially alternative fungicide for citrus fruit postharvest treatment, with a low environmental risk [131].

Bloem et al. [132] reported on another chelating agent, Ethylenediaminetetraacetic acid (EDTA), which is a compound mainly used in micronutrients and as a fertilizer component. The application of EDTA<sub>Na</sub> controlled *Fusarium graminearum* in wheat, by up to 90% in

field experiments. In in vitro assays, the presence of EDTANa<sub>2</sub> inhibited the deoxynivalenol production and chitin synthase activity of *Fusarium* spp. [14].

Chelators have also been studied for use in phytoremediation systems. Chelator addition (e.g., EDTA) may increase the removal of pollutants by plants, with reduced plant growth observed at the same time. In addition, these compounds have high half-life time values and are persistent in the environment; thus, the exposure of high concentrations of chelating agents to plants and soil microorganisms can be toxic [133]. Following these studies, researchers have recently focused on more eco-friendly chelating agent applications. The application of the degradable chelators glutamic acid, nitrilotriacetic acid, ethylenediamine disuccinic acid, and citric acid were examined on their own and in combination to increase the phytoextraction effectiveness of amaranth (*Amaranthus hypochondriacus* L.) in two Cd-contaminated agricultural soils. First, the plants were resistant to the chelators that were used, and the addition of glutamic acid and nitrilotriacetic acid effectively aided plant biomass production. Additionally, the combination of glutamic acid and nitrilotriacetic acid showed significantly higher soil enzyme activity compared with individual applications [134]. Despite the lack of similar studies related to fungicide removal, the use of chelators in CWs is a promising tool for enhanced plant uptake and phytoaccumulation in CWs for fungicide removal [135].

#### Bioaugmentation with PGPR Plant Growth-Promoting Rhizobacteria and AMF Arbuscular Mycorrhizal Fungi

In the field of agriculture, some isolated beneficial microorganisms, plant extracts, and essential oils can have lethal effects on crop pathogens (Table 3). Consequently, various biofungicides have been formulated and registered [9–11].

**Table 3.** List of common biofungicides [37].

Group Name and Mode of Action	Chemical/Biological Family	Organism Origin	Target Site
Plant extracts, Biologicals with multiple modes of action	Polypeptide	Extract from lupine plantlets	Multiple effects on ion membrane
	Phenols, sesquiterpenes, triterpenoids, and coumarins	Extract from <i>Swinglea glutinosa</i>	Affects fungal spores and germ tubes, induces plant defense
	Terpene hydrocarbons, terpene alcohols, and terpene phenols	Extract from <i>Melaleuca alternifolia</i>	Competition, cell membrane disruption, induced plant defense
Microbial, Biologicals with multiple modes of action	Bacterial <i>Bacillus</i> spp.	<i>Bacillus amyloliquefaciens</i> strain QST713	Competition, mycoparasitism, antibiosis, induced plant defense
	Bacterial <i>Pseudomonas</i> spp.	<i>Pseudomonas chlororaphis</i> strain AFS009	
	Fungal <i>Trichoderma</i> spp.	<i>Trichoderma atroviride</i> strain I-1237	

The application of several microbial-based fungicides is called, by many researchers, “augmentative biological control” [136]. Plant Growth Promoting Microorganisms (PGPM) are among the most popular microbial-based fungicides that can control disease directly or indirectly. Moreover, PGPM are able to increase crop yield by enhancing nutrient uptake, improving disease tolerance and enhancing abiotic or biotic stress adaptation. Plants can host PGPM in plant parts such as the rhizosphere and the endosphere. The most commonly applied microbial inoculants are the PGPR (e.g., *Pseudomonas* spp., *Bacillus* spp., and *Rhizobia* spp.), followed by the AMF (e.g., *Glomus* spp. and *Rhizoglyphus* spp.). Numerous studies reported successful applications of PGPM against plant diseases. *Bacillus amyloliquefaciens* (former subtilis) is a rhizobacterium that can produce endospores and

antibiotics. The mode of action is based on pathogen cell membrane disruption, and it is mainly used for *B. cinerea* on horticultural species [137]. Ongena et al. [138] reported that *Bacillus amyloliquefaciens* has the potential to induce plant defense mechanisms of disease-affected plants.

PGPR promote the production of vital compounds, including organic acids, siderophores, antibiotics, enzymes, and phytohormones. These compounds are responsible for various beneficial impacts on the vegetation of phytoremediation systems, which improve pollutant removal efficiency, including greater plant biomass, tolerance development under abiotic (e.g., weather conditions) or biotic (e.g., pathogens) stressful conditions, higher nutrient uptake, higher microbial activity on the root system, increase of pesticide biodegradation, and pesticide bioaccumulation [139–142]. In addition, PGPR can reduce ethylene production through enzyme 1-aminocyclopropane-1-carboxylate (ACC) deaminase synthesis, which dissipates the ethylene precursor ACC, achieving a significant plant growth enhancement [143]. Furthermore, DalCorso et al. [144] suggests that plant growth promotion and phytoaccumulation improvement can be achieved by bacterial auxin (IAA), which is stimulated by PGPR. Despite the lack of studies related to acibenzolar-S-methyl fate in phytoremediation systems, some studies prove its translocation ability within plant tissues and its interactions with PGPR strains. For instance, the PGPR strains *Bacillus subtilis* GB03 and *Bacillus pumilus* SE34 had a significant influence on ASM translocation in tomatoes, promoting higher concentrations of fungicide in aerial parts compared with non-PGPR treated plants [145].

Xiao et al. [146] investigated the remediation ability of hyperaccumulator *Sedum alfredii* to remove the benzimidazole fungicide carbendazim. The plantation of *S. alfredii* was combined with carbendazim-degrading bacterial strains (*Bacillus subtilis*, *Paracoccus* spp., *Flavobacterium* spp., and *Pseudomonas* spp.). The removal performance was increased by 32.1–42.5% when inoculated with bacteria *S. alfredii* treatments. Additionally, bacterial inoculation improved the microbial activity and structure of soil and enhanced carbendazim biodegradation. Parlakidis et al. [19] investigated the influence of bioaugmentation of PGPR *Pseudomonas putida* on *P. australis* growth and uptake ability in CWs, targeting the fungicide fluopyram. The results showed that the inoculated plants absorbed higher concentrations of fluopyram, and they had higher plant biomass compared with the non-bioaugmented plants.

Furthermore, PGPR can be directly involved in fungicide biodegradation. The influence of *Bacillus subtilis* on the fungicide penthiopyrad was investigated on a laboratory scale. *B. subtilis* can enhance by approximately 5% the dissipation rate of penthiopyrad. However, its combination with fungi *Trichoderma harzianum* increased the degradation by 29.1% [147]. Myresiotis et al. [148] studied the interactions between various PGPR strains inoculated in soil and fungicides propamocarb hydrochloride and acibenzolar-S-methyl. In all cases, soil inoculated with PGPR showed a remarkable dissipation rate of acibenzolar-S-methyl, approximately six times higher than non-inoculated soils. From all tested strains, *Bacillus amyloliquefaciens* strain IN937a and *Bacillus Pumilus* strain SE34 had the most significant impact on propamocarb hydrochloride degradation.

Another essential microbe group that can be bioaugmented in CW plants is the AMF. AMF presents similar benefits to PGPR. The increased production of phytohormones, which enhance plant growth and contaminant plant uptake, was associated with AMF inoculation in accordance with Liao et al. [149]. Furthermore, Göhre and Paszkowski [150] demonstrated that plant rhizosphere growth and the nutrients and water adsorption by plant roots was much greater in inoculated AMF plants as compared to untreated plants.

## 5. Conclusions, Perspectives, and Challenges

Fungicides are considered a vital tool for agriculture, protecting crops against fungal diseases and therefore securing high agricultural productivity. Nowadays, the evolution of technology can provide novel chemical fungicides, such as nanofungicides and chiral fungi-

cides, to address plant resistance development. However, the extensive use of chemical fungicides leads to a risk for public health, natural waters, and non-target organisms.

Various studies indicate that CWs can prevent the distribution of pesticides to the natural environment, but few of them target fungicides. Fungicides can have adverse impacts on plants and beneficial microorganisms in CWs. As a result, plant accumulators and microbial communities may have a low contribution to fungicide removal. In the agricultural industry, microbial formulations (e.g., PGPR and AMF) are proposed as an alternative eco-friendly solution for plant diseases. Similarly, synthetic plant growth regulators, chelators, and phytohormones have a beneficial effect on plants under biotic or abiotic stress, showing much promise as an effective anti-fungal strategy. Their mode of action is based on plant defense induction, enhanced plant biomass, and production of vital compounds. Recently, these formulations have been investigated in phytoremediation systems, presenting effective removal of organic pollutants, metals, and nutrients. This review demonstrates strong evidence to suggest that the addition of exogenous microorganisms, plant growth regulators, and phytohormones can overcome the fungicide toxicity side-effects for CW vegetation and microbial communities and enhance fungicide removal. However, there are information gaps regarding their interactions with chemical fungicides within CWs.

The exogenous application of PGRs can be used to improve phytoremediation, but there is still a lack of information on the appropriate application stages and dosages. Hence, the synergistic effects of indigenous and exogenous melatonin on toxic compounds are still unclear. The application of chelating agents in CWs could be an efficient strategy to remove inorganic fungicides and nanofungicides. However, low-biodegradable chelators at high doses can have a toxic effect on CW vegetation and microbial communities. Therefore, the choice of plant species, chelator type, and dose treatment are critical for optimum removal performance.

The microbial populations in the substrate or water of the CW systems, as well as those in the plant material, could be altered by inoculation. Many studies found that the inoculation strains were persistent (e.g., [151]), while some investigations found that the density of the inoculated strains decreased over time (e.g., [152]). In addition, the length of the study appears to have an effect on the outcomes. The inoculation process can be done immediately, or waiting for the system to be stabilized [89]. Hussain et al. [151] found that removal performance of the tested bioaugmented CWs was optimum in the first three months, but then deteriorated in the following months. Plant health declined at the same time, with plant shoots beginning to become yellowish.

However, available data for the microbial inoculation effects on microbial communities in CW system are provided mostly by molecular fingerprinting such as restriction fragment, length polymorphism, real-time quantitative PCR, etc., which indicate only the dominant members of the microbial communities. Next-generation sequencing techniques are necessary to determine the population size of inoculants, their competition with indigenous microorganisms, and their interactions with the fungicides, reaching taxa identification. Furthermore, the hydrological conditions of CWs have received little attention, which could have a crucial impact on microbial development, considering that free water surface CWs are mostly aerobic, whereas HF CWs are mostly anoxic/anaerobic.

**Author Contributions:** Conceptualization, G.D.G. and Z.V.; perspectives, G.D.G. and Z.V.; writing—original draft preparation, P.P. and T.M.; writing—review and editing, Z.V. and G.D.G.; supervision, Z.V. and G.D.G. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was supported by the Hellenic Foundation for Research and Innovation (HFRI) under the HFRI PhD Fellowship grant (Fellowship Number: 353).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.



**Data Availability Statement:** The data supporting the research findings of this study are available from the corresponding author upon request.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Zubrod, J.P.; Bundschuh, M.; Arts, G.; Bruhl, C.A.; Imfeld, G.; Knabel, A.; Payraudeau, S.; Rasmussen, J.J.; Rohr, J.; Scharmuller, A.; et al. Fungicides: An overlooked pesticide class? *Environ. Sci. Technol.* **2019**, *53*, 3347–3365. [CrossRef] [PubMed]
2. Sharma, A.; Kumar, V.; Shahzad, B.; Tanveer, M. Worldwide Pesticide Usage and its Impacts on Ecosystem. *SN Appl. Sci.* **2019**, *1*, 1446. [CrossRef]
3. EUROSTAT. 2021. Available online: [https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Agrienvironmental\\_indicator\\_consumption\\_of\\_pesticides](https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Agrienvironmental_indicator_consumption_of_pesticides) (accessed on 9 September 2021).
4. Nicolopoulou-Stamati, P.; Maipas, S.; Kotampasi, C.; Stamatis, P.; Hens, L. Chemical Pesticides and Human Health: The Urgent Need for a New Concept in Agriculture. *Front. Public Health* **2016**, *4*, 148. [CrossRef] [PubMed]
5. Siddhartha; Verma, A.; Bashyal, B.M.; Gogoi, R.; Kumar, R. New nano-fungicides for the management of sheath blight disease (*Rhizoctonia solani*) in rice. *Int. J. Pest Manag.* **2020**, 1–10. [CrossRef]
6. Duhan, J.S.; Kumar, R.; Kumar, N.; Kaur, P.; Nehra, K.; Duhan, S. Nanotechnology: The new perspective in precision agriculture. *Biotechnol. Rep.* **2017**, *24*, 11–23. [CrossRef]
7. Jeschke, P. Current status of chirality in agrochemicals. *Pest Manag. Sci.* **2018**, *74*, 2389–2404. [CrossRef]
8. Zhou, M.; Wang, W. Recent Advances in Synthetic Chemical Inducers of Plant Immunity. *Front. Plant Sci.* **2018**, *9*, 1613. [CrossRef]
9. Sheir, S.K.; Osman, G.Y.; Galal, M.A.; Soliman, M.M. Biohazards of the biofungicide, *Trichoderma harzianum* on the crayfish, *Procambarus clarkii*: Histological and biochemical implications. *Egypt. J. Basic Appl. Sci.* **2015**, *2*, 87–97. [CrossRef]
10. Abbeya, J.A.; Percivala, D.; Abbeya, L.; Asiedua, S.K.; Prithiviraja, B.; Schilder, A. Biofungicides as alternative to synthetic fungicide control of grey mould (*Botrytis cinerea*)—Prospects and challenges. *Biocontrol Sci. Technol.* **2019**, *29*, 207–228. [CrossRef]
11. Brown, M.S.; Baysal-Gurel, F.; Oliver, J.B.; Addesso, K.M. Comparative Performance of Fungicides, Biofungicides, and Host Plant Defense Inducers in Suppression of Phytophthora Root Rot in Flowering Dogwood during Simulated Root Flooding Events. *Plant Dis.* **2019**, *103*, 1703–1711. [CrossRef]
12. Legein, M.; Smets, W.; Vandenheuvel, D.; Eilers, T.; Muysshondt, B.; Prinsen, E.; Samson, R.; Lebeer, S. Modes of action of microbial biocontrol in the phyllosphere. *Front. Microbiol.* **2020**, *11*, 1619. [CrossRef] [PubMed]
13. Kaewchangwat, N.; Dueansawang, S.; Tumcharern, G.; Suttisintong, K. Synthesis of Copper-Chelates Derived from Amino Acids and Evaluation of Their Efficacy as Copper Source and Growth Stimulator for *Lactuca sativa* in Nutrient Solution Culture. *J. Agric. Food Chem.* **2017**, *65*, 9828–9837. [CrossRef] [PubMed]
14. Song, X.S.; Gu, K.X.; Gao, J.; Wang, J.X.; Ding, S.C.; Zhou, M. Ethylenediaminetetraacetic Acid Disodium Salt Acts as an Antifungal Candidate Molecule against *Fusarium graminearum* by Inhibiting DON Biosynthesis and Chitin Synthase Activity. *Toxins* **2020**, *13*, 17. [CrossRef] [PubMed]
15. Papaevangelou, V.A.; Gikas, G.D.; Vryzas, Z.; Tsihrintzis, V.A. Treatment of agricultural equipment rinsing water containing a fungicide in pilot-scale horizontal subsurface flow constructed wetlands. *Ecol. Eng.* **2017**, *101*, 193–200. [CrossRef]
16. Gikas, G.D.; Pérez-Villanueva, M.; Tsiaras, M.; Alexoudis, C.; Pérez-Rojas, G.; Masís-Mora, M.; Lizano-Fallas, V.; Rodríguez-Rodríguez, C.E.; Vryzas, Z.; Tsihrintzis, V.A. Low-cost approaches for the removal of terbutylazine from agricultural wastewater: Constructed wetlands and biopurification system. *Chem. Eng. J.* **2018**, *335*, 647–656. [CrossRef]
17. Rajmohan, K.S.; Chandrasekaran, R.; Varjani, S. A Review on Occurrence of Pesticides in Environment and Current Technologies for Their Remediation and Management Indian. *J. Microbiol.* **2020**, *60*, 125–138.
18. Vymazal, J.; Březinová, T. The use of constructed wetlands for removal of pesticides from agricultural runoff and drainage: A review. *Environ. Int.* **2015**, *75*, 11–20. [CrossRef]
19. Parlakidis, P.; Mavropoulos, T.; Vryzas, Z.; Gikas, G.D. Fluopyram removal from agricultural equipment rinsing water using HSF pilot-scale constructed wetlands. *Environ. Sci. Pollut. Res.* **2021**, in press. [CrossRef]
20. Gikas, G.D.; Vryzas, Z.; Tsihrintzis, V.A. S-metolachlor herbicide removal in pilot-scale horizontal subsurface flow constructed wetlands. *Chem. Eng. J.* **2018**, *339*, 108–116. [CrossRef]
21. Kiaune, L.; Singhasemanon, N. Pesticidal copper (I) oxide: Environmental fate and aquatic toxicity. *Rev. Environ. Contam. Toxicol.* **2011**, *213*, 1–26.
22. Georgopoulos, P.G.; Roy, A.; Yonone-Lioy, M.J.; Opiekun, R.E.; Lioy, P.J. Copper: Environmental dynamics and human exposure issues. *J. Toxicol. Environ. Health Part B* **2001**, *4*, 341–394. [CrossRef] [PubMed]
23. Rader, K.J.; Carbonaro, R.F.; van Hullebusch, E.D.; Baken, S.; Delbeke, K. The Fate of Copper Added to Surface Water: Field, Laboratory, and Modeling Studies. *Environ. Toxicol. Chem.* **2019**, *7*, 1386–1399. [CrossRef] [PubMed]
24. Beck, A.J.; Sañudo-Wilhelmy, S.A. Impact of water temperature and dissolved oxygen on copper cycling in an urban estuary. *Environ. Sci. Technol.* **2007**, *41*, 6103–6108. [CrossRef] [PubMed]
25. U.S. EPA. ECOTOX Database. 2009. Available online: <http://cfpub.epa.gov/ecotox/> (accessed on 10 September 2021).



26. Ballabio, C.; Panagos, P.; Lugato, E.; Huang, J.H.; Orgiazzi, A.; Jones, A.; Fernández-Ugalde, O.; Borrelli, P.; Montanarella, L. Copper distribution in European topsoils: An assessment based on LUCAS soil survey. *Sci. Total Environ.* **2018**, *636*, 282–298. [CrossRef]
27. Lamichhane, J.R.; Osdaghi, E.; Behlau, F.; Kohl, J.; Jones, J.B.; Noel, J. Thirteen decades of antimicrobial copper compounds applied in agriculture. A review. *Agron. Sustain. Dev.* **2018**, *38*, 28. [CrossRef]
28. Kuklińska, K.; Wolska, L.; Namieśnik, J.; Monika Cieszyńska, M. Analytical and bioanalytical problems associated with the toxicity of elemental sulfur in the environment. *Trends. Analyt. Chem.* **2013**, *48*, 14–21. [CrossRef]
29. Díaz-Raviña, M.; Calvo de Anta, R.; Bååth, E. Tolerance (PICT) of the bacterial communities to copper in vineyards soils from Spain. *J. Environ. Qual.* **2007**, *36*, 1760–1764. [CrossRef]
30. Ambrosini, V.G.; Voges, J.G.; Canton, L.; da Rosa Couto, R.; Ferreira, P.A.A.; Comin, J.J.; Soares, C.R.F.S. Effect of arbuscular mycorrhizal fungi on young vines in copper-contaminated soil. *Braz. J. Microbiol.* **2015**, *46*, 1045–1052. [CrossRef]
31. Schoffer, J.T.; Sauvé, S.; Neaman, A.; Ginocchio, R. Role of Leaf Litter on the Incorporation of Copper-Containing Pesticides into Soils Under Fruit Production: A Review. *J. Soil Sci. Plant Nutr.* **2020**, *20*, 990–1000. [CrossRef]
32. Cetkauskaitė, A.; Pessala, P.; Södergren, A. Elemental sulfur: Toxicity in vivo and in vitro to bacterial luciferase, in vitro yeast alcohol dehydrogenase, and bovine liver catalase. *Environ. Toxicol.* **2004**, *19*, 372–386. [CrossRef]
33. Huber, B.; Herzog, B.; Drewes, J.E.; Koch, K.; Müller, E. Characterization of sulfur oxidizing bacteria related to biogenic sulfuric acid corrosion in sludge digesters. *BMC Microbiol.* **2016**, *16*, 153. [CrossRef]
34. Papadakis, E.N.; Vryzas, Z.; Kotopoulou, A.; Kintzikoglou, K.; Makris, K.C.; Papadopoulou-Mourkidou, E. A Pesticide monitoring survey in rivers and lakes of northern Greece and its human and ecotoxicological risk assessment. *Ecotoxicol. Environ. Saf.* **2018**, *116*, 1–9. [CrossRef] [PubMed]
35. Marsala, R.Z.; Capri, E.; Russo, E.; Bisagni, M.; Colla, R.; Lucini, L.; Gallo, A.; Suci, N.A. First evaluation of pesticides occurrence in groundwater of Tidone Valley, an area with intensive viticulture. *Sci. Total Environ.* **2020**, *736*, 139730. [CrossRef] [PubMed]
36. PPDB (Pesticide Properties Data Base). *Pesticides General Information*; University of Hertfordshire: Hatfield, UK, 2016; Available online: <https://sitem.herts.ac.uk/aeru/ppdb/en/Reports/1362.htm> (accessed on 1 July 2021).
37. FRAC Code List 2021: Fungal Control Agents Sorted by Cross Resistance Pattern and Mode of Action. Available online: [https://www.frac.info/docs/default-source/publications/frac-code-list/fraccode-list-2021--final.pdf?sfvrsn=f7ec499a\\_2](https://www.frac.info/docs/default-source/publications/frac-code-list/fraccode-list-2021--final.pdf?sfvrsn=f7ec499a_2) (accessed on 15 October 2021).
38. Komárek, M.; Čadková, E.; Chrástný, V.; Bordas, F.; Bollinger, J.C. Contamination of vineyard soils with fungicides: A review of environmental and toxicological aspects. *Environ. Int.* **2010**, *36*, 138–151. [CrossRef] [PubMed]
39. Herrero-Hernández, E.; Andrades, M.S.; Álvarez-Martín, A.; Pose-Juan, E.; Rodríguez-Cruz, M.S.; Sánchez-Martín, M.J. Occurrence of pesticides and some of their degradation products in waters in a Spanish wine region. *J. Hydrol.* **2013**, *486*, 234–245. [CrossRef]
40. Wightwick, A.M.; Bui, A.D.; Zhang, P.; Rose, G.; Allinson, M.; Myers, J.H.; Reichman, S.M.; Menzies, N.W.; Pettigrove, V.; Allinson, G. Environmental fate of fungicides in surface waters of a horticultural-production catchment in southeastern Australia. *Arch. Environ. Contam. Toxicol.* **2012**, *62*, 380–390. [CrossRef]
41. Merel, S.; Benzing, S.; Gleiser, C.; Di Napoli-Davis, G.; Zwiener, C. Occurrence and overlooked sources of the biocide carbendazim in wastewater and surface water. *Environ. Pollut.* **2018**, *239*, 512–521. [CrossRef]
42. Jiang, J.; Wu, S.; Wang, Y.; An, X.; Cai, L.; Zhao, X.; Wu, C. Carbendazim has the potential to induce oxidative stress, apoptosis, immunotoxicity and endocrine disruption during zebrafish larvae development. *Toxicol. Vitro* **2015**, *29*, 1473–1481. [CrossRef]
43. Altenhofen, S.; Nabinger, D.D.; Wiprich, M.T.; Pereira, T.C.B.; Bogo, M.R.; Bonan, C.D. Tebuconazole alters morphological, behavioral and neurochemical parameters in larvae and adult zebrafish (*Danio rerio*). *Chemosphere* **2017**, *180*, 483–490. [CrossRef]
44. Cui, F.; Chai, T.; Liu, X.; Wang, C. Toxicity of three strobilurins (kresoxim-methyl, pyraclostrobin, and trifloxystrobin) on *Daphnia magna*. *Environ. Toxicol. Chem.* **2017**, *36*, 182–189. [CrossRef]
45. Giddings, J.M.; Arts, G.; Hommen, U. The relative sensitivity of macrophyte and algal species to herbicides and fungicides: An analysis using species sensitivity distributions. *Integr. Environ. Assess. Manag.* **2013**, *9*, 308–318. [CrossRef] [PubMed]
46. Walsh, É.; Coughlan, N.E.; O'Brien, S.; Jansen, M.; Kuehnhold, H. Density Dependence Influences the Efficacy of Wastewater Remediation by *Lemna minor*. *Plants* **2021**, *10*, 1366. [CrossRef] [PubMed]
47. Ekperusi, A.O.; Nwachukwu, E.O.; Sikoki, F.D. Assessing and Modelling the Efficacy of *Lemna paucicostata* for the Phytoremediation of Petroleum Hydrocarbons in Crude Oil-Contaminated Wetlands. *Sci. Rep.* **2020**, *10*, 8489. [CrossRef] [PubMed]
48. Megateli, S.; Dosnon-Olette, R.; Trotel-Aziz, P.; Geffrad, A.; Semsari, S.; Couderchet, M. Simultaneous effects of two fungicides (copper and dimethomorph) on their phytoremediation using *Lemna minor*. *Ecotoxicology* **2013**, *22*, 683–692. [CrossRef] [PubMed]
49. Cedergreen, N. Quantifying Synergy: A Systematic Review of Mixture Toxicity Studies within Environmental Toxicology. *PLoS ONE* **2014**, *9*, e96580. [CrossRef]
50. Sgolastra, F.; Arnan, X.; Cabbri, R.; Isani, G.; Medrzycki, P.; Teper, D.; Bosch, J. Combined exposure to sublethal concentrations of an insecticide and a fungicide affect feeding, ovary development and longevity in a solitary bee. *Proc. Biol. Sci.* **2018**, *285*, 20180887. [CrossRef]
51. Han, W.; Yang, Y.; Gao, J.; Zhao, D.; Ren, C.; Wang, S.; Zhao, S.; Zhong, Y. Chronic toxicity and biochemical response of *Apis cerana cerana* (Hymenoptera: Apidae) exposed to acetamiprid and propiconazole alone or combined. *Ecotoxicology* **2019**, *28*, 399–411. [CrossRef]

52. Wade, A.; Lin, C.H.; Kurkul, C.; Regan, E.R.; Johnson, R.M. Combined Toxicity of Insecticides and Fungicides Applied to California Almond Orchards to Honey Bee Larvae and Adults. *Insects* **2019**, *10*, 20. [\[CrossRef\]](#)
53. Kingsley, K.C. Acute toxicity of copper hydroxide and glyphosate mixture in *Clarias gariepinus*: Interaction and prediction using mixture assessment models. *Environ. Health Toxicol.* **2019**, *34*, e2019003. [\[CrossRef\]](#)
54. Mullin, C.A.; Chen, J.; Fine, J.D.; Frazier, M.T.; Frazier, J.L. The formulation makes the honey bee poison. *Pestic. Biochem. Physiol.* **2015**, *120*, 27–35. [\[CrossRef\]](#)
55. Hage-Ahmed, K.; Rosner, K.; Steinkellner, S. Arbuscular mycorrhizal fungi and their response to pesticides. *Pest Manag. Sci.* **2019**, *75*, 583–590. [\[CrossRef\]](#) [\[PubMed\]](#)
56. Khan, S.; Shahid, M.; Khan, M.S.; Syed, A.; Bahkali, A.H.; Elgorban, A.M.; Pichtel, J. Fungicide-Tolerant Plant Growth-Promoting Rhizobacteria Mitigate Physiological Disruption of White Radish Caused by Fungicides Used in the Field Cultivation. *Int. J. Environ. Res. Public Health* **2020**, *17*, 7251. [\[CrossRef\]](#) [\[PubMed\]](#)
57. Buysens, C.; Dupré de Boulois, H.; Declerck, S. Do fungicides used to control *Rhizoctonia solani* impact the non-target arbuscular mycorrhizal fungus *Rhizophagus irregularis*? *Mycorrhiza* **2015**, *25*, 277–288. [\[CrossRef\]](#) [\[PubMed\]](#)
58. Bielská, L.; Hale, S.E.; Škulcová, L. A review on the stereospecific fate and effects of chiral conazole fungicides. *Sci. Total Environ.* **2021**, *750*, 141600. [\[CrossRef\]](#)
59. Deng, Y.; Zhang, W.; Qin, Y.; Liu, R.; Zhang, L.; Wang, Z.; Zhou, Z.; Diao, J. Stereoselective toxicity of metconazole to the antioxidant defenses and the photosynthesis system of *Chlorella pyrenoidosa*. *Aquat. Toxicol.* **2019**, *210*, 129–138. [\[CrossRef\]](#)
60. Liu, C.; Wang, B.; Diao, J.; Zhou, Z. Enantioselective toxicity and bioaccumulation of epoxiconazole enantiomers to the green alga *Scenedesmus obliquus*. *RSC Adv.* **2016**, *6*, 59842–59850. [\[CrossRef\]](#)
61. Zhang, X.; Wang, X.; Luo, F.; Sheng, H.; Zhou, L.; Zhong, Q.; Lou, Z.; Sun, H.; Yang, M.; Cui, X.; et al. Application and enantioselective residue determination of chiral pesticide penconazole in grape, tea, aquatic vegetables and soil by ultraperformance liquid chromatography-tandem mass spectrometry. *Ecotoxicol. Environ. Saf.* **2019**, *15*, 530–537. [\[CrossRef\]](#)
62. Pan, X.; Cheng, Y.; Dong, F.; Liu, N.; Xu, J.; Liu, X.; Wu, X.; Zheng, Y. Stereoselective bioactivity, acute toxicity and dissipation in typical paddy soils of the chiral fungicide propiconazole. *J. Hazard. Mater.* **2018**, *359*, 194–202. [\[CrossRef\]](#)
63. Worrall, E.A.; Hamid, A.; Mody, K.T.; Mitter, N.; Pappu, H.R. Nanotechnology for Plant Disease Management. *Agronomy* **2018**, *8*, 285. [\[CrossRef\]](#)
64. Malandrakis, A.A.; Kavroulakis, N.; Chrysikopoulos, C.V. Use of copper, silver and zinc nanoparticles against foliar and soil-borne plant pathogens. *Sci. Total Environ.* **2019**, *670*, 292–299. [\[CrossRef\]](#)
65. Shyla, K.; Natarajan, N.; Nakkeeran, S. Antifungal Activity of Zinc Oxide, Silver and Titanium Dioxide Nanoparticles against *Macrophomina phaseolina*. *J. Mycol. Plant Pathol.* **2014**, *44*, 269–274.
66. Safari, Z.S.; Ding, P.; Nakasha, J.J.; Yusoff, S.F. Controlling *Fusarium oxysporum* Tomato Fruit Rot under Tropical Condition Using Both Chitosan and Vanillin. *Coatings* **2021**, *11*, 367. [\[CrossRef\]](#)
67. Xu, L.; Cao, L.; Li, F.; Wang, X.; Huang, Q. Utilization of chitosan-lactide copolymer nanoparticles as controlled release pesticide carrier for pyraclostrobin against *Colletotrichum gossypii* Southw. *J. Dispers. Sci. Technol.* **2014**, *35*, 544–550. [\[CrossRef\]](#)
68. Janatova, A.; Bernardos, A.; Smid, J.; Frankova, A.; Lhotka, M.; Kourimská, L.; Pulkrabek, J.; Kloucek, P. Long-term antifungal activity of volatile essential oil components released from mesoporous silica materials. *Ind. Crop. Prod.* **2015**, *67*, 216–220. [\[CrossRef\]](#)
69. Chaud, M.; Souto, E.B.; Zielinska, A.; Severino, P.; Batain, F.; Oliveira-Junior, J.; Alves, T. Nanopesticides in Agriculture: Benefits and Challenge in Agricultural Productivity, Toxicological Risks to Human Health and Environment. *Toxics* **2021**, *9*, 131. [\[CrossRef\]](#) [\[PubMed\]](#)
70. Ameen, F.; Alsamhary, K.; Alabdullatif, J.A.; AlNadhari, S. A review on metal-based nanoparticles and their toxicity to beneficial soil bacteria and fungi. *Ecotoxicol. Environ. Saf.* **2021**, *213*, 112027. [\[CrossRef\]](#)
71. Goswami, P.; Yadav, S.; Mathur, J. Positive and negative effects of nanoparticles on plants and their applications in agriculture. *Plant Sci. Today* **2019**, *6*, 232–234. [\[CrossRef\]](#)
72. Grillo, R.; Fraceto, L.F.; Amorim, M.J.B.; Scott-Fordsmand, J.J.; Schoonjans, R.; Chaudhry, Q. Ecotoxicological and regulatory aspects of environmental sustainability of nanopesticides. *J. Hazard Mater.* **2021**, *404*, 124148. [\[CrossRef\]](#)
73. Zhao, L.; Hu, Q.; Huang, Y.; Keller, A.A. Response at Genetic, Metabolic, and Physiological Levels of Maize (*Zea mays*) Exposed to a Cu(OH)<sub>2</sub> Nanopesticide. *ACS Sustain. Chem. Eng.* **2017**, *5*, 8294–8301. [\[CrossRef\]](#)
74. Abd-Alla, M.H.; Nafady, N.A.; Khalaf, D.M. Assessment of silver nanoparticles contamination on faba bean *Rhizobium leguminosarum* bv. *Viciae*, *Glomus aggregatum* symbiosis: Implications for induction of autophagy process in root nodule. *Agric. Ecosyst. Environ.* **2016**, *218*, 163–177. [\[CrossRef\]](#)
75. Zhang, L.; Wu, L.; Si, Y.; Shu, K. Size-dependent cytotoxicity of silver nanoparticles to *Azotobacter vinelandii*: Growth inhibition, cell injury, oxidative stress and internalization. *PLoS ONE* **2018**, *13*, e0209020. [\[CrossRef\]](#) [\[PubMed\]](#)
76. Jennings, V.; Goodhead, R.; Tyler, C.R. Ecotoxicology of Nanomaterials in Aquatic Systems. *Front. Nanosci.* **2015**, *8*, 3–45.
77. Sun, H.; Luo, F.; Zhang, X.; Zhou, L.; Lou, Z.; Chen, Z. Residue analysis and dietary exposure risk assessment of acibenzolar-S-methyl and its metabolite acibenzolar acid in potato, garlic, cabbage, grape and tomato. *Ecotoxicol. Environ. Saf.* **2021**, *207*, 111178. [\[CrossRef\]](#) [\[PubMed\]](#)
78. Oliveira, T.A.S.; Blum, L.E.B.; Duarte, E.A.A.; Luz, E.D.M.N. Reduction of Papaya Rot *Phytophthora palmivora* with Phosnite and Acibenzolar-s-methyl in Preharvest and Postharvest. *J. Biosci.* **2018**, *34*, 1522–1531. [\[CrossRef\]](#)

79. Ganiyu, S.A.; Popoola, A.R.; Agbolade, J.O.; Enikuomehim, O.A. Thymol and acibenzolar-s-methyl reduce incidence and severity of bacterial wilt of tomato caused by race I biovar III (R1B3) strain of *Ralstonia solanacearum* in Nigeria. *Arch. Phytopathol. Plant Prot.* **2018**, *51*, 1106–1129. [\[CrossRef\]](#)
80. Buiarelli, F.; Di Filippo, P.; Riccardi, C.; Pomata, D.; Marsiglia, R.; Console, C.; Puri, D. Hydrophilic Interaction Liquid Chromatography-Tandem Mass Spectrometry Analysis of Fosetyl-Aluminum in Airborne Particulate Matter. *J. Anal. Methods Chem.* **2018**, *7*, 8792085. [\[CrossRef\]](#)
81. Barreto, A.; Santos, J.; Amorim, M.J.B.; Maria, V.L. Is the Synthetic Fungicide Fosetyl-Al Safe for the Ecotoxicological Models *Danio rerio* and *Enchytraeus crypticus*? *Appl. Sci.* **2021**, *11*, 7209. [\[CrossRef\]](#)
82. Guziejewski, D.; Brycht, M.; Nosal-Wiercińska, A.; Smarzewska, S.; Ciesielski, W.; Skrzypek, S. Electrochemical study of the fungicide acibenzolar-s-methyl and its voltammetric determination in environmental samples. *J. Environ. Sci. Health B* **2014**, *49*, 550–556. [\[CrossRef\]](#)
83. Carvalho, P.N.; Basto, M.C.P.; Almeida, C.M.R.; Brix, H. A review of plant–pharmaceutical interactions: From uptake and effects in crop plants to phytoremediation in constructed wetlands. *Environ. Sci. Pollut. Res.* **2014**, *21*, 11729–11763. [\[CrossRef\]](#)
84. Guo, W.; Zhang, H.; Huo, H. Organochlorine pesticides in aquatic hydrophyte tissues and surrounding sediments in Baiyangdian wetland, China. *Ecol. Eng.* **2014**, *67*, 150–155. [\[CrossRef\]](#)
85. Vryzas, Z. The Plant as Metaorganism and Research on Next-Generation Systemic Pesticides—Prospects and Challenges. *Front. Microbiol.* **2016**, *7*, 1968. [\[CrossRef\]](#) [\[PubMed\]](#)
86. Bauer, A.; Luetjohann, J.; Rohn, S.; Jantzen, E.; Kuballa, J. Development of a suspect screening strategy for pesticide metabolites in fruit and vegetables by UPLC–Q–TOF–MS. *Food Anal. Methods* **2018**, *11*, 1591–1607. [\[CrossRef\]](#)
87. Zhang, C.; Zhou, T.; Xu, Y.; Du, Z.; Li, B.; Wang, J.; Wang, J.; Zhu, L. Ecotoxicology of strobilurin fungicides. *Sci. Total Environ.* **2020**, *742*, 140611. [\[CrossRef\]](#) [\[PubMed\]](#)
88. Sánchez, O. Constructed Wetlands Revisited: Microbial Diversity in the–Omics Era. *Microb. Ecol.* **2017**, *73*, 722–733. [\[CrossRef\]](#) [\[PubMed\]](#)
89. Tondera, K.; Chazarenc, F.; Chagnon, P.L.; Brisson, J. Bioaugmentation of treatment wetlands—A review. *Sci. Total Environ.* **2021**, *775*, 145820. [\[CrossRef\]](#)
90. Lyu, T.; Zhang, L.; Xu, X.; Arias, C.A.; Brix, H.; Carvalho, P.N. Removal of the pesticide tebuconazole in constructed wetlands: Design comparison, influencing factors and modelling. *Environ. Pollut.* **2018**, *233*, 71–80. [\[CrossRef\]](#)
91. Rios-Montes, K.A.; Casas-Zapata, J.C.; Briones-Gallardo, R.; Peñuela, G. Optimal conditions for chlorothalonil and dissolved organic carbon in horizontal subsurface flow constructed wetlands. *J. Environ. Sci. Health B* **2017**, *52*, 274–281. [\[CrossRef\]](#)
92. Lv, T.; Carvalho, P.N.; Zhang, T.; Zhang, Y.; Button, M.; Arias, C.A.; Weber, K.; Brix, H. Functionality of microbial communities in constructed wetlands used for pesticide remediation: Influence of system design and sampling strategy. *Water Res.* **2017**, *110*, 241–251. [\[CrossRef\]](#)
93. Lv, T.; Zhang, Y.; Casas, M.E.; Carvalho, P.N.; Arias, C.A.; Bester, K.; Brix, H. Phytoremediation of imazalil and tebuconazole by four emergent wetland plant species in hydroponic medium. *Chemosphere* **2016**, *148*, 459–466. [\[CrossRef\]](#)
94. Amaro, A.C.E.; Baron, D.; Ono, E.O.; Rodrigues, J.D. Physiological effects of strobilurin and carboxamides on plants: An overview. *Acta Physiol. Plant* **2020**, *42*, 4. [\[CrossRef\]](#)
95. Wang, S.; Wang, J.; Zhang, X.; Xu, X.T.; Wen, Y.; He, J.; Zhao, Y.H. Freshwater quality criteria of four strobilurin fungicides: Interspecies correlation and toxic mechanism. *Chemosphere* **2021**, *284*, 131340. [\[CrossRef\]](#) [\[PubMed\]](#)
96. Maillard, E.; Payraudeau, S.; Faivre, E.; Grégoire, C.; Gangloff, S.; Imfeld, G. Removal of pesticide mixtures in a stormwater wetland collecting runoff from a vineyard catchment. *Sci. Total Environ.* **2011**, *409*, 2317–2324. [\[CrossRef\]](#) [\[PubMed\]](#)
97. McKnight, A.M.; Gannon, T.W.; Yelverton, F. Phytoremediation of azoxystrobin and imidacloprid by wetland plant species *Juncus effusus*, *Pontederia cordata* and *Sagittaria latifolia*. *Int. J. Phytoremediat.* **2021**, *14*, 196–204. [\[CrossRef\]](#)
98. Elsaesser, D.; Stang, C.; Bakanov, N.; Schulz, R. The Landau Stream Mesocosm Facility: Pesticide mitigation in vegetated flow-through streams. *Bull. Environ. Contam. Toxicol.* **2013**, *90*, 640–645. [\[CrossRef\]](#)
99. Pedersen, M.; Wegner, C.; Phansak, P.; Sarath, G.; Gaussoin, R.; Schlegel, V. Monitoring wheat mitochondrial compositional and respiratory changes using Fourier transform mid-infrared spectroscopy in response to agrochemical treatments. *Spectrochim. Acta Part A Mol. Biomol. Spectrosc.* **2017**, *173*, 727–732. [\[CrossRef\]](#) [\[PubMed\]](#)
100. Hellin, P.; King, R.; Urban, M.; Hammond-Kosack, K.E.; Legrève, A. The adaptation of *Fusarium culmorum* to DMI Fungicides Is Mediated by Major Transcriptome Modifications in Response to Azole Fungicide, Including the Overexpression of a PDR Transporter (FcABC1). *Front. Microbiol.* **2018**, *9*, 1385. [\[CrossRef\]](#)
101. Stang, C.; Bakanov, N.; Schulz, R. Experiments in water-macrophyte systems to uncover the dynamics of pesticide mitigation processes in vegetated surface waters/streams. *Environ. Sci. Pollut. Res.* **2016**, *23*, 673–682. [\[CrossRef\]](#) [\[PubMed\]](#)
102. Kiiker, R.; Juurik, M.; Heick, T.M.; Mäe, A. Changes in DMI, SDHI, and QoI Fungicide Sensitivity in the Estonian *Zymoseptoria tritici* Population between 2019 and 2020. *Microorganisms* **2021**, *9*, 814. [\[CrossRef\]](#)
103. Jiménez, R.R.; Alvarado, G.; Ruepert, C.; Ballester, E.; Sommer, S. The Fungicide Chlorothalonil Changes the Amphibian Skin Microbiome: A Potential Factor Disrupting a Host Disease-Protective Trait. *Appl. Microbiol.* **2021**, *1*, 26–37. [\[CrossRef\]](#)
104. Mackie, K.A.; Müller, T.; Kandeler, E. Remediation of copper in vineyards—A mini review. *Environ. Pollut.* **2012**, *167*, 16–26. [\[CrossRef\]](#)



105. Mosbach, A.; Edel, D.; Farmer, A.D.; Widdison, S.; Barchietto, T.; Dietrich, R.A.; Corran, A.; Scalliet, G. Anilinopyrimidine Resistance in *Botrytis cinerea* Is Linked to Mitochondrial Function. *Front. Microbiol.* **2017**, *8*, 2361. [[CrossRef](#)] [[PubMed](#)]
106. Dosnon-Olette, R.; Couderchet, M.; Eullaffroy, P. Phytoremediation of fungicides by aquatic macrophytes: Toxicity and removal rate. *Ecotoxicol. Environ. Saf.* **2009**, *72*, 2096–2101. [[CrossRef](#)] [[PubMed](#)]
107. Dosnon-Olette, R.; Couderchet, M.; El Arfaoui, A.; Sayen, S.; Eullaffroy, P. Influence of initial pesticide concentrations and plant population density on dimethomorph toxicity and removal by two duckweed species. *Sci. Total Environ.* **2010**, *408*, 2254–2259. [[CrossRef](#)] [[PubMed](#)]
108. Cao, C.; Huang, J.; Yan, C.; Liu, J.; Hu, Q.; Guan, W. Shifts of system performance and microbial community structure in a constructed wetland after exposing silver nanoparticles. *Chemosphere* **2018**, *199*, 661–669. [[CrossRef](#)]
109. Cao, C.; Huang, J.; Yan, C.; Zhang, X. Hydraulic flow direction alters impacts of AgNPs on pollutant removal and silver spatial distribution in vertical flow constructed wetlands. *Environ. Sci. Pollut. Res.* **2021**, *28*, 67736–67747. [[CrossRef](#)]
110. Xu, L.; Wu, C.; Oelmüller, R.; Zhang, W. Role of Phytohormones in *Piriformospora indica*-Induced Growth Promotion and Stress Tolerance in Plants: More Questions Than Answers. *Front. Microbiol.* **2018**, *9*, 1646. [[CrossRef](#)]
111. Saleem, M.; Arshad, M.; Hussain, S.; Bhatti, A.S. Perspective of plant growth promoting rhizobacteria (PGPR) containing ACC deaminase in stress agriculture. *J. Ind. Microbiol. Biotechnol.* **2007**, *34*, 635–648. [[CrossRef](#)]
112. Yang, X.; Chen, L.; Yang, Y.; Guo, X.; Chen, G.; Xiong, X.; Dong, D.; Li, G. Transcriptome analysis reveals that exogenous ethylene activates immune and defense responses in a high late blight resistant potato genotype. *Sci. Rep.* **2020**, *10*, 21294. [[CrossRef](#)]
113. Tripathi, G.D.; Javedm, Z.; Mishram, M.; Fasakem, V.; Dashora, K. Phytomelatonin in stress management in agriculture. *Heliyon* **2021**, *7*, e06150. [[CrossRef](#)]
114. Zhang, S.; Zheng, X.; Reiter, R.J.; Feng, S.; Wang, Y.; Liu, S.; Jin, L.; Li, Z.; Datla, R.; Ren, M. Melatonin attenuates potato late blight by disrupting cell growth, stress tolerance, fungicide susceptibility and homeostasis of gene expression in *Phytophthora infestans*. *Front. Plant Sci.* **2017**, *8*, 1993. [[CrossRef](#)]
115. Arnao, M.B.; Hernández-Ruiz, J. Role of Melatonin to Enhance Phytoremediation Capacity. *Appl. Sci.* **2019**, *9*, 5293. [[CrossRef](#)]
116. Yan, Y.; Sun, S.; Zhao, N.; Yang, W.; Shi, Q.; Gong, B. COMT1 overexpression resulting in increased melatonin biosynthesis contributes to the alleviation of carbendazim phytotoxicity and residues in tomato plants. *Environ. Pollut.* **2019**, *252*, 51–61. [[CrossRef](#)] [[PubMed](#)]
117. Koo, Y.M.; Heo, A.Y.; Choi, H.W. Salicylic Acid as a Safe Plant Protector and Growth Regulator. *Plant Pathol. J.* **2020**, *36*, 1–10. [[CrossRef](#)] [[PubMed](#)]
118. Li, L.; Zou, Y. Induction of disease resistance by salicylic acid and calcium ion against *Botrytis cinerea* in tomato (*Lycopersicon esculentum*). *Emir. J. Food Agric.* **2017**, *29*, 78–82. [[CrossRef](#)]
119. Li, Y.; Qiu, L.; Zhang, Q.; Zhuansun, X.; Li, H.; Chen, X.; Krugman, T.; Sun, Q.; Xie, C. Exogenous sodium diethyldithiocarbamate, a Jasmonic acid biosynthesis inhibitor, induced resistance to powdery mildew in wheat. *Plant Direct.* **2020**, *4*, e00212. [[CrossRef](#)]
120. Zalewski, K.; Lahuta, L.B.; Martysiak-Żurowska, D.; Okorski, A.; Nitkiewicz, B.; Zielonka, Ł. Effect of Exogenous Application of Methyl Jasmonate on the Lipid and Carbohydrate Content and Composition of Winter Triticale (*Triticosecale* Wittm.) Grain and the Severity of Fungal Infections in Triticale Plants and Grain. *J. Agric. Food Chem.* **2019**, *67*, 5932–5939. [[CrossRef](#)]
121. Rostami, S.; Azhdarpoor, A. The application of plant growth regulators to improve phytoremediation of contaminated soils: A review. *Chemosphere* **2019**, *220*, 818–827. [[CrossRef](#)]
122. Rostami, S.; Azhdarpoor, A.; Rostami, M.; Samaei, M.R. The effects of simultaneous application of plant growth regulators and bioaugmentation on improvement of phytoremediation of pyrene contaminated soils. *Chemosphere* **2016**, *161*, 219–223. [[CrossRef](#)]
123. Khan, A.; Halo, B.A.; Elyassi, A.R.; Ali, S.; Ali, S.; Al-Hosni, K.; Hussain, J.; Al-Harrasi, A.; Lee, I. Indole acetic acid and ACC deaminase from endophytic bacteria improves the growth of *Solanum lycopersicum*. *Electron. J.* **2016**, *21*, 58–64. [[CrossRef](#)]
124. Siddiqui, M.H.; Al-Waibi, M.H.; Basalah, M.O. Interactive effect of calcium and gibberellin on nickel tolerance in relation to antioxidant systems in *Triticum aestivum* L. *Protoplasma* **2011**, *248*, 503–511. [[CrossRef](#)]
125. Ji, P.; Tang, X.; Jiang, Y.; Tong, Y.; Gao, P.; Han, W. Potential of Gibberellic Acid 3 (GA3) for Enhancing the Phytoremediation Efficiency of *Solanum nigrum* L. *Bull. Environ. Contam. Toxicol.* **2015**, *95*, 810–814. [[CrossRef](#)] [[PubMed](#)]
126. Senaratna, T.; Touchell, D.; Bunn, E.; Dixon, K. Acetyl salicylic acid (Aspirin) and salicylic acid induce multiple stress tolerance in bean and tomato plants. *J. Plant Growth Regul.* **2000**, *30*, 157–161. [[CrossRef](#)]
127. Mateo, A.; Funck, D.; Mühlenbock, P.; Kular, B.; Mullineaux, P.M.; Karpinski, S. Controlled levels of salicylic acid are required for optimal photosynthesis and redox homeostasis. *J. Exp. Bot.* **2006**, *57*, 1795–1807. [[CrossRef](#)] [[PubMed](#)]
128. Afrousheh, M.; Tehranifar, A.; Shoor, M.; Safari, V.R. Salicylic Acid Alleviates the Copper Toxicity in *Tagetes Erecta*. *Int. J. Farming Allied Sci.* **2015**, *4*, 232–238.
129. Mirbolook, A.; Rasouli-Sadaghiani, M.; Sepehr, E.; Lakzian, A.; Hakimi, M. Synthesized Zn(II)-Amino Acid and -Chitosan Chelates to Increase Zn Uptake by Bean (*Phaseolus vulgaris*) Plants. *J. Plant Growth Regul.* **2021**, *40*, 831–847. [[CrossRef](#)]
130. Buzón-Durán, L.; Martín-Gil, J.; Marcos-Robles, J.L.; Fombellida-Villafuella, Á.; Pérez-Lebeña, E.; Martín-Ramos, P. Antifungal Activity of Chitosan Oligomers–Amino Acid Conjugate Complexes against *Fusarium culmorum* in Spelt (*Triticum spelta* L.). *Agronomy* **2021**, *10*, 1427. [[CrossRef](#)]
131. Dibona-Villanueva, L.; Fuentealba, D. Novel Chitosan-Riboflavin Conjugate with Visible Light-Enhanced Antifungal Properties against *Penicillium digitatum*. *J. Agric. Food Chem.* **2021**, *69*, 945–954. [[CrossRef](#)]

132. Bloem, E.; Haneklaus, S.; Haensch, R.; Schnug, E. EDTA application on agricultural soils affects microelement uptake of plants. *Sci. Total Environ.* **2017**, *577*, 166–173. [\[CrossRef\]](#)
133. Zaier, H.; Ghnaya, T.; Ben Rejeb, K.; Lakhdar, A.; Rejeb, S.; Jemal, F. Effects of EDTA on phytoextraction of heavy metals (Zn, Mn and Pb) from sludge-amended soil with *Brassica napus*. *Bioresour. Technol.* **2010**, *101*, 3978–3983. [\[CrossRef\]](#)
134. Wang, K.; Liu, Y.; Song, Z.; Wang, D.; Qiu, W. Chelator complexes enhanced *Amaranthus hypochondriacus* L. phytoremediation efficiency in Cd-contaminated soils. *Chemosphere* **2019**, *237*, 124480. [\[CrossRef\]](#)
135. Batool, A.; Saleh, T.A. Removal of toxic metals from wastewater in constructed wetlands as a green technology; catalyst role of substrates and chelators. *Ecotoxicol. Environ. Saf.* **2020**, *189*, 109924. [\[CrossRef\]](#)
136. Köhl, J.; Kolnaar, R.; Ravensberg, W.J. Mode of Action of Microbial Biological Control Agents Against Plant Diseases: Relevance Beyond Efficacy. *Front. Plant Sci.* **2019**, *10*, 845. [\[CrossRef\]](#) [\[PubMed\]](#)
137. Matzen, N.; Heick, M.S.; Jørgensen, L.N. Control of powdery mildew (*Blumeria graminis* spp.) in cereals by Serenade® ASO (*Bacillus amyloliquefaciens* (former subtilis) strain QST 713). *Biol. Control* **2019**, *139*, 104067. [\[CrossRef\]](#)
138. Ongena, M.; Jourdan, E.; Adam, A.; Paquot, M.; Brans, A.; Joris, B.; Arpigny, J.L.; Thonart, P. Surfactin and fengycin lipopeptides of *Bacillus subtilis* as elicitors of induced systemic resistance in plants. *Environ. Microbiol.* **2007**, *9*, 1084–1090. [\[CrossRef\]](#) [\[PubMed\]](#)
139. Tang, M.; Zhang, F.; Yao, S.; Liu, Y.; Chen, J. Application of *Pseudomonas flava* WD600 3 for sewage treatment in constructed wetland in winter. *Environ. Technol.* **2015**, *36*, 1205–1211. [\[CrossRef\]](#) [\[PubMed\]](#)
140. Syranidou, E.; Christofilopoulos, S.; Gkavrou, G.; Thijs, S.; Weyens, N.; Vangronsveld, J.; Kalogerakis, N. Exploitation of Endophytic Bacteria to Enhance the Phytoremediation Potential of the Wetland Helophyte *Juncus acutus*. *Front. Microbiol.* **2016**, *7*, 1016. [\[CrossRef\]](#) [\[PubMed\]](#)
141. Riva, V.; Riva, F.; Vergani, L.; Crotti, E.; Borin, S.; Mapelli, F. Microbial assisted phytodepuration for water reclamation: Environmental benefits and threats. *Chemosphere* **2020**, *241*, 1248432. [\[CrossRef\]](#)
142. Shahid, M.J.; AL-surhane, A.A.; Kouadri, F.; Ali, S.; Nawaz, N.; Afzal, M.; Rizwan, M.; Ali, B.; Soliman, M.H. Role of microorganisms in the remediation of wastewater in floating treatment wetlands: A review. *Sustainability* **2020**, *12*, 5559. [\[CrossRef\]](#)
143. Glick, B.R. Plant growth-promoting bacteria: Mechanisms and applications. *Scientifica* **2012**, *2012*, 963401. [\[CrossRef\]](#)
144. DalCorso, G.; Fasani, E.; Manara, A.; Visioli, G.; Furini, A. Heavy Metal Pollutions: State of the Art and Innovation in Phytoremediation. *Int. J. Mol. Sci.* **2019**, *20*, 3412. [\[CrossRef\]](#)
145. Myresiotis, C.K.; Vryzas, Z.; Papadopoulou-Mourkidou, E. Enhanced root uptake of acibenzolar-S-methyl (ASM) by tomato plants inoculated with selected *Bacillus* plant growth-promoting rhizobacteria (PGPR). *Appl. Soil Ecol.* **2014**, *77*, 26–33. [\[CrossRef\]](#)
146. Xiao, W.; Wang, H.; Li, T.; Zhu, Z.; Zhang, J.; He, Z.; Yang, X. Bioremediation of Cd and carbendazim co-contaminated soil by Cd-hyperaccumulator *Sedum alfredii* associated with carbendazim-degrading bacterial strains. *Environ. Sci. Pollut. Res.* **2013**, *20*, 380–389. [\[CrossRef\]](#) [\[PubMed\]](#)
147. Podbielska, M.; Kus-Liśkiewicz, M.; Jagusztyn, B.; Piechowicz, B.; Sadło, S.; Słowik-Borowiec, M.; Twarużek, M.; Szpyrka, E. Influence of *Bacillus subtilis* and *Trichoderma harzianum* on Penthiopyrad Degradation under Laboratory and Field Studies. *Molecules* **2020**, *25*, 1421. [\[CrossRef\]](#) [\[PubMed\]](#)
148. Myresiotis, C.K.; Vryzas, Z.; Papadopoulou-Mourkidou, E. Biodegradation of soil-applied pesticides by selected strains of plant growth-promoting rhizobacteria (PGPR) and their effects on bacterial growth. *Biodegradation* **2012**, *23*, 297–310. [\[CrossRef\]](#)
149. Liao, D.; Wang, S.; Cui, M.; Liu, J.; Chen, A.; Xu, G. Phytohormones Regulate the Development of Arbuscular Mycorrhizal Symbiosis. *Int. J. Mol. Sci.* **2018**, *19*, 3146. [\[CrossRef\]](#)
150. Göhre, V.; Paszkowski, U. Contribution of the arbuscular mycorrhizal symbiosis to heavy metal phytoremediation. *Planta* **2016**, *223*, 1115–1122. [\[CrossRef\]](#)
151. Hussain, Z.; Arslan, M.; Shabir, G.; Malik, M.H.; Mohsin, M.; Iqbal, S.; Afzal, M. Remediation of textile bleaching effluent by bacterial augmented horizontal flow and vertical flow constructed wetlands: A comparison at pilot scale. *Sci. Total Environ.* **2019**, *685*, 370–379. [\[CrossRef\]](#)
152. Fahid, M.; Arslan, M.; Shabir, G.; Younus, S.; Yasmeen, T.; Rizwan, M.; Siddique, K.; Ahmad, S.R.; Tahseen, R.; Iqbal, S.; et al. *Phragmites australis* in combination with hydrocarbons degrading bacteria is a suitable option for remediation of diesel-contaminated water in floating wetlands. *Chemosphere* **2020**, *240*, 124890. [\[CrossRef\]](#)