

Article

Antiobesity Effects of Ginsenoside Rg1 on 3T3-L1 Preadipocytes and High Fat Diet-Induced Obese Mice Mediated by AMPK

Huimin Liu ^{1,2}, Jing Wang ¹, Meihong Liu ^{2,3}, Hongyu Zhao ⁴, Sanabil Yaqoob ^{2,3}, Mingzhu Zheng ^{2,3}, Dan Cai ^{2,3} and Jingsheng Liu ^{2,3,*}

¹ College of Life Science, Jilin Agricultural University, Changchun, Jilin 130118, China; liuhuimin@jlau.edu.cn (H.L.); wangjing11151204@163.com (J.W.)

² National Engineering Laboratory for Wheat and Corn Deep Processing, Changchun, Jilin 130118, China; liumh@jlau.edu.cn (M.L.); sanabily67@gmail.com (S.Y.); zhengmzhu@163.com (M.Z.); dan1980623@163.com (D.C.)

³ College of Food Science and Engineering, Jilin Agricultural University, Changchun, Jilin 130118, China

⁴ Chinese Medicine Science Academy of Jilin Province, Changchun, Jilin 130118, China; fixiov5815@126.com

* Correspondence: liujs1007@vip.sina.com; Tel.: +86-431-8453-2886

Received: 16 May 2018; Accepted: 21 June 2018; Published: 27 June 2018

Supplementary File

Table S1. Primers sequences used in qRT-PCR analysis.

Gene	Forward primers	Reverse Primers
PPAR γ	AGACCACTCGCATTCCTTT	CCACAGACTCGGCACTCAAT
C/EBP α	ACCGGGTTTCGGGACTTGA	CCCGCAGGAACATCTTTAAGTGA
HSL	TCCTGGAACCTAAGTGGACGCAAG	CAGACACACTCCTGCGCATAGAC
Perilipin	GTCAATGAACAAGGGCCCAAC	CACAGGCAGCTGCAGAACTCTC
FABP4	TGGGAACCTGGAAGCTTGTCTC	GAATTCCACGCCCAGTTTGA
FAS	TGCTGTTGGAAGTCAGCTATGAA	GATGCCTCTGAACCACTCACAC
SREBP1-c	CCGAGATGTGCGAACTGGA	GAAGTCACTGTCTTGTTGTTGATG
ACC	AGTGATGGTGGCCTGCTCTTG	AGCAGACGGTGAGCGCATTA
β -actin	CATCCGTAAAGACCTCTATGCCAAC	ATGGAGCCACCGATCCACA