

Table S1. KDQOL scores according to diet cluster.

	hiSSB (n=20)	loSSB (n = 28)	<i>P</i> value between groups
Average Symptom KDQOL	77 (68 - 82)	81 (73 -88)	0.228
Average Effects KDQOL	73 (63 - 89)	81 (70 -94)	0.176
Average Burden KDQOL	41 (25 - 72)	69 (31 -88)	0.058
Average S12 PCS KDQOL	33 (27 -42)	38 (29 -44)	0.363
Average SF MCS KDQOL	47 (39 - 58)	57 (48 -61)	0.026

Data is presented as Median (Q1-Q3); Mann-Whitney U test

Table S2: Comparison of KDQOL score by selected variables

Variable	Symptoms	Effects	Burden	PCS	MCS
Diet Cluster					
hiSSB	73.8±13.1	74.1±17.7	48.8±26.7	33.5±11.1	46.1±12.7 ^a
loSSB	79.2±10.7	81.1±15.6	64.8±29.0	36.5±11.0	53.0± 9.2 ^a
Gender					
Male	77.4±11.6	78.2±16.8	56.3±28.6	35.4±11.7	50.7±12.5
Female	76.5±12.6	78.1±17.1	59.8±29.7	35.0±10.3	50.6±10.0
Tobacco use					
Yes	74.8±14.1	77.9±21.5	56.3±33.2	35.0±10.7	50.6±12.6
No	77.9±10.9	78.3±14.4	58.5±27.1	35.3±11.3	50.7±11.0
Antidepressant use					
Yes	66.1±17.6	66.3±24.9	42.5±29.1	28.0±9.2	43.6±10.9
No	78.2±10.7	79.6±15.3	59.6±28.6	36.1±11.0	51.5±11.3
Age [β(se)]	0.36(0.15)*	0.52(0.19)*	0.32(0.36)	0.03(0.13)	0.27(0.14)
Vintage [β(se)]	-0.03(0.03)	0.06(0.03)	0.13(0.06)*	-0.06(0.02)*	0.01(0.02)
BMI [β(se)]	-0.03(0.35)	-0.20(0.45)	0.56(0.80)	-0.35(0.28)	-0.46(0.30)

^a significant difference at 0.05. *significant association at 0.05. Values are mean ± SD. BMI: Body Mass Index.