

Article

Association between Non-Alcoholic Fatty Liver Disease and Dietary Habits, Stress, and Health-Related Quality of Life in Korean Adults

A Lum Han

Department of Family Medicine, Wonkwang University Hospital, Iksan 54538, Korea; qibosarang@naver.com; Tel.: +82-063-859-1301

Received: 8 April 2020; Accepted: 24 May 2020; Published: 27 May 2020



Abstract: Considering the increasing prevalence of non-alcoholic fatty liver disease (NAFLD), this study aimed to evaluate the association between NAFLD and dietary habits, stress, and health-related quality of life (HRQoL) in Korean individuals by using data from the Korea National Health and Nutrition Examination Survey (KNHANES) VI 2013–2015. NAFLD was defined in individuals with a hepatic steatosis index (HSI) value ≥ 36 . Eating habits were assessed based on the frequencies of eating and eating out; stress was assessed through the stress perception rate; and the EuroQol-5D (EQ-5D) questionnaire was used to assess the HRQoL. We performed a complex sample logistic regression analysis and estimated the odds ratios by adjusting for significant factors to evaluate associations between NAFLD and dietary habits, stress, and HRQoL. Occurrence of NAFLD was not significantly associated with meal frequencies over one week. With an increase in stress, based on the stress perception rate, the risk of NAFLD increased 1.316-fold (95% confidence interval (CI): 1.175–1.469, $p < 0.05$). Additionally, a decrease in the EQ-5D score by 1 increased the risk of NAFLD 3.38-fold (95% CI: 1.893–4.844, $p < 0.05$). Thus, NAFLD treatment should include stress management, and underlying HRQoL should be considered during treatment.

Keywords: non-alcoholic fatty liver disease; health-related quality of life; stress; dietary habits

1. Introduction

The incidence of non-alcoholic fatty liver disease (NAFLD) has been increasing globally, including in children and adolescents. NAFLD currently accounts for 75% of the chronic liver disease cases in the United States and its prevalence has doubled in Korea, Japan, and the United Kingdom [1–3]. NAFLD has been linked to obesity, various metabolic diseases, and even cardiovascular diseases and mortality. NAFLD occurs independently of alcohol consumption and is associated with an abnormality in insulin sensitivity and fatty acid uptake in the liver, and metabolic alterations mediated by an inflammatory mechanism [2,4]. As with obesity, lifestyle habits, such as lack of exercise and unhealthy diet, have a great impact on the development and exacerbation of NAFLD [1,3,5]. Several studies have shown that consumption of certain nutrients, such as carbohydrates and fructose, affect NAFLD. High intake of carbohydrates, contributing to over 60% of daily energy consumption, or excessive simple sugar intake have been reported to exacerbate NAFLD [3,6,7].

Many studies have also shown that obesity and metabolic disorders can be associated with breakfast habits; particularly when the least of a day's food is eaten at breakfast, breakfast is skipped or of poor quality, or a nocturnal eating habit affects subsequent breakfast intake [8–12]. Notably, a small breakfast size exacerbated obesity in mouse experiments [8], and nocturnal syndrome was found to be correlated to obesity [9]. Furthermore, consumption of snacks along with skipping breakfast was related to obesity in adolescents [10]. In contrast, a regular diet reduced incidence of obesity and chronic diseases [11] and breakfast has been shown to be important for diet success [12].

Similarly, several studies have investigated the relationship between dietary habits and NAFLD. Poor eating habits, such as high consumption of calories, from carbohydrates and fats, and low intake of vitamins and minerals, are linked to the development of NAFLD [5]. Moreover, use of nutritional supplements can reduce the incidence of NAFLD [13]. In a study that analyzed meal patterns and NAFLD in Korea; traditional, Western and high-carbohydrate, and simple meal patterns were investigated [14]. The traditional eating pattern increased NAFLD, while a simple meal pattern decreased NAFLD [14]. NAFLD patients had a low Korean dietary diversity score, a tool for assessing a balanced diet [15]. However, most existing studies have been performed across different races and countries, and by assessing dietary habits in various ways. As the data on Korean population is limited, there is a need for further studies in Korean patients with NAFLD.

NAFLD is also associated with depression, psychological distress, and psychiatric disease [16–20]. Stress can induce and exacerbate chronic metabolic diseases through excessive release of neurotransmitters such as norepinephrine, epinephrine, hormones such as cortisol, and inflammatory cytokines [18–23]. These findings suggest that mental state affects the development of NAFLD through hormones and inflammatory mechanisms. In addition, stress also exacerbates chronic diseases and metabolic diseases through poor lifestyle habits such as overeating, binge eating, excessive alcohol consumption, and smoking [21]. However, studies linking stress with NAFLD are lacking.

The above studies indicate that poor lifestyle, depression, stress, chronic illness, and mortality are associated with NAFLD. Therefore, this provides a good basis for investigating the potential link between NAFLD and health-related quality of life (HRQoL). Several studies report that patients with NAFLD score low on the quality of life assessment tool, in terms of physical, emotional, and economical assessments [24–27]. The EuroQol- 5D (EQ-5D), developed in Europe, has proven useful as an HRQoL assessment tool in many studies [28–31], including those involving Koreans [32–34].

The hepatic steatosis index (HSI), which is based on levels of alanine aminotransferase (ALT), aspartate aminotransferase (AST), and body-mass index (BMI), is a useful index to diagnose NAFLD that has been validated in various studies [35–40]. HSI was reported to have an area under the receiver operating characteristic (AUROC) of 0.81 for the diagnosis of NAFLD [35], which, although a slightly lower AUROC than that of the Framingham Steatosis Index, was of diagnostic value [36]. Another study reported an AUROC of 0.83 for HSI and the tool was deemed to be convenient [37]. Moreover, HSI is cost effective and can be performed before computed tomography (CT) and magnetic resonance imaging (MRI) based diagnoses [38]. In a large-scale cohort study, HSI was found to be an effective indicator for predicting NAFLD in comparison to results using ultrasound [39]. HSI was used in a Korean study that revealed the correlation between NAFLD and asymptomatic hypothyroidism [40].

Based on the results from these studies, we were interested in the correlation between NAFLD, and dietary habits, stress, and HRQoL in a Korean population. We investigated the association between NAFLD, diagnosed using HSI, and these factors using representative data from Koreans.

2. Materials and Methods

2.1. Study Design

The data are based on a survey by the Korean Centers for Disease Control and Prevention (KCDC) of Korea National Health and Nutrition Examination Survey (KNHANES) VI 2013–2015. Since 1998, KNHANES has conducted a continuous nationwide survey to monitor the health and nutrition of Koreans every year [41]. The survey consists of three components: a health interview, a health checkup, and a nutrition survey. Health interviews and examinations are carried out by trained clinicians. The survey uses a stratified, multi-stage clustered probability sampling. KNHANES has been approved by the KCDC's Agency Review Board (ARB) [41]. This study followed ethical standards laid out in the declaration of Helsinki. The study was approved by the clinical trial screening committee of Wonkwang University Hospital (ARB approval number 2020-03-012). The KNHANES VI survey was conducted on those aged 19 or older who faithfully completed the survey.

The following persons were excluded: patients with excessive alcohol use (defined as >20 g/day for males and >10 g/day for females); hepatic cellular carcinoma; hepatitis virus B and/or C (HBV, HCV) with systematic history and blood tests; autoimmune hepatitis, drug-induced liver damage, or other liver diseases; liver cirrhosis; or other medical conditions that may affect HSI levels.

The subjects were divided into two groups, one group comprising of suspected NAFLD cases and the other a control group (Figure 1). The suspected NAFLD cases were defined as those with an HSI > 36, while the control group had HSI ≤ 36; where HSI was calculated as follows: $HSI = 8 \times (ALT/AST \text{ ratio}) + BMI (+2 \text{ for women})$. We used HSI to define NAFLD since the fatty liver index (FLI) cannot be used as gamma glutamyl transferase data are not presented in KNHANES, and ultrasound or CT were not performed.




Figure 1. Flow diagram of subjects recruited in this study. HCC, hepatic cellular carcinoma; HCV, hepatitis virus C; HBV, hepatitis virus B; LC, liver cirrhosis; HSI, hepatic steatosis index; KNHANES, Korea National Health and Nutrition Examination Survey.

The usual level of stress perception was investigated with questions that required the individual to respond by selecting one of the following answers: (1) I feel very much, (2) I feel a lot, (3) I feel a little, (4). A person who responded with a (1) or (2) was given a score of 1, and those who marked answers (3) or (4) were scored 0, with 1 indicating high stress and 0 indicating low stress.

HRQoL was evaluated by the widely used EQ-5D [29]. EQ-5D consists of five multiple choice questions and one subjective health level. The health status was determined with regards to five areas: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression. The individuals were asked to choose one of the following three levels: “No problem at all”, “There are some problems”, or “There are many problems.” The responses to the five questions were subjected to the score conversion system, and expressed as a point between 1, which means excellent health, and −1, which indicates health worse than death. The EQ-5D index was then calculated using the following equation:

$$\text{EQ-5D index} = 1 - (0.0081 + (0.1140 \times M2) + (0.6274 \times M3) + (0.0572 \times SC2) + (0.2073 \times SC3) + (0.0615 \times UA2) + (0.2812 \times UA3) + (0.0581 \times PD2) + (0.2353 \times PD3) + (0.0675 \times AD2) + (0.2351 \times AD3))$$

We compared the general factors, stress perception, and EQ-5D values between the normal (HSI ≤ 36) and NAFLD (HSI > 36) groups. We also examined the general factors, cardiovascular-related factors (from blood tests), dietary habits (meal patterns), and whether the individuals had companions at meals in both groups. The relationship between suspected NAFLD and control group and general

characteristics, dietary habits, stress, and EQ-5D are presented as an odds ratio (OR). Furthermore, to determine the most important variables in the relationship between the NAFLD and control groups and dietary habits, stress, and EQ-5D, the other variables were corrected, and the correlations are presented as OR.

2.2. Anthropometric Measurements

Anthropometric and biochemical measurements were performed by an experienced inspector. The height (cm) and weight (kg) were measured to the nearest 0.1 value, using a Seca 225 instrument (Seca, Hamburg, Germany) and GL-6000-20 scale (G-tech, Seoul, Korea), respectively. BMI was calculated as weight (kg)/square of height (m²). Waist circumference (WC) was measured between the margin under the ribs and the iliac crest and recorded to 0.1 cm.

Laboratory tests using blood tests were taken after fasting for 8 h, and samples were immediately transported to a central laboratory (Neodin Medical Institute, Seoul, Korea) for processing. The Hitachi 7600 automatic analyzer (Hitachi, Tokyo, Japan) was used to measure fasting plasma glucose, HbA_{1c} (hemoglobin A_{1c}), total cholesterol, triglyceride (TG), high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL) cholesterol, aspartate aminotransferase (AST), and alanine aminotransferase (ALT) levels.

2.3. Sociodemographic and Lifestyle Variables

A self-written questionnaire was used to investigate the subject's sociodemographic and lifestyle variables. Based on the recent smoking status, individuals were divided into "smokers who are currently smoking" (within two years) and "non-smokers." Alcohol consumption was expressed as grams of alcohol consumed per week. Physical exercise was measured using the International Physical Activity Questionnaire [42]. Regular exercise was defined as three or more times per week and 20 min or more per trial. The education level was divided into four categories: under elementary school graduation, middle school graduation, high school graduation, and above university graduation.

Dietary habits were investigated for the number of breakfasts, lunches, and dinners during a week over the past year and the presence or absence of companions at the time of the meals.

2.4. Statistical Analysis

Statistical analyses were performed using PASW (Predictive Analytics SoftWare) statistics software (SPSS version 24.0, IBM SPSS Inc., Chicago, IL, USA). Overall, frequency analysis was performed using the complex sample plan for frequencies analysis. A complex sample Rao-Scott adjusted chi-square test and a complex sample generalized linear model were used to compare general characteristics and dietary habits, stress, HRQoL, and NAFLD. Values are presented as the number (%) or the mean \pm standard deviation. A complex sample logistic regression test was used to correlate NAFLD with general characteristics and dietary habits, stress, and HRQoL. We performed a complex sample Rao-Scott adjusted Chi-square test and complex sample generalized linear model t-test (Table 1) to investigate the differences between general characteristics and dietary habits, stress, HRQoL, and NAFLD.

We performed a complex sample logistic regression analysis and estimated the OR and 95% confidence intervals (CI) by adjusting for statistically significant factors (Table 2) to investigate the associations between dietary habits, stress, and HRQoL and potential NAFLD. All data are presented as means \pm standard errors (SEs) or percentages (%) for categorical variables. *p* values < 0.05 were considered statistically significant.

3. Results

In total, 17,726 subjects (50.6% male; 49.4% female) were recruited in this study, with an average age of 43.93 \pm 0.25 years. Table 1 summarizes the differences between the general characteristics, dietary habits, stress perception, and HRQoL among individuals with and without NAFLD. The rate

of practice of aerobic exercise was lower among individuals with than in those without NAFLD (41.7% vs. 42.1%; $p < 0.0001$). In a weekly dinner frequency survey over the past year, the rate of almost no daily consumption of dinner was slightly but not significantly lower in the NAFLD group than in the non-NAFLD group. The correlation analysis between EQ-5D and stress perception rate showed a strong correlation with the parameter estimation value of -0.994 . In other words, it was found that EQ-5D decreased whenever the stress perception rate increased.

Table 1. Differences between general characteristics and dietary habits, stress, HRQoL, and NAFLD.

		Total (17,726)	HSI \leq 36	HSI $>$ 36	<i>p</i> -Value ^a
Sex	Male	7986 (50.6)	6004 (47.5)	1982 (61.3)	<0.0001
	Female	9740 (49.4)	7958 (52.5)	1782 (38.7)	
Age	years	43.93 \pm 0.25	43.59 \pm 0.27	45.1 \pm 0.34	<0.0001
Educational level	\leq Elementary	4129 (17.3)	3360 (18.1)	769 (14.5)	<0.0001
	Middle	2064 (11.2)	1623 (11.3)	441 (10.8)	
	High	5010 (31)	3868 (30.7)	1142 (32.3)	
	\geq College	5753 (35.9)	4515 (35.5)	1238 (37.4)	
Smoking		2905 (19.5)	2110 (18.1)	795 (24.4)	<0.0001
Exercise		6956 (42)	5495 (42.1)	1461 (41.7)	<0.0001
Alcohol		8790 (54.1)	6860 (53.5)	1930 (56)	<0.0001
Stress		4637 (28.2)	3496 (26.9)	1141(32.7)	<0.0001
Breakfast frequency	5–7/week	10,153 (60.2)	8129 (61.1)	2024 (57)	0.002
	3–4/week	1702 (12.3)	1332 (12.2)	370 (12.6)	
	1–2/week	1603 (12.2)	1234 (11.7)	369 (14.2)	
	Rare	1908 (15.2)	1475 (15)	433 (16.2)	
Lunch frequency	5–7/week	13,933 (90.4)	11,074 (90.8)	2859 (89)	0.047
	3–4/week	917 (6.3)	708 (6.2)	209 (7)	
	1–2/week	261 (1.7)	198 (1.6)	63 (2.1)	
	Rare	255 (1.5)	190 (1.4)	65 (2)	
Dinner frequency	5–7/week	13,980 (90.1)	11,079 (90.1)	2901 (90.1)	0.783
	3–4/week	1106 (8.1)	859 (8)	247 (8.3)	
	1–2/week	218 (1.5)	180 (1.6)	38 (1.3)	
	Rare	62 (0.4)	52 (0.4)	10 (0.3)	
Companion for breakfast		7120 (42.3)	5692 (42.8)	1428 (40.4)	0.002
Companion for lunch		10,343 (69.1)	8234 (69.3)	2109 (68)	0.042
Companion for dinner		11,676 (77)	9240 (76.8)	2436 (77.7)	0.485
WC		81.09 \pm 0.13	77.84 \pm 0.12	92.98 \pm 0.17	<0.0001
BMI		23.66 \pm 0.04	22.37 \pm 0.03	28.3 \pm 0.06	<0.0001
Fasting glucose		99.28 \pm 0.23	96.71 \pm 0.2	108.38 \pm 0.6	<0.0001
HbA _{1c}		5.61 \pm 0.01	5.52 \pm 0.01	5.92 \pm 0.02	<0.0001
TC		190.3 \pm 0.38	187.96 \pm 0.38	198.54 \pm 0.82	<0.0001
HDL-C		51.11 \pm 0.14	52.63 \pm 0.15	45.8 \pm 0.2	<0.0001
TG		134.18 \pm 1.23	120.86 \pm 1.32	180.56 \pm 2.56	<0.0001
LDL-C		119.07 \pm 0.82	114.85 \pm 1	124.47 \pm 1.27	<0.0001
Stress		0.28 \pm 0.004	0.27 \pm 0.005	0.33 \pm 0.01	<0.0001
EQ-5D		0.95 \pm 0.001	0.96 \pm 0.001	0.95 \pm 0.002	<0.0001

Abbreviations: HRQoL: health-related quality of life; NAFLD: Non-alcoholic fatty liver disease; WC: waist circumference; BMI, body-mass index; HbA_{1c}, hemoglobin A_{1c}; TC, total cholesterol; HDL-C, high-density lipoprotein cholesterol; TG, triglyceride; LDL-C, low-density lipoprotein cholesterol; Stress, if '1' is recorded for the stress question; EQ-5D, EuroQol- 5 Dimension; HSI, hepatic steatosis index. Definitions: Smoking: the percentage of individuals who are currently smoking (within two years); alcohol, the percentage of alcohol intake more than once a month in the last year; breakfast (lunch and dinner) frequency: breakfast (lunch and dinner) frequency for 1 week in the last year; HSI, hepatic steatosis index = $8 \times$ (alanine aminotransferase / aspartate aminotransferase ratio) + BMI (+2 for women); NAFLD, HSI \geq 36. Values are presented as the number (%) or the mean \pm standard deviation. ^a The *p*-value was determined through the complex sample Rao–Scott adjusted Chi-square test and complex sample generalized linear model *t*-test.

Table 2 delineates the association between NAFLD risk and dietary habits, stress, and HRQoL. The risk of developing NAFLD was higher when skipping breakfast and in the absence of a companion for breakfast (OR: 1.165 and CI: 1.016–1.337, OR: 1.008 and CI: 0.903–1.126, respectively, $p < 0.05$). The NAFLD risk was higher when eating out than when not eating out (OR: 1.250 and CI: 1.006–1.552, $p < 0.05$). Every 1-in increase in WC and an increase in BMI increased the risk of NAFLD (OR: 1.237 and CI: 1.227–1.247, OR: 2.389 and CI: 2.309–2.471, respectively, $p < 0.05$). Every 1-unit increase in FBS (fasting blood sugar), HbA_{1c}, total cholesterol, HDL cholesterol, triglycerides, and LDL cholesterol also increased the NAFLD risk (OR: 1.021 and CI: 1.018–1.023, OR: 1.829 and CI: 1.707–1.96, OR: 1.008 and CI: 1.006–1.009, OR: 0.948 and CI: 0.944–0.952, OR: 1.004 and CI: 1.003–1.005, and OR: 1.008 and CI: 1.005–1.011, respectively, $p < 0.05$). Furthermore, the NAFLD risk was higher in the stress-perceiving group (OR: 1.316 and CI: 1.203–1.44, $p < 0.05$). When the EQ-5D score decreased by 1, the risk of developing NAFLD increased (OR: 2.255 and CI: 1.533–3.317, $p < 0.05$).

Table 3 delineates the association between NAFLD and eating habits, HRQoL, and stress after correcting for significant variables. The frequency of breakfast, lunch, and dinner for one week were not significantly associated with NAFLD occurrence. In the stress-perceiving group, the risk of developing NAFLD was higher (OR: 1.314 and CI: 1.175–1.469, $p < 0.05$), and when EQ-5D decreased by 1, the risk of developing NAFLD increased (OR: 3.028 and CI: 1.893–4.844, $p < 0.05$).

Table 2. General characteristics and the association between NAFLD and dietary habits, stress, and HRQoL.

Characteristics		OR	(95% CI)	<i>p</i> -Value ^a
Sex	Male	1.747	(1.609–1.898)	<0.0001
	Female	1	Reference	
Age		1.047	(1.025–1.069)	<0.0001
Educational level	≤Elementary	0.762	(0.678–0.856)	0.003
	Middle	0.903	(0.788–1.034)	0.190
	High	0.998	(0.897–1.11)	0.660
	≥College	1	Reference	
Smoking		1.349	(1.221–1.49)	<0.0001
Exercise		0.892	(0.821–0.97)	<0.0001
Drinking		1.057	(0.966–1.156)	<0.0001
Breakfast frequency	5–7/week	1	Reference	
	3–4/week	1.104	(0.952–1.28)	0.545
	1–2/week	1.301	(1.123–1.508)	0.245
	Rare	1.165	(1.016–1.337)	0.029
Lunch frequency	5–7/week	1	Reference	
	3–4/week	1.153	(0.946–1.406)	0.312
	1–2/week	1.330	(0.943–1.876)	0.791
	Rare	1.415	(1.005–1.992)	0.046
Dinner frequency	5–7/week	1	Reference	
	3–4/week	1.034	(0.878–1.218)	0.692
	1–2/week	0.833	(0.548–1.267)	0.897
	Rare	0.882	(0.407–1.912)	0.751
Eating out	≥2/day	1.250	(1.006–1.552)	
	1/day	1.103	(0.906–1.343)	
	5–6/week	0.899	(0.739–1.094)	
	3–4/week	0.971	(0.78–1.208)	
	1–2/week	1.008	(0.833–1.22)	
	1–3/month	1.064	(0.868–1.304)	
	Rare	1	Reference	

Table 2. Cont.

Characteristics		OR	(95% CI)	<i>p</i> -Value ^a
Companion for breakfast	Yes	1	Reference	0.001
	No	1.008	(0.903–1.126)	
Companion for lunch	Yes	1	Reference	0.018
	No	1.029	(0.932–1.136)	
Companion for dinner	Yes	1	Reference	0.335
	No	0.958	(0.857–1.07)	
WC		1.237	(1.227–1.247)	<0.0001
BMI		2.389	(2.309–2.471)	<0.0001
Fasting glucose		1.021	(1.018–1.023)	<0.0001
HbA _{1c}		1.829	(1.707–1.96)	<0.0001
TC		1.008	(1.006–1.009)	<0.0001
HDL-C		0.948	(0.944–0.952)	<0.0001
TG		1.004	(1.003–1.005)	<0.0001
LDL-C		1.008	(1.005–1.011)	<0.0001
Stress recognition		1.316	(1.203–1.44)	<0.0001
EQ-5D		2.255	(1.533–3.317)	<0.0001

Abbreviations: HRQoL, health-related quality of life; NAFLD, Non-alcoholic fatty liver disease; WC, waist circumference; BMI, body-mass index; HbA_{1c}, hemoglobin A_{1c}; TC, total cholesterol; HDL-C, high-density lipoprotein cholesterol; TG, triglyceride; LDL-C, low-density lipoprotein cholesterol; Stress, if '1' is recorded for the stress question; EQ-5D, EuroQoL- 5 Dimension; OR, odds ratio; CI, confidence interval. Definitions: Age, 10-year increase; Smoking, the percentage of individuals who is currently smoking (within two year); alcohol, the percentage of alcohol intake more than once a month in the last year; breakfast (lunch and dinner) frequency, breakfast (lunch and dinner) dinner frequency for 1 week in the last year. ^a The *p*-value was determined through the complex sample logistic regression test.

Table 3. Association between NAFLD and dietary habits, stress, and HRQoL.

		OR	(95% CI)	<i>p</i> -Value ^a
Breakfast frequency	5–7/week	1	Reference	0.925
	3–4/week	1.052	0.890–1.243	
	1–2/week	1.246	1.053–1.474	
	Rare	1.042	0.890–1.221	
Lunch frequency	5–7/week	1	Reference	0.399
	3–4/week	1.116	0.903–1.378	
	1–2/week	1.291	0.907–1.836	
	Rare	1.333	0.940–1.890	
Dinner frequency	5–7/week	1	Reference	0.313
	3–4/week	1.025	0.852–1.233	
	1–2/week	0.861	0.563–1.317	
	Rare	0.653	0.278–1.533	
Stress		1.314	1.175–1.469	<0.0001
EQ-5D		3.028	1.893–4.844	<0.0001

Abbreviations: HRQoL, health-related quality of life; NAFLD, Non-alcoholic fatty liver disease; EQ-5D, EuroQoL- 5 Dimension; OR, odds ratio; CI, confidence interval. Definitions: Age, 10-year increase; breakfast (lunch and dinner) frequency, breakfast (lunch and dinner) dinner frequency for 1 week in the last year; Stress, if '1' is recorded for the stress question. Adjusted for sex, age, education level, smoking, and exercise. ^a The ORs and 95% CI are determined through the complex sample logistic regression test.

4. Discussion

This study investigated the association between NAFLD occurrence and dietary habits, stress, and HRQoL among Korean adults. In particular, stress and HRQoL were associated with NAFLD. Herein, the HSI was considered for diagnosing NAFLD, since it is a simple and effective diagnostic index often used among Korean individuals. In a cohort study, a logistic regression model was performed using high serum ALT to serum AST ratio, high BMI, and diabetes mellitus as variables, which are independent risk factors for NAFLD. HSI was shown to have an AUROC of 0.812 [39]. Another study reported that the optimal cut-off value of HSI was 35 (sensitivity, 81%; specificity, 60%) [43]. Several studies have also been conducted considering HSI > 36 to predict NAFLD; we used the same definition in the current study.

Numerous studies have investigated the association between NAFLD and nutrient intake, lack of exercise, and carbohydrate intake [44–49]; however, few studies have focused on the pattern of dietary habits and NAFLD, or have reported conflicting findings [50–52]. To determine whether breakfast consumption influences NAFLD risk, similar to weight loss, it is necessary to assess its association with meal patterns. Herein, the NAFLD risk was lower upon daily breakfast consumption; however, it appeared irrelevant after correcting for the relevant variables, suggesting that total calories, daily carbohydrate intake, and the lack of essential nutrients are better associated with NAFLD than with dietary habits including the frequency of breakfast consumption, dinner consumption, and the frequency of eating out.

The relevance of breakfast in metabolic diseases and obesity has been adequately clarified [10–12,53,54]. Furthermore, the frequency of breakfast consumption is low among obese children with a high-energy intake at dinner [55]. The reduction in breakfast consumption in obesity has emerged as a global trend [11]. People do not have breakfast and cannot control their appetite, and so they binge. However, in the current study, the frequency of breakfast consumption and NAFLD risk in Korean subjects were not associated after correcting for related variables. A previous study has examined the composition of meals and NAFLD by investigating the connection between the Mediterranean diet and NAFLD [56]. The study involved 584 subjects with one or more cardiovascular disease risk factors. In the study, NAFLD diagnosis was performed by ultrasound, and adherence to the Mediterranean diet was classified as low, intermediate, or high. The study results indicated that the intermediate and high adherence groups had a lower risk of developing NAFLD than the low adherence group. The study examined the composition of the meal and diagnosed NAFLD using ultrasound, which is different from our study.

Although further studies are required, the following mechanisms could potentially promote the pathogenesis of NAFLD. Consumption of high-energy foods stimulates insulin secretion to regulate blood glucose levels. Insulin resistance results in fat breakdown in peripheral fat tissue to increase the generation of free fatty acids, which eventually increases the intrahepatic influx of free fatty acids. Furthermore, insulin increases fat biosynthesis in the liver, thus influencing the pathogenesis of fatty liver [1–3]. There is also a contribution of diet to NAFLD, as opposed to increasing insulin. The results of a mouse study show that intermittent fasting is effective for NAFLD [57]. Intermittent fasting in obese diet mice reduced hepatic lipogenesis and reduced β -oxidation markers to alleviate hepatic steatosis and inflammation.

Herein, the pathophysiological basis is defined as the interaction between psychological status, insulin resistance-related metabolic disorders, and NAFLD [16–20]. Psychological issues are associated with metabolic diseases via the accumulation of visceral fat through hormonal dysregulation, including that of insulin. Psychological factors including stress and depression promote corticotrophin-releasing hormone and cortisol secretion through the hypothalamic-pituitary-adrenal (HPA) axis [16–20]. Dysregulation of the HPA axis leads to the accumulation of visceral fat through hormone secretion and the secretion of inflammatory cytokines like tumor necrosis factor (TNF) [16–20,58]. Inflammatory substances undergo a vicious cycle that causes the relapse of metabolic abnormalities, again through insulin resistance and leptin disruptions [58]. These metabolic abnormalities are a major factor

accelerating NAFLD progression [1–3]. This hypothesis suggests a direct and indirect link between psychological factors and NAFLD, depending on personal susceptibility.

There have been mouse studies reporting that chronic stress has a negative effect on NAFLD [59]. It was reported that hepatic *TNF α* , *MCP-1*, and *HMOX* mRNA expression levels were high in chronic stress-exposed mice, which was attributed to accelerated liver oxidative stress and inflammatory mechanisms [59]. Studies have revealed that NAFLD is a stress-sensitive disease, as an alteration of the HPA axis exacerbates NAFLD [60,61]. Chronic psychiatric problems are closely related to NAFLD and obesity, and some studies report that psychiatric treatment must be considered when treating NAFLD [61]. This is consistent with the results of the current study, which suggest that stress management is necessary for NAFLD treatment.

NAFLD itself results in other metabolic diseases and is associated with stress and depression. Moreover, its treatment and clinical management are costly [62]. This is speculated to result in the deterioration of the HRQoL in NAFLD, and is supported by our results. Although not presented in a table, additional analysis in the current study showed that EQ-5D had a negative correlation with stress index, which is expected because EQ-5D evaluates various quality of life parameters, including stress.

A previous study has shown that NAFLD leads to economic losses, as shown by health economics, a tool to measure the economic and social consequences of the disease [63]. In addition, various metabolic conditions that can accompany NAFLD can lead to poor HRQoL [64–66]. Conversely, people with poor HRQoL are more likely to develop chronic diseases, especially metabolic diseases [64,65,67,68]. These studies, along with our current study, suggest that NAFLD is closely related to HRQoL.

The EQ-5D used in this study as an HRQoL assessment tool has already been demonstrated in Korean studies [69]. EQ-5D was developed by the EuroQoL group for the purpose of clinical and economic evaluation. It is a simple and widely used tool for assessing overall HRQoL. QoL scores have been previously reported in NAFLD including NASH (non-alcoholic steatohepatitis) and cirrhosis in a previous study, which investigated the association between NAFLD and QoL using the Non-alcoholic Steatohepatitis Clinical Research Network and the Short-Form 36 (SF-36) survey [26]. Previous studies investigated the association between NAFLD severity and QoL and reported a poor QoL among individuals with NAFLD [26,62], consistent with the present study. A previous study compared the relative effects of NAFLD on the HRQoL with those of other chronic liver diseases. NAFLD patients had a significantly lower HRQoL than hepatitis B or hepatitis C patients, as revealed through multiple domains of scores of the 29-item Chronic Liver Disease Questionnaire, suggesting that the HRQoL was severely impaired in NAFLD patients [24]. Herein, the EQ-5D indicator was used to evaluate the HRQoL. The closer this score is to 1 point, the higher the QoL [70,71]. Nam et al. (2007) used the EQ-5D to estimate the weightage of the QoL questionnaire, and this indicator was extensively used to assess the QoL among Korean individuals [71]. Furthermore, herein, the probability of NAFLD occurrence was 2.255-fold among the HRQoL indicators after correcting for related variables. Multiple approaches should be considered for the clinical management of NAFLD for improving stress and the HRQoL should be accompanied by classical treatments including pharmacotherapy, lifestyle improvement, and diet.

This study has some limitations. First, since this study included cross-sectional data, it is difficult to explain the causal relationship between stress dietary habits, HRQoL, and NAFLD. Second, the gold-standard diagnosis of NAFLD is liver biopsy and abdominal ultrasound; however, it was difficult to conduct these procedures when surveying the entire population. Furthermore, although the FLI is a commonly used NAFLD indicator, it was difficult to use here because γ -glutamyl transferase levels were not determined. Third, all other variables potentially affecting NAFLD were not considered herein. Despite these limitations, this study has the following strengths. First, this study delineated the importance of psychological aspects by elucidating the association between stress and NAFLD. Second, the number of breakfast, lunch, and dinner meals per week as a dietary habit was not associated with NAFLD. In the future, more specific dietary habits should be investigated to confirm the association.

Most importantly, this study shows that the HRQoL is associated with NAFLD among Korean adults. These results may provide essential strategies to facilitate the treatment of NAFLD.

Based on the results of this study, future studies should investigate these associations using more accurate tools for NAFLD diagnosis, while evaluating and comparing HRQoL using various tools. These additional studies may provide evidence for confirming the causal relationship between NAFLD and quality of life.

Funding: This research was funded by Wonkwang University.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Hassan, K.; Bhalla, V.; El Regal, M.E.; A-Kader, H.H. Nonalcoholic fatty liver disease: A comprehensive review of a growing epidemic. *World J. Gastroenterol. WJG* **2014**, *20*, 12082. [[CrossRef](#)]
2. Sozio, M.S.; Liangpunsakul, S.; Crabb, D. The role of lipid metabolism in the pathogenesis of alcoholic and nonalcoholic hepatic steatosis. In *Seminars in Liver Disease*; Thieme Medical Publishers: New York, NY, USA, 2010.
3. Jun, D.W. The role of diet in non-alcoholic fatty liver disease. *Korean J. Gastroenterol.* **2013**, *61*, 243–251. [[CrossRef](#)] [[PubMed](#)]
4. Singal, A.K.; Hasanin, M.; Kaif, M.; Wiesner, R.; Kuo, Y.F. Nonalcoholic steatohepatitis is the most rapidly growing indication for simultaneous liver kidney transplantation in the United States. *Transplantation* **2016**, *100*, 607–612. [[CrossRef](#)]
5. Yasutake, K.; Kohjima, M.; Kotoh, K.; Nakashima, M.; Nakamuta, M.; Enjoji, M. Dietary habits and behaviors associated with nonalcoholic fatty liver disease. *World J. Gastroenterol. WJG* **2014**, *20*, 1756. [[CrossRef](#)] [[PubMed](#)]
6. Toshimitsu, K.; Matsuura, B.; Ohkubo, I.; Niiya, T.; Furukawa, S.; Hiasa, Y.; Kawamura, M.; Ebihara, K.; Onji, M. Dietary habits and nutrient intake in non-alcoholic steatohepatitis. *Nutrition* **2007**, *23*, 46–52. [[CrossRef](#)] [[PubMed](#)]
7. Abdelmalek, M.F.; Suzuki, A.; Guy, C.; Unalp-Arida, A.; Colvin, R.; Johnson, R.J.; Diehl, A.M. Nonalcoholic Steatohepatitis Clinical Research Network Increased fructose consumption is associated with fibrosis severity in patients with nonalcoholic fatty liver disease. *Hepatology* **2010**, *51*, 1961–1971. [[CrossRef](#)]
8. Fuse, Y.; Hirao, A.; Kuroda, H.; Otsuka, M.; Tahara, Y.; Shibata, S. Differential roles of breakfast only (one meal per day) and a bigger breakfast with a small dinner (two meals per day) in mice fed a high-fat diet with regard to induced obesity and lipid metabolism. *J. Circadian Rhythm.* **2012**, *10*, 4. [[CrossRef](#)]
9. Stunkard, A.J.; Grace, W.J.; Wolff, H.G. The night-eating syndrome: A pattern of food intake among certain obese patients. *Am. J. Med.* **1955**, *19*, 78–86. [[CrossRef](#)]
10. Summerbell, C.D.; Moody, R.C.; Shanks, J.; Stock, M.J.; Geissler, C. Relationship between feeding pattern and body mass index in 220 free-living people in four age groups. *Eur. J. Clin. Nutr.* **1996**, *50*, 513–519.
11. Timlin, M.T.; Pereira, M.A. Breakfast frequency and quality in the etiology of adult obesity and chronic diseases. *Nutr. Rev.* **2007**, *65*, 268–281. [[CrossRef](#)]
12. Wyatt, H.R.; Grunwald, G.K.; Mosca, C.L.; Klem, M.L.; Wing, R.R.; Hill, J.O. Long-term weight loss and breakfast in subjects in the National Weight Control Registry. *Obes. Res.* **2002**, *10*, 78–82. [[CrossRef](#)] [[PubMed](#)]
13. Lei, S.; Liu, Z.W.; Li, Y.; Gong, C.; Zhang, H.; Song, L.J.; Huang, C.Y.; Li, M. The prevalence of nonalcoholic fatty liver disease and its association with lifestyle/dietary habits among university faculty and staff in Chengdu. *Biomed. Environ. Sci.* **2012**, *25*, 383–391.
14. Chung, G.E. Dietary patterns are associated with the prevalence of nonalcoholic fatty liver disease in Korean adults. *Nutrition* **2019**, *62*, 32–38. [[CrossRef](#)]
15. Chang, J.H.; Lee, H.S.; Kang, E.H. A study on dietary habits, nutrient intakes and dietary quality in adults of a health screening and promotion center according to non-alcoholic fatty liver disease. *J. Nutr. Health* **2014**, *47*, 330–341. [[CrossRef](#)]
16. Chung, G.E.; Youn, J.; Kim, Y.S.; Lee, J.E.; Yang, S.Y.; Lim, J.H.; Song, J.H.; Doo, E.Y.; Kim, J.S. Non-Alcoholic Fatty Liver Disease and Its Association with Depression in Korean General Population. *J. Korean Med. Sci.* **2019**, *34*. [[CrossRef](#)]

17. Surdea-Blaga, T.; Dumitraşcu, D. Depression and anxiety in nonalcoholic steatohepatitis: Is there any association? *Rom. J. Intern. Med. Rev. Roum. Med. Interne* **2011**, *49*, 273–280.
18. Russ, T.C.; Kivimäki, M.; Morling, J.R.; Starr, J.M.; Stamatakis, E.; Batty, G.D. Association between psychological distress and liver disease mortality: A meta-analysis of individual study participants. *Gastroenterology* **2015**, *148*, 958–966. [[CrossRef](#)]
19. Nagaraja, A.S.; Sadaoui, N.C.; Dorniak, P.L.; Lutgendorf, S.K.; Sood, A.K. SnapShot: Stress and disease. *Cell Metab.* **2016**, *23*, 388.e1. [[CrossRef](#)]
20. Lundberg, U. Stress hormones in health and illness: The roles of work and gender. *Psychoneuroendocrinology* **2005**, *30*, 1017–1021. [[CrossRef](#)]
21. Schneiderman, N.; Ironson, G.; Siegel, S.D. Stress and health: Psychological, behavioral, and biological determinants. *Annu. Rev. Clin. Psychol.* **2005**, *1*, 607–628. [[CrossRef](#)]
22. Björntorp, P. Visceral fat accumulation: The missing link between psychosocial factors and cardiovascular disease? *J. Inter. Med.* **1991**, *230*, 195–201. [[CrossRef](#)] [[PubMed](#)]
23. Yudkin, J.S.; Kumari, M.; Humphries, S.E.; Mohamed-Ali, V. Inflammation, obesity, stress and coronary heart disease: Is interleukin-6 the link? *Atherosclerosis* **2000**, *148*, 209–214. [[CrossRef](#)]
24. Dan, A.A.; Kallman, J.B.; Wheeler, A.; Younoszai, Z.; Collantes, R.; Bodini, S.; Gerber, L.; Younoszai, Z.M. Health-related quality of life in patients with non-alcoholic fatty liver disease. *Aliment. Pharmacol. Ther.* **2007**, *26*, 815–820. [[CrossRef](#)] [[PubMed](#)]
25. Assimakopoulos, K.; Karaivazoglou, K.; Tsermpini, E.E.; Diamantopoulou, G.; Triantos, C. Quality of life in patients with nonalcoholic fatty liver disease: A systematic review. *J. Psychosom. Res.* **2018**, *112*, 73–80. [[CrossRef](#)]
26. David, K.; Kowdley, K.V.; Unalp, A.; Kanwal, F.; Brunt, E.M.; Schwimmer, J.B. Quality of life in adults with nonalcoholic fatty liver disease: Baseline data from the nonalcoholic steatohepatitis clinical research network. *Hepatology* **2009**, *49*, 1904–1912. [[CrossRef](#)]
27. Kennedy-Martin, T.; Bae, J.P.; Paczkowski, R.; Freeman, E. Health-related quality of life burden of nonalcoholic steatohepatitis: A robust pragmatic literature review. *J. Patient Rep. Outcomes* **2018**, *2*, 28. [[CrossRef](#)]
28. Tsuchiya, A.; Ikeda, S.; Ikegami, N.; Nishimura, S.; Sakai, I.; Fukuda, T.; Hamashima, C.; Hisashige, A.; Tamura, M. Estimating an EQ-5D population value set: The case of Japan. *Health Econ.* **2002**, *11*, 341–353. [[CrossRef](#)]
29. Balestroni, G.; Bertolotti, G. EuroQol-5D (EQ-5D): An instrument for measuring quality of life. *Monaldi Arch. Chest Dis.* **2012**, *78*, 155–159. [[CrossRef](#)]
30. Dolan, P.; Gudex, C.; Kind, P.; Williams, A. *A Social Tariff for EuroQol: Results from a UK General Population Survey*; University of York, Center for Health Economics: Heslington, UK, 1995.
31. Luo, N.; Johnson, J.A.; Shaw, J.W.; Feeny, D.; Coons, S.J. Self-reported health status of the general adult US population as assessed by the EQ-5D and Health Utilities Index. *Med. Care* **2005**, *43*, 1078–1086. [[CrossRef](#)]
32. Han, M.A.; Ryu, S.Y.; Park, J.; Kang, M.; Park, J.K.; Kim, K.S. Health-related Quality of Life Assessment by the EuroQol-5D in Some Rural Adults. *J. Prev. Med. Public Health* **2008**, *41*, 173–180. [[CrossRef](#)]
33. Jo, M.-W.; Yun, S.C.; Lee, S.I. Estimating quality weights for EQ-5D health states with the time trade-off method in South Korea. *Value Health* **2008**, *11*, 1186–1189. [[CrossRef](#)] [[PubMed](#)]
34. Kang, E.J.; Shin, H.S.; Park, H.J.; Jo, M.W.; Kim, N.Y. A valuation of health status using EQ-5D. *Korean J. Health Econ. Policy* **2006**, *12*, 19–43.
35. Papagianni, M.; Sofogianni, A.; Tziomalos, K. Non-invasive methods for the diagnosis of nonalcoholic fatty liver disease. *World J. Hepatol.* **2015**, *7*, 638. [[CrossRef](#)] [[PubMed](#)]
36. Bateman, R.M.; Sharpe, M.D.; Jagger, J.E.; Ellis, C.G.; Solé-Violán, J.; López-Rodríguez, M.; Herrera-Ramos, E.; Ruíz-Hernández, J.; Borderías, L.; Horcajada, J.; et al. Development, external validation, and comparative assessment of a new diagnostic score for hepatic steatosis. *Am. J. Gastroenterol.* **2014**, *109*, 1404.
37. Shen, Y.N.; Yu, M.X.; Gao, Q.; Li, Y.Y.; Huang, J.J.; Sun, C.M.; Qiao, N.; Zhang, H.X.; Wang, H.; Lu, Q.; et al. External validation of non-invasive prediction models for identifying ultrasonography-diagnosed fatty liver disease in a Chinese population. *Medicine* **2017**, *96*, e7610. [[CrossRef](#)]
38. Zhang, Z.; Wang, G.; Kang, K.; Wu, G.; Wang, P. Diagnostic accuracy and clinical utility of a new noninvasive index for hepatic steatosis in patients with hepatitis B virus infection. *Sci. Rep.* **2016**, *6*, 32875. [[CrossRef](#)]

39. Lee, J.H.; Kim, D.; Kim, H.J.; Lee, C.H.; Yang, J.I.; Kim, W.; Kim, Y.J.; Yoon, J.H.; Cho, S.H.; Sung, M.W.; et al. Hepatic steatosis index: A simple screening tool reflecting nonalcoholic fatty liver disease. *Dig. Liver Dis.* **2010**, *42*, 503–508. [[CrossRef](#)]
40. Lee, J.; Ha, J.; Jo, K.; Lim, D.J.; Lee, J.M.; Chang, S.A.; Kang, M.I.; Cha, B.Y.; Kim, M.H. Male-specific association between subclinical hypothyroidism and the risk of non-alcoholic fatty liver disease estimated by hepatic steatosis index: Korea National Health and Nutrition Examination Survey 2013 to 2015. *Sci. Rep.* **2018**, *8*, 1–7. [[CrossRef](#)]
41. Kweon, S.; Kim, Y.; Jang, M.J.; Kim, Y.; Kim, K.; Choi, S.; Chun, C.; Khang, Y.H.; Oh, K. Data resource profile: The Korea national health and nutrition examination survey (KNHANES). *Int. J. Epidemiol.* **2014**, *43*, 69–77. [[CrossRef](#)]
42. Craig, C.L.; Marshall, A.L.; Sjöström, M.; Bauman, A.E.; Booth, M.L.; Ainsworth, B.E.; Pratt, M.; Ekelund, U.; Yngve, A.; Sallis, J.F.; et al. International physical activity questionnaire: 12-country reliability and validity. *Med. Sci. Sports Exerc.* **2003**, *35*, 1381–1395. [[CrossRef](#)]
43. Chen, L.D.; Huang, J.F.; Chen, Q.S.; Lin, G.F.; Zeng, H.X.; Lin, X.F.; Lin, X.J.; Lin, L.; Lin, Q.C. Validation of fatty liver index and hepatic steatosis index for screening of non-alcoholic fatty liver disease in adults with obstructive sleep apnea hypopnea syndrome. *Chin. Med. J.* **2019**, *132*, 2670–2676. [[CrossRef](#)] [[PubMed](#)]
44. Da Silva, H.E.; Arendt, B.M.; Noureldin, S.A.; Therapondos, G.; Guindi, M.; Allard, J.P. A cross-sectional study assessing dietary intake and physical activity in Canadian patients with nonalcoholic fatty liver disease vs healthy controls. *J. Acad. Nutr. Diet.* **2014**, *114*, 1181–1194. [[CrossRef](#)] [[PubMed](#)]
45. Kris-Etherton, P.M.; Pearson, T.A.; Wan, Y.; Hargrove, R.L.; Moriarty, K.; Fishell, V.; Etherton, T.D. High-monounsaturated fatty acid diets lower both plasma cholesterol and triacylglycerol concentrations. *Am. J. Clin. Nutr.* **1999**, *70*, 1009–1015. [[CrossRef](#)] [[PubMed](#)]
46. Zelber-Sagi, S.; Nitzan-Kaluski, D.; Goldsmith, R.; Webb, M.; Blendis, L.; Halpern, Z.; Oren, R. Long term nutritional intake and the risk for non-alcoholic fatty liver disease (NAFLD): A population based study. *J. Hepatol.* **2007**, *47*, 711–717. [[CrossRef](#)] [[PubMed](#)]
47. Ganji, S.H.; Kashyap, M.L.; Kamanna, V.S. Niacin inhibits fat accumulation, oxidative stress, and inflammatory cytokine IL-8 in cultured hepatocytes: Impact on non-alcoholic fatty liver disease. *Metabolism* **2015**, *64*, 982–990. [[CrossRef](#)] [[PubMed](#)]
48. Ullah, R.; Rauf, N.; Nabi, G.; Ullah, H.; Shen, Y.; Zhou, Y.D.; Fu, J. Role of nutrition in the pathogenesis and prevention of non-alcoholic fatty liver disease: Recent updates. *Int. J. Biol. Sci.* **2019**, *15*, 265. [[CrossRef](#)] [[PubMed](#)]
49. Yang, H.Y.; Tzeng, Y.H.; Chai, C.Y.; Hsieh, A.T.; Chen, J.R.; Chang, L.S.; Yang, S.S. Soy protein retards the progression of non-alcoholic steatohepatitis via improvement of insulin resistance and steatosis. *Nutrition* **2011**, *27*, 943–948. [[CrossRef](#)]
50. Trovato, F.M.; Martines, G.F.; Brischetto, D.; Catalano, D.; Musumeci, G.; Trovato, G.M. Fatty liver disease and lifestyle in youngsters: Diet, food intake frequency, exercise, sleep shortage and fashion. *Liver Int.* **2016**, *36*, 427–433. [[CrossRef](#)]
51. Koot, B.G.; van der Baan-Slootweg, O.H.; Tamminga-Smeulders, C.L.; Rijcken, T.H.; Korevaar, J.C.; van Aalderen, W.M.; Jansen, P.L.; Benninga, M.A. Lifestyle intervention for non-alcoholic fatty liver disease: Prospective cohort study of its efficacy and factors related to improvement. *Arch. Dis. Child.* **2011**, *96*, 669–674. [[CrossRef](#)]
52. Nseir, W.; Hellou, E.; Assy, N. Role of diet and lifestyle changes in nonalcoholic fatty liver disease. *World J. Gastroenterol. WJG* **2014**, *20*, 9338.
53. Cho, S.; Dietrich, M.; Brown, C.J.; Clark, C.A.; Block, G. The effect of breakfast type on total daily energy intake and body mass index: Results from the Third National Health and Nutrition Examination Survey (NHANES III). *J. Am. Coll. Nutr.* **2003**, *22*, 296–302. [[CrossRef](#)] [[PubMed](#)]
54. Song, W.O.; Chun, O.K.; Obayashi, S.; Cho, S.; Chung, C.E. Is consumption of breakfast associated with body mass index in US adults? *J. Am. Diet. Assoc.* **2005**, *105*, 1373–1382. [[CrossRef](#)] [[PubMed](#)]
55. Moreno, L.A.; Rodriguez, G. Dietary risk factors for development of childhood obesity. *Curr. Opin. Clin. Nutr. Metab. Care* **2007**, *10*, 336–341. [[CrossRef](#)]
56. Baratta, F.; Pastori, D.; Polimeni, L.; Bucci, T.; Ceci, F.; Calabrese, C.; Ernesti, I.; Pannitteri, G.; Violi, F.; Angelico, F.; et al. Adherence to mediterranean diet and non-alcoholic fatty liver disease: Effect on insulin resistance. *Am. J. Gastroenterol.* **2017**, *112*, 1832–1839. [[CrossRef](#)] [[PubMed](#)]

57. De Souza Marinho, T.; Ornellas, F.; Barbosa-da-Silva, S.; Mandarim-de-Lacerda, C.A.; Aguilu, M.B. Beneficial effects of intermittent fasting on steatosis and inflammation of the liver in mice fed a high-fat or a high-fructose diet. *Nutrition* **2019**, *65*, 103–112. [[CrossRef](#)]
58. Hotamisligil, G.S.; Shargill, N.S.; Spiegelman, B.M. Adipose expression of tumor necrosis factor-alpha: Direct role in obesity-linked insulin resistance. *Science* **1993**, *259*, 87–91. [[CrossRef](#)]
59. Czech, B.; Neumann, I.D.; Müller, M.; Reber, S.O.; Hellerbrand, C. Effect of chronic psychosocial stress on nonalcoholic steatohepatitis in mice. *Int. J. Clin. Exp. Pathol.* **2013**, *6*, 1585. [[PubMed](#)]
60. Zhang, S.; Ma, C.; Wang, X.; Wang, Q.; Song, W.; Li, C.; Zhai, C.; Qi, Y.; Fan, S.; Cheng, F. Impact of chronic psychological stress on nonalcoholic fatty liver disease. *Int. J. Clin. Exp. Med.* **2019**, *12*, 7991–7998.
61. Stewart, K.E.; Levenson, J.L. Psychological and psychiatric aspects of treatment of obesity and nonalcoholic fatty liver disease. *Clin. Liver Dis.* **2012**, *16*, 615–629. [[CrossRef](#)] [[PubMed](#)]
62. Tapper, E.B.; Sengupta, N.; Hunink, M.G.; Afdhal, N.H.; Lai, M. Cost-effective evaluation of nonalcoholic fatty liver disease with NAFLD fibrosis score and vibration controlled transient elastography. *Am. J. Gastroenterol.* **2015**, *110*, 1298–1304. [[CrossRef](#)]
63. Cortesi, P.; Scalone, L.; Ciampichini, R.; Cozzolino, P.; Cesana, G.; Mantovani, L.; Okolicsanyi, S.; Ciaccio, A.; Rota, M.; Gentiluomo, M.; et al. Health related quality of life in the major liver conditions. *Hepatology* **2013**, *58* (Suppl. 1), 1210A.
64. Fontaine, K.; Barofsky, I. Obesity and health-related quality of life. *Obes. Rev.* **2001**, *2*, 173–182. [[CrossRef](#)] [[PubMed](#)]
65. Swallen, K.C.; Reither, E.N.; Haas, S.A.; Meier, A.M. Overweight, obesity, and health-related quality of life among adolescents: The National Longitudinal Study of Adolescent Health. *Pediatrics* **2005**, *115*, 340–347. [[CrossRef](#)] [[PubMed](#)]
66. Hassan, M.K.; Joshi, A.V.; Madhavan, S.S.; Amonkar, M.M. Obesity and health-related quality of life: A cross-sectional analysis of the US population. *Int. J. Obes.* **2003**, *27*, 1227–1232. [[CrossRef](#)]
67. Jia, H.; Lubetkin, E.I. The impact of obesity on health-related quality-of-life in the general adult US population. *J. Public Health* **2005**, *27*, 156–164. [[CrossRef](#)]
68. Ford, E.S.; Li, C. Metabolic syndrome and health-related quality of life among US adults. *Ann. Epidemiol.* **2008**, *18*, 165–171. [[CrossRef](#)]
69. Kim, M.H.; Cho, Y.S.; Uhm, W.S.; Kim, S.; Bae, S.C. Cross-cultural adaptation and validation of the Korean version of the EQ-5D in patients with rheumatic diseases. *Qual. Life Res.* **2005**, *14*, 1401–1406. [[CrossRef](#)]
70. Herdman, M.; Gudex, C.; Lloyd, A.; Janssen, M.F.; Kind, P.; Parkin, D.; Bonse, G.; Badia, X. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual. Life Res.* **2011**, *20*, 1727–1736. [[CrossRef](#)]
71. Nam, H.; Kim, K.; Kwon, S.; Koh, K.; Poul, K.; Ko, K.W.; Kind, P. *EQ-5D Korean Valuation Study Using Time Trade Off Method*; ScienceOpen, Inc.: Burlington, MA, USA, 2011.

