Table S1. LLD score-2018 and food components in GI cancers survivors and reference population, Lifelines 2006-2009

	GI cancers survivors (n=307) Mean (SD)	Reference population (n=3070) Mean (SD)	(P-value)2
LDS	25.71 (5.92)	25.92 (5.91)	0.35
esired food component			
Vegetables	102.51 (54.93)	107.39 (57.51)	0.05
Fruits	180.85 (115.53)	173.06 (126.94)	0.45
Whole grain products	101.69 (51.57)	109.28 (53.39)	0.11
Legumes and Nuts	24.06 (26.49)	22.85 (22.26)	0.72
Fish	17.67 (17.81)	14.52 (13.56)	0.80
Oil and soft margarine	15.03 (16.40)	16.07 (16.99)	0.71
Unsweetened dairy	210.25 (193.76)	194.41 (157.37)	0.91
Coffee	440.14 (226.37)	460.63 (234.53)	0.46
Tea	266.80 (227.80)	251.41 (230.31)	0.00
Indesired food component			
Red and Processed meat	61.54 (33.46)	62.65 (32.08)	0.09
Butter, hard margarine	31.26 (24.39)	29.81 (23.53)	0.33
Sugar sweetened beverages	109.51 (157.24)	96.13 (133.16)	0.26
Daily alcohol intake	7.49 (8.88)	7.77 (9.09)	0.41

Abbreviations: CI, confidence interval; LLD, life Line diet; MD, mean difference; SD, standard deviation.

¹All food components are presented in mean intake in grams per day.
²Results from a multivariate linear regression adjusted for age, sex, smoking, BMI, education level and physical




Figure S1: Mean LLD score according to age and sex category among GI cancers survivors stratified by time since diagnosis, Lifeline Study (2006-2009).

Abbreviations: GI, gastrointestinal; LLD, life Line diet.




Figure S2. Mean LLD score according to age and sex category among GI cancers survivors stratified by tumour site, Lifeline Study (2006-2009).