

Supplementary Material

Supplementary Table S1. Nucleotide sequences of primers used for real time quantitative PCR.

Gene	Forward Primer (5' to 3')	Reverse Primer (5' to 3')
<i>Acc1</i>	TGCAGGTATCCCCACTCTTC	TTCTGATTCCCTTCCCTCCT
<i>Cd36</i>	GTCCTGGCTGTGTTGGA	GCTCAAAGATGGCTCCATTG
<i>Cpt1α</i>	GCTCGCACATTACAAGGACAT	TGGACACCACATAGAGGCAG
<i>Fas1</i>	CTATTGTGGACGGAGGTATC	TGCTGTAGCCCAGAAGAG
<i>Fatp5</i>	CCTGCCAAGCTTCTGCTTAAT	GCTCATGTGATAGGATGGCTGG
<i>Had</i>	ATCGTGAACCGTCTCTTGGT	AGGACTGGGCTGAAATAAGG
<i>Srebp-1c</i>	CCCACCCCCCTTACACACC	GCCTGCGGTCTTCATTGT

The table shows the nucleotide sequences of primers used for PCR amplification. *Acc1*, acetyl CoA carboxylase; *Cd36*, fatty acid translocase, homologue of CD36; *Cpt1α*, carnitine palmitoyltransferase 1 alpha; *Fas1*, sterol regulatory element-binding protein 1; *Fatp5*, fatty acid transport protein 5; *Had*, hydroxyacyl-CoA dehydrogenase; *Srebp1c*, sterol regulatory element-binding protein 1c.