

Article

Effects of Cannabidiol Supplementation on Skeletal Muscle Regeneration after Intensive Resistance Training

Eduard Isenmann ^{1,2}, Sebastian Veit ¹, Lynn Starke ¹, Ulrich Flenker ¹ and Patrick Diel ^{1,*}

¹ Department of Molecular and Cellular Sports Medicine, Institute for Cardiovascular Research and Sports Medicine, German Sports University, 50933 Cologne, Germany; e.isenmann@dshs-koeln.de (E.I.); s.veit@dshs-koeln.de (S.V.); lynn@starke-team.de (L.S.); u.flenker@dshs-koeln.de (U.F.)

² Department of Fitness and Health, IST-University of Applied Sciences, 40233 Dusseldorf, Germany

* Correspondence: diel@dshs-koeln.de; Tel.: +49-221-4982-5860

Abstract: Cannabidiol supplements (CBD) are increasingly consumed by athletes to improve regeneration. However, the evidence for the pro-regenerative effects of CBD in sports is quite limited. Therefore, our aim was to investigate the effects of a single CBD supplementation in a six-arm placebo-controlled crossover study after resistance training on performance and muscle damage. Before and after the resistance training, one-repetition maximum in the back squat (1RM BS), countermovement jump (CMJ), and blood serum concentrations of creatine kinase (CK) and myoglobin (Myo) were measured in healthy, well-trained participants. Sixteen of 21 participants completed the study and were included in the analysis. In 1RM BS, a significant decrease was observed after 24 h ($p < 0.01$) but not 48 or 72 h. A significant group difference was detected after 72 h ($p < 0.05$; ES = 0.371). In CMJ, no significant changes were observed. The CK and Myo concentrations increased significantly after 24 h (CK: $p < 0.001$; Myo: $p < 0.01$), 48 h (CK: $p < 0.001$; Myo: $p < 0.01$) and 72 h (CK: $p < 0.001$; Myo: $p < 0.001$). After 72 h, a significant group difference was observed in both muscle damage biomarkers (CK: $p < 0.05$ ES = 0.236; Myo: $p < 0.05$; ES = 0.214). The results show small and significant effects on muscle damage and recovery of squat performance after 72 h. However, more data are necessary for clearer statements about the pro-regenerative effects of CBD supplementation after resistance training.

Keywords: cannabidiol; CBD; recovery; muscle damage; performance; strength training; resistance training



Citation: Isenmann, E.; Veit, S.; Starke, L.; Flenker, U.; Diel, P. Effects of Cannabidiol Supplementation on Skeletal Muscle Regeneration after Intensive Resistance Training. *Nutrients* **2021**, *13*, 3028. <https://doi.org/10.3390/nu13093028>

Academic Editor: Yosuke Yamada

Received: 3 August 2021

Accepted: 25 August 2021

Published: 30 August 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Intense training and performances result in damage and inflammation of the skeletal muscles [1–3]. Consequently, the performance of the trained skeletal muscles decreases after the intensive training sessions and needs time to recover [4–6]. In competitive sports, it is essential to minimize the recovery time to provide further training stimuli as quickly as possible and to increase performance. Dietary supplements are often used to achieve this goal [7–9]. Cannabidiol (CBD) products in particular are often being consumed by elite athletes [10–12]. CBD is a non-psychoactive compound of Cannabis Sativa [13]. Due to its structural differences to Δ^9 -tetrahydrocannabinol (Δ^9 -THC), it has different affinities to bind to the cannabinoid receptors 1 and 2 [14–17]. Mechanistically, CBD can inhibit the NF- κ B signaling pathway and activate JAK/STAT kinase [18]. Therefore, CBD is used in a variety of diseases such as autoimmune encephalitis [19], rheumatoid arthritis [20], colitis [21], diabetes [22], and psoriasis [23], as well as Alzheimer's [24], Parkinson's [25], and Huntington's Disease [26]. It has also anticonvulsant [14,27–29], antipsychotic [14,27,30], antiemetic [14,27], anticarcinogenic [31,32], and antidepressant [14,33] effects. Although regulations regarding its legality vary among countries worldwide [34,35], CBD was removed from the World Anti-Doping Agency (WADA) prohibited substances list in 2018 [36], based on evidence that CBD is safe and well-tolerated in humans [37–39]. While the effects

and associated mechanisms of CBD in the human body have been studied [40–42], data in the context of sports and performance are still limited. Previous research has shown that roughly 27–42% of CBD users state improvement of sleep as a reason for administration [43,44]. Improved sleep quality has been shown mainly in animal models [45,46] and not clearly in humans [47–49]. Additionally, first studies have shown a small effect on muscle soreness after CBD supplementation [50,51]. For the pro-regenerative effects of CBD in sport, there are currently only two studies that have investigated biomarkers and performance parameters. In untrained men, the intake of CBD oil after six sets of ten maximal eccentric isokinetic elbow flexions did not lead to any significant differences in perceived soreness, arm circumference, hanging joint angle, and peak torque between the groups [51]. In contrast, a pilot study [52] in highly trained men (>150% of body weight in the back squat [53]), showed an inhibitory effect of CBD on serum creatine kinase (CK) concentrations after three sets of 12 back squats at 70% 1RM plus three sets of 15 drop jumps landed in a deep squat. Consequently, based on the current lack of data, no definite conclusion can be drawn about the effects of CBD supplementation to promote regeneration processes after intensive strength training. Therefore, the aim of this study was to investigate the effects of a single CBD supplementation 24, 48, and 72 h after intensive strength training on muscle damage and performance. For this purpose, a placebo-controlled, double-blind, six-arm crossover study was conducted.

2. Martials and Methods

2.1. Participants

The study design was approved by the ethics committee of the German Sports University Cologne and conducted in accordance with the Declaration of Helsinki. All participants were informed about the study design and confirmed their voluntary participation in written form. A total of 21 subjects were recruited for the study. People who were ill, injured, or were taking medication or dietary supplements were excluded. All subjects were healthy and experienced (at least one year) in strength training with a back squat performance of at least 125% of their body mass [53]. A total of 16 subjects (Table 1) finished the study, while five dropped out due to reasons not related to the study. To confirm the performance requirements, maximum strength was tested prior to the investigation. During the actual days of the study as well as 24 h before, no further physical activities were allowed, the natural diet was left unchanged.

Table 1. Subjects information.

Parameter	$\mu \pm \sigma$
Age [Years]	24 $\pm$ 3
Height [cm]	181.4 $\pm$ 10.0
Body Mass [kg]	79.2 $\pm$ 13.7
CMJ [cm]	55.2 $\pm$ 8.5
1RM [kg]	118.5 $\pm$ 24.7
1RM/BM [kg]	1.4 $\pm$ 0.2

$\mu \pm \sigma$ = mean $\pm$ standard deviation; CMJ = Countermovement Jump; 1RM = One-Repetition Maximum; 1RM/BM = One-Repetition Maximum/Body Mass.

2.2. Study Design

The study was carried out as a randomized, double-blind study in a six-arm crossover design and compared two conditions after an intensive strength training: the experimental condition using CBD (CBD) and the control condition using a placebo (PLA). A washout period of two weeks was taken between all interventions. For neuromuscular measures, maximum strength and power were tested for the lower body. CK and Myoglobin (Myo) were used as biochemical measures. Pretests (T 0) were performed for all four parameters prior to each intervention. Additionally, the same tests were executed again in three different sessions 24 (T 24), 48 (T 48), and 72 (T 72) hours after a training protocol. This results

in three different PRE-POST interventions per condition for a total of six interventions (Figure 1). While the general order of conditions was the same, subjects were randomly assigned to start at one of six conditions according to age, gender, and maximal strength.




Figure 1. Study design. BS = Back Squat; reps = Repetitions; s = second; 1RM; One-Repetition Maximum; h = hours.

2.3. Testing Protocol

All subjects arrived at the facility in the morning rested and fasted for the initial examination. Anthropometric data were collected, and a blood sample was taken (T 0). Afterwards, subjects consumed a standardized breakfast consisting of 100 g of banana, 5 g of honey, and 40 g or 60 g of oatmeal for the women and men, respectively. Thus, nutritional values were about 270 kcal for female subjects and about 350 kcal for male subjects. After a 30-min break, the warm-up started with five minutes of running and five minutes of dynamic stretching. After the warm-up, maximal power and strength were tested. This testing protocol was performed both PRE (T 0) and POST training (T 24, T 48, T 72). It was based on the guidelines of the National Strength and Conditioning Association (NSCA) for performance testing [54].

2.4. Skeletal Muscle Serum Creatine Kinase (CK) and Myoglobin (Myo)

CK and Myo serum concentrations were determined for all time points until subjects performed their POST intervention using the COBAS h 232 Point-of-Care-System (Roche Diagnostic Systems, Rotkreuz, Switzerland). Therefore, the sample size decreases with each POST intervention as values after the POST intervention would be comparable.

2.5. Strength and Power 1RM Back Squat (1RM) and Counter Movement Jump (CMJ)

Maximal strength and power performance tests were conducted at each PRE (T 0) and POST (T 24, T 48, T 72) intervention. To measure maximal power output, subjects completed a jump and reach test (CMJ). Three jumps were averaged to determine the jump height. Maximal strength was measured as the one-repetition maximum (1RM) in the back squat. Subjects performed four warm-up sets, starting with 10 repetitions at 50% of their estimated 1RM, followed by eight repetitions at 60–70%, four–six repetitions at 70–85%, and, finally, two–four repetitions at 80–95%. All loads were rounded to the nearest 2.5 kg step. Subjects rested for two minutes between warm-up sets. Before the first 1RM attempt, another three minutes of rest was taken. Subjects performed up to four attempts to establish

the 1RM. The first attempts were taken at 90%, followed by a four-minute rest that was taken between all attempts.

2.6. Training Protocol

Following the T0 testing protocol, subjects performed a training protocol that intended to induce muscle damage. It included three sets of 12 back squats at an intensity of 70% of their 1RM with 150 s of rest between sets. In addition, subjects performed a combination of drop jumps from a 45 cm high box, landed in a deep squat, for three sets of 15 repetitions with 60 s of rest between sets. To be clear, this combination is a classic drop jump off the box where athletes are supposed to minimize contact time with the floor while still jumping as high as possible. This jump off the floor is then to be landed in a deep squat.

2.7. Supplementation

Following the training protocol, subjects drank 60 mg CBD solubilised with 250 mL water or a PLA drink directly after exercise. Specific CBD and PLA caps that fitted on a standard plastic bottle were produced that could only be distinguished by their color (red/white cap) (ATHENION GmbH, Berlin, Germany). There was no difference in taste or appearance between the two beverages. Other foods, dietary supplements, or caffeinated drinks were prohibited for the following three hours.

2.8. Statistical Analysis

Raw data were analyzed using the statistical software R version 4.0.4 (R Core Team, 2021). The raw data of CK and Myo were transformed to their natural logarithms ($\ln$) before data analysis. The values of CMJ and of 1RM were used as-is. The individual time interval (Δt [h]) since the beginning of the investigation was introduced as an additional covariate. Data analysis was performed by means of linear mixed-effects models (LME). By definition, $\ln$ (CK), $\ln$ (Myo), CMJ, and 1RM represented the dependent variables. Model building was performed independently for each of these measures. We were precisely interested in the potential effects of CBD on the course of recovery. Therefore, all models invariably encompassed the interaction term of CBD with the recovery interval (RI, levels T0 through T) as a fixed effect. Likewise, RI itself axiomatically was included as a fixed effect. Effect sizes (ES) of the fixed effects were calculated as where t represents the t -test statistic and DF the corresponding degrees of freedom and classified as trivial ($d < 0.2$), small ($0.2 \leq d \leq 0.5$), medium ($0.5 \leq d \leq 0.8$), or large ($d > 0.8$). Initially, random effects were merely assumed between the individual intercepts of each measure. Subsequently, Δt was included as a random effect, where linear individual trends were assumed. The presence of potentially non-linear individual trends was then investigated by upgrading to 2nd or 3rd order natural splines of Δt . Moreover, Δt was invariably incorporated into the models as 1st order continuous autoregressive term, i.e., the data were assumed to be individually autocorrelated in time. After the development of an appropriate random effects structure, it was tested whether Δt also contributes to general trends in the population, i.e., whether it represents a significant fixed effect. In either case, model comparisons were based on likelihood statistics and of changes of the Akaike information criterion (AIC).

3. Results

The mean and standard error (mean $\pm$ SE) of all parameters for each group can be seen in Table 2. The means $\pm$ SE for each parameter is depicted in Figure 2. Individual values can be found in Tables S5–S8 as well as Figures S1–S6 in the Supplemental Material.

Table 2. All average values of the muscle damage parameters CK and Myo as well as the performance in 1RM BS and CMJ.

Parameter	Units	PLA (<i>n</i> = 16)			CBD (<i>n</i> = 16)			ES
		PRE	POST	Δ	PRE	POST	Δ	
Creatine Kinase T 24	[U/L]	211.6 ± 16.9	399.9 ± 68.4	188.3 ± 58.4 ***	197.4 ± 19.1	351.5 ± 61.3	154.1 ± 61.1 ***	-
Creatine Kinase T 48	[U/L]	196.8 ± 39.1	530.8 ± 237.5	333.9 ± 238.6 ***	188.6 ± 25.2	320.9 ± 65.9	132.4 ± 66.0	-
Creatine Kinase T 72	[U/L]	347.9 ± 151.3	3417.7 ± 2119.9	3069.7 ± 2106.6 ***	181.9 ± 25.2	232.6 ± 35.0	50.7 ± 44.7 ***/#	-0.24
Myoglobin T 24	[ng/mL]	31.8 ± 8.3	38.4 ± 15.1	6.6 ± 9.2 **	31.9 ± 15.0	31.8 ± 8.3	-0.1 ± 12.7 **	-
Myoglobin T 48	[ng/mL]	30.5 ± 10.0	33.9 ± 15.4	3.5 ± 19.0	28.3 ± 6.4	37.5 ± 25.6	9.2 ± 25.0 **	-
Myoglobin T 72	[ng/mL]	30.3 ± 5.8	37.1 ± 15.8	6.8 ± 15.0 ***	28.1 ± 5.6	34.4 ± 11.3	6.3 ± 8.6 ***/#	0.21
Back Squat T 24	[kg]	118.0 ± 6.0	114.7 ± 5.5	-3.3 ± 1.5 **	118.1 ± 5.5	117.5 ± 5.5	-0.6 ± 0.8 **	-
Back Squat T 48	[kg]	118.3 ± 6.0	116.3 ± 6.7	-2.0 ± 1.9	119.1 ± 5.6	118.0 ± 5.4	-1.1 ± 1.0	-
Back Squat T 72	[kg]	118.2	116.2 ± 6.8	-2.0 ± 1.6	118.8 ± 5.8	120.0 ± 5.6	1.3 ± 0.9 #	0.37
Counter Movement Jump T 24	[cm]	55.7 ± 2.0	54.6 ± 1.5	-1.2 ± 0.7	54.4 ± 1.8	53.6 ± 1.7	-0.8 ± 0.4	-
Counter Movement Jump T 48	[cm]	54.9 ± 2.0	55.0 ± 2.1	0.1 ± 0.6	54.9 ± 1.8	54.0 ± 1.8	-0.8 ± 0.6	-
Counter Movement Jump T 72	[cm]	54.4 ± 2.0	54.1 ± 2.1	-0.4 ± 0.8	55.3 ± 1.7	55.5 ± 1.7	0.2 ± 0.6	-

The mean ± standard error ($\mu \pm \sigma x$) of all parameters for each group before (PRE) and after (POST) training, the corresponding individual differences (Δ) and the level of significance (significant difference over time: * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$; significant difference between groups: differences. # = $p < 0.05$, ##) as well as the effect size ($ES = 2 \cdot t/DF^2$) in case of significant group.




Figure 2. The mean ± SE for both groups at all time points. (a) Skeletal Muscle Specific Creatine Kinase. (b) Myoglobin. (c) Back Squat, (d) Countermovement Jump. *** indicates a significant difference over time with $p < 0.001$; ** = $p < 0.01$. Indicates a significant difference between groups at given time point with # = $p < 0.05$.

3.1. Physiological Adaptations over Time (Random Effects)

The models for $\ln$ (CK) and for $\ln$ (Myo) required 2nd natural splines of Δt in order to appropriately describe individual trends. Both, the introduction of linear and 2nd order terms resulted in significant improvements of the model fits ($p < 0.01$, respectively). The introduction of higher-order terms of Δt corrupted the models as inferred by AIC increases. The individual trends of CMJ and 1RM, by contrast, were sufficiently described by linear terms. In both cases, the introduction yielded significant model improvements ($p < 0.001$, respectively). Second-order natural splines, subsequently, yielded no further improvements but undesirable increases in AIC. $\ln$ (CK) and $\ln$ (Myo) showed significant decreases during the course of the study (CK: $p < 0.01$, ES: -0.286 ; Myo: $p < 0.05$, ES: -0.25). Meanwhile, 1RM showed a significant increase ($p < 0.05$, ES: 0.342) (See Tables S1–S4 for the corresponding parameter estimates (Δt)).

3.2. Effects on Skeletal Muscle Damage

Tables S1–S4 (Supplemental Material) show the fixed effects of the final models for $\ln$ (CK), $\ln$ (Myo), CMJ, and 1RM, respectively. In the case of statistically significant effects ($p < 0.05$), the corresponding effect sizes (ES) have been added. In addition to the individual trends, $\ln$ (CK), $\ln$ (Myo), and 1RM required the introduction of a linear term for Δt . $\ln$ (CK) was significantly elevated in both groups after 24 h ($p < 0.001$; ES: 0.69), 48 h ($p < 0.001$; ES: 0.531), and 72 h ($p < 0.001$; ES: 0.517). CBD attenuated the increase after 72 h, resulting in a significant group difference ($p < 0.05$; ES: -0.24). No significant effects of CBD on $\ln$ (CK) were apparent at T 24 and T 48. In a similar fashion, $\ln$ (Myo) showed a significant increase in both groups after 24 h ($p < 0.01$; ES: 0.301), 48 h ($p < 0.01$; ES: 0.281), and 72 h ($p < 0.001$; ES: 0.417). Again, after 72 h, CBD showed a dampening effect on $\ln$ (Myo), which led to a significant group difference ($p < 0.05$; ES: -0.214).

3.3. Effects on Performance

CMJ did not show any significance at any time point in both groups. It also did not change significantly over the course of the study (Δt). 1RM was significantly reduced in both groups after 24 h ($p < 0.01$ for both groups) but not after 48 or 72 h. Although changes from T 0 to T 72 were no significant, CBD resulted in a significant group difference ($p < 0.05$; ES: 0.371). Changes in performance differ quite substantially between subjects. Therefore, it is worth looking at the individual changes to see how many subjects increased, decreased, or maintained performance. After 24 h, five subjects of the CBD group and nine subjects of the PLA group showed decreases in their performance, while three subjects of each group improved. Eight subjects of the CBD group and four subjects of the PLA group were able to maintain their performance. After 48 h, five subjects of the CBD group and six subjects of the PLA group showed decreases, while three of the CBD group and six of the PLA group showed increasing performances. Eight and four subjects of the CBD and PLA group maintained performance, respectively. After 72 h, the CBD group had two subjects with decreases in performance while the PLA group included eight. Seven subjects of the CBD group even increased their performance, while only three did so in the PLA group. Seven subjects of the CBD and four of the PLA group maintained performance, respectively.

4. Discussion

This study examined the effect of a single CBD supplementation following an intensive strength protocol on skeletal muscle damage and lower body performance. The results clearly show an increase in CK and Myo concentration in both groups at all time points (T 24, T 48, T 72) with significant differences between the groups at T 72 (ES: CK = -0.236 ; Myo = -0.214). The exercise induced muscle damage and the increase in CK and Myo concentration after T 24 confirms observations from previous studies [1–3,52]. However, a single supplementation of CBD did not reduce muscle damage within 48 h. A longer time interval of at least 72 h, however, might be sufficient for effects of CBD to occur to a detectable extends. Therefore, this study cannot confirm the inhibitory effect of CBD

on CK concentration after 24 h seen in a pilot project with very well-trained strength athletes [52]. Furthermore, it can be assumed that different nutritional supplements can promote recovery at different rates despite similar mechanisms. Research has shown that a protein–carbohydrate combination can inhibit the increase in CK concentration after the same training protocol in athletes with similar performance levels [3]. A reason regarding the effects of CBD on muscle damage is possibly the absorption rate. Previous studies have shown that the maximum absorption of CBD can take between one–four hours [55]. Therefore, it can be speculated that post-exercise use may be too late to reduce muscle damage. Consequently, it may be beneficial to add CBD before training to bring the rate of absorption in line with the end of training. Regarding performance outcomes, a significant decrease was observed in 1RM but not in CMJ. The results of the 1RM are confirmed with observations of previous studies [1,3,52,56]. However, the decrease in performance in the CBD group is smaller than in the pilot study [52]. This could be caused by the different performance levels of the studies. The absolute strength values and training loads were significantly higher in the pilot study than in the present study (146.5 ± 21.6 vs. 118.6 ± 22.6 [kg]). This could explain the different changes in CK concentrations. An important observation of this study is the group difference between the CBD and PLA groups at T 72 ($p < 0.05$; ES: 0.371). In contrast to the PLA group, the CBD group was able to recover the squat performance and even slightly increase it on average (Table 2). This observation is in line with the kinetics of CK and Myo concentration. It seems that after 72 h, the CBD group has completely recovered and restored its maximum performance. However, these effects cannot be reproduced in the CMJ. In CMJ, neither significant changes at any time point nor differences between groups were observed. As a result, it can be assumed that the training protocol did not induce sufficient power reduction or that the Jump and Reach Test is not the optimal test for power performance. Consequently, jumping ability should ideally be measured with a force plate, and possibly the loading protocol should be revised to measure power change in the CMJ. In addition to the influence of CBD supplementation on performance, the influence of measurement repetition was examined due to the six-arm study design. All models required the introduction of Δt in a slightly differing manner. Fundamentally, $\ln$ (CK) and $\ln$ (Myo) showed significant negative population trends, while 1RM showed a linear increase. Although small in size (ES: $\ln$ (CK): -0.286 ; $\ln$ (Myo): -0.25 ; 1RM: 0.342 —see Tables S1–S3), this indicates the presence of significant adaptation processes throughout the study. Physiologically, the findings are mutually consistent. However, the imperative precondition of being experienced in strength training was not met sufficiently by all participants. Otherwise, the comparably small number of training bouts should not have induced detectable physiological changes. Formally, individual intercepts or trends could also have been considered to constitute random effects nested within the respective arm of the study. This, however, would have presumed random scatter of the individual parameters between arms. After all, the order of the arms had been individually randomized to this end. This presumption, however, is false. Rather, Δt must be considered to consistently take significant individual and possibly non-linear effects. For practical reasons, CBD and PLA were not administered in a strictly alternating order. Rather, the administration patterns were randomized independently within each of the subjects. Therefore, the factors Δt , i.e., adaptation, and CBD confound fundamentally. Hence, the potential effects of CBD might be slightly obliterated. In fact, the first out of the six testing sequences merely featured six CBD but ten PLA treatments. But the highest proportions of CBD treatments precisely occurred during the midsection of the study (sequences 3 and 4, 9/16 and 10/16 CBD treatments, respectively) while the remaining sequences happened to be largely balanced. The potential effects of CBD are, thus, unlikely to become significantly biased by adaptation. One of the most advantageous features of LMEs is exactly to account for time-dependent covariates, if appropriately modeled. As is fundamentally desirable in sports science and sports medicine, adaption processes at individual and population levels have been reflected as far as possible here. The validity of the model estimates, therefore, appears largely untainted with this potential

confounder. Although this study determined important findings to the effect of a single CBD supplementation after intensive strength training, there are also some limitations. The results of this study cannot be exactly compared to previous studies without mentioning the differing strength levels. Due to the training restriction before and during the intervention, subjects were not able to follow their normal strength training routine. Very well-trained strength athletes (squat performance above 150% of their BW) usually train more than four times a week to achieve the appropriate training volume for all muscle groups [57]. Therefore, it was impossible to recruit a population of highly well-trained athletes in this study design. In addition, the use of a jump-and-reach test might not be sufficient to monitor performance in this study design. Furthermore, no statements can currently be made about the effects of CBD after endurance training or after chronic supplementation and training. In order to minimize the limitations and to prevent possible adaptation processes due to the intervention, future studies should mainly include well- to very well-trained athletes [53,58] to assess the effect of CBD in the recovery process. Additionally, to performance and muscle damage parameters, inflammatory, immune, and antioxidative biomarkers should be also included in future studies for a more detailed analysis of the muscle healing process and adaptation.

5. Conclusions

This study investigated the effects of CBD after 24, 48, and 72 h following a single resistance training in a placebo-controlled, six-arm crossover study. No strong effects were observed in biomarkers or performance parameters. However, small and significant effects of a single supplementation of CBD on CK and Myo concentrations were observed after 72 h. For CBD to have a stronger effect on recovery processes after intense strength training, continuous and repetitive supplementation is probably needed. For clearer statements, however, further studies on pro-regenerative and recovery effects of CBD after strength training or other sports are essential.

Supplementary Materials: The following are available online at <https://www.mdpi.com/article/10.3390/nu13093028/s1>, Figure S1: Individual ln (CK) trends (coloured) and the mean $\pm$ 95% confidence interval (black) for each group and time point (indicated above each panel), Figure S2: Individual ln (CK) trends (colored) and the mean $\pm$ 95% confidence interval (black) for each group and time point (indicated above each panel) cut at ln (CK) = 7.5 to eliminate outliers for better visibility of changes over time, Figure S3: Individual ln (Myo) trends (colored) and the mean $\pm$ 95% confidence interval (black) for each group and time point (indicated above each panel), Figure S4: Individual ln (Myo) trends (coloured) and the mean $\pm$ 95% confidence interval (black) for each group and time point (indicated above each panel) cut at ln (Myo) = 5.0 to eliminate outliers for better visibility of changes over time, Figure S5: Individual 1RM Back Squat trends (coloured) and the mean $\pm$ 95% confidence interval (black) for each group and time point (indicated above each panel), Figure S6: Individual CMJ trends (coloured) and the mean $\pm$ 95% confidence interval (black) for each group and time point (indicated above each panel), Table S1: Effects of recovery interval (RI T 0—RI T 72) on ln (CK), Table S2: Effects of recovery interval (RI T 0—RI T 72) on ln (Myo), Table S3: Effects of recovery interval (RI T 0—RI T 72) on 1RM, Table S4: Effects of recovery interval (RI T 0—RI T 72) on CMJ, Table S5: Individual CK values for all time points and each group [in U/L], Table S6: Individual Myo values for all time points and each group [in ng/mL], Table S7: Individual 1RM Back Squat values for all time points and each group [in kg], Table S8: Individual CMJ values for all time points and each group [in cm].

Author Contributions: Conceptualization, P.D. and E.I.; methodology, P.D. and E.I.; formal analysis, U.F. and S.V.; investigation, E.I. and L.S.; resources, P.D.; data curation, U.F. and S.V.; writing—original draft preparation, E.I., S.V., L.S. and U.F.; writing—review and editing, E.I., S.V. and P.D.; visualization, S.V. and U.F.; supervision, P.D.; project administration, E.I.; funding acquisition, P.D. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by ATHENION GmbH, Berlin, Germany.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Ethics Committee of GERMAN SPORTS UNIVERSITY COLOGNE, GERMANY (155/2018; approved on 5 December 2018).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The data presented in this study are available in Tables S5–S8 in the Supplemental Material.

Acknowledgments: The authors would like to thank all subjects for their voluntary participation.

Conflicts of Interest: The funders had no role in the design of the study; in the collection, analyses, or interpretation of data in the writing of the manuscript.

Abbreviations

The following abbreviations are used in this manuscript:

1RM	One-Repetition Maximum
BS	Back Squat
CMJ	Countermovement Jump
CK	Skeletal Muscle Specific Creatine Kinase
Myo	Myoglobin
ln	Logarithm
SE	Standard Error
ES	Effect Size
CBD	Cannabidiol
PLA	Placebo
Δ^9 -THC	Δ^9 -tetrahydrocannabinol
h	hours
FDA	U.S. Food and Drug Administration
WADA	World Anti-Doping Agency
NCSA	National Strength and Conditioning Association
AIC	Akaike Information Criterion
Δ	Delta
t	time
NSAIDs	Non-Steroidal Anti-Inflammatory Drugs
NFL	National Football League
RI	Recovery Interval
LME	Linear Mixed Effect Model
DF	Degrees of Freedom
JAK	Janus Kinase
STAT	Signal Transducer and Activator of Transcription
EU	European Union

References

- Isenmann, E.; Trittel, L.; Diel, P. The effects of alpha lipoic acid on muscle strength recovery after a single and a short-term chronic supplementation—A study in healthy well-trained individuals after intensive resistance and endurance training. *J. Int. Soc. Sports Nutr.* **2020**, *17*, 1–13. [[CrossRef](#)]
- Isenmann, E.; Blume, F.; Bizjak, D.; Hundsdörfer, V.; Pagano, S.; Schibrowski, S.; Simon, W.; Schmandra, L.; Diel, P. Comparison of pro-regenerative effects of carbohydrates and protein administered by shake and non-macro-nutrient matched food items on the skeletal muscle after acute endurance exercise. *Nutrients* **2019**, *11*, 744. [[CrossRef](#)]
- Janning, M.; Isenmann, E.; Diel, P. Comparison of pro-regenerative nutritive actions given by foodstuff or shake after strength exercise. *Ger. J. Sports Med.* **2020**, *71*, 9. [[CrossRef](#)]
- Bloch, W. Muscle healing: Physiology and adverse factors. In *Muscle Injuries in Sports*; Müller-Wohlfahrt, H.W., Ueblicher, P., Hänsel, L., Garrett, W.E., Eds.; Thieme: Stuttgart, Germany, 2013.
- Koh, T.J. Physiology and mechanism of skeletal muscle damage. In *Skeletal Muscle Damage and Repair*; Tiidus, P.M., Ed.; Human Kinetics: Champaign, IL, USA, 2008.
- Sayers, S.P.; Hubal, M.J. Histological, chemical, and functional manifestation of muscle damage. In *Skeletal Muscle Damage and Repair*; Tiidus, P.M., Ed.; Human Kinetics: Champaign, IL, USA, 2008.
- Lieberman, H.R.; Marriott, B.P.; Williams, C.; Judelson, D.A.; Glickman, E.L.; Geiselman, P.J.; Dotson, L.; Mahoney, C.R. Patterns of dietary supplement use among college students. *Clin. Nutr.* **2015**, *34*, 976–985. [[CrossRef](#)] [[PubMed](#)]

8. National Study on Substance Use Habits of College Student-Athletes. 2018. Available online: https://www.wada-ama.org/sites/default/files/prohibited_list_2018_summary_of_modifications_en.pdf (accessed on 10 July 2021).
9. Tscholl, P.; Alonso, J.M.; Dollé, G.; Junge, A.; Dvorak, J. The use of drugs and nutritional supplements in top-level track and field athletes. *Am. J. Sports Med.* **2010**, *38*, 133–140. [[CrossRef](#)] [[PubMed](#)]
10. Zeiger, J.S.; Silvers, W.S.; Fleegler, E.M.; Zeiger, R.S. Cannabis use in active athletes: Behaviors related to subjective effects. *PLoS ONE* **2019**, *14*, e0218998. [[CrossRef](#)]
11. Ware, M.A.; Jensen, D.; Barrette, A.; Vernec, A.; Derman, W. Cannabis and the health and performance of the elite athlete. *Clin. J. Sport Med.* **2018**, *28*, 480–484. [[CrossRef](#)]
12. Docter, S.; Khan, M.; Gohal, C.; Ravi, B.; Bhandari, M.; Gandhi, R.; Leroux, T. Cannabis use and sport: A systematic review. *Sports Health A Multidiscip. Approach* **2020**, *12*, 189–199. [[CrossRef](#)]
13. Hartsel, J.A.; Eades, J.; Hickory, B.; Makriyannis, A. Cannabis sativa and Hemp. In *Nutraceuticals*; Gupta, R.C., Ed.; Academic Press: Boston, CA, USA, 2016; Chapter 53; pp. 735–754. [[CrossRef](#)]
14. Ligresti, A.; Petrocellis, L.D.; Marzo, V.D. From phytocannabinoids to cannabinoid receptors and endocannabinoids: Pleiotropic physiological and pathological roles through complex pharmacology. *Physiol. Rev.* **2016**, *96*, 1593–1659. [[CrossRef](#)]
15. Laprairie, R.B.; Bagher, A.M.; Kelly, M.E.M.; Denovan-Wright, E.M. Cannabidiol is a negative allosteric modulator of the cannabinoid CB1 receptor. *Br. J. Pharmacol.* **2015**, *172*, 4790–4805. [[CrossRef](#)] [[PubMed](#)]
16. Thomas, A.; Baillie, G.L.; Phillips, A.M.; Razdan, R.K.; Ross, R.A.; Pertwee, R.G. Cannabidiol displays unexpectedly high potency as an antagonist of CB1 and CB2 receptor agonists in vitro. *Br. J. Pharmacol.* **2007**, *150*, 613–623. [[CrossRef](#)]
17. Maurya, N.; Velmurugan, B.K. Therapeutic applications of cannabinoids. *Chem.-Biol. Interact.* **2018**, *293*, 77–88. [[CrossRef](#)]
18. Kozela, E.; Pietr, M.; Juknat, A.; Rimmerman, N.; Levy, R.; Vogel, Z. Cannabinoids Δ^9 -Tetrahydrocannabinol and Cannabidiol differentially inhibit the Lipopolysaccharide-activated NF- κ B and Interferon- β /STAT proinflammatory pathways in BV-2 microglial cells. *J. Biol. Chem.* **2010**, *285*, 1616–1626. [[CrossRef](#)]
19. Kozela, E.; Lev, N.; Kaushansky, N.; Eilam, R.; Rimmerman, N.; Levy, R.; Ben-Nun, A.; Juknat, A.; Vogel, Z. Cannabidiol inhibits pathogenic T cells, decreases spinal microglial activation and ameliorates multiple sclerosis-like disease in C57BL/6 mice. *Br. J. Pharmacol.* **2011**, *163*, 1507–1519. [[CrossRef](#)]
20. Malfait, A.M.; Gallily, R.; Sumariwalla, P.F.; Malik, A.S.; Andreaskos, E.; Mechoulam, R.; Feldmann, M. The nonpsychoactive cannabis constituent cannabidiol is an oral anti-arthritis therapeutic in murine collagen-induced arthritis. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 9561–9566. [[CrossRef](#)]
21. Borrelli, F.; Aviello, G.; Romano, B.; Orlando, P.; Capasso, R.; Maiello, F.; Guadagno, F.; Petrosino, S.; Capasso, F.; Marzo, V.D.; et al. Cannabidiol, a safe and non-psychotropic ingredient of the marijuana plant *Cannabis sativa*, is protective in a murine model of colitis. *J. Mol. Med.* **2009**, *87*, 1111–1121. [[CrossRef](#)] [[PubMed](#)]
22. Weiss, L.; Zeira, M.; Reich, S.; Har-Noy, M.; Mechoulam, R.; Slavin, S.; Gallily, R. Cannabidiol lowers incidence of diabetes in non-obese diabetic mice. *Autoimmunity* **2006**, *39*, 143–151. [[CrossRef](#)]
23. Wilkinson, J.D.; Williamson, E.M. Cannabinoids inhibit human keratinocyte proliferation through a non-CB1/CB2 mechanism and have a potential therapeutic value in the treatment of psoriasis. *J. Dermatol. Sci.* **2007**, *45*, 87–92. [[CrossRef](#)] [[PubMed](#)]
24. Ramirez, B.G. Prevention of Alzheimer's disease pathology by cannabinoids: Neuroprotection mediated by blockade of microglial activation. *J. Neurosci.* **2005**, *25*, 1904–1913. [[CrossRef](#)]
25. García-Arencibia, M.; González, S.; de Lago, E.; Ramos, J.A.; Mechoulam, R.; Fernández-Ruiz, J. Evaluation of the neuroprotective effect of cannabinoids in a rat model of Parkinson's disease: Importance of antioxidant and cannabinoid receptor-independent properties. *Brain Res.* **2007**, *1134*, 162–170. [[CrossRef](#)] [[PubMed](#)]
26. Consroe, P.; Laguna, J.; Allender, J.; Snider, S.; Stern, L.; Sandyk, R.; Kennedy, K.; Schram, K. Controlled clinical trial of cannabidiol in Huntington's disease. *Pharmacol. Biochem. Behav.* **1991**, *40*, 701–708. [[CrossRef](#)]
27. Campos, A.C.; Moreira, F.A.; Gomes, F.V.; Bel, E.A.D.; Guimarães, F.S. Multiple mechanisms involved in the large-spectrum therapeutic potential of cannabidiol in psychiatric disorders. *Philos. Trans. R. Soc. B Biol. Sci.* **2012**, *367*, 3364–3378. [[CrossRef](#)] [[PubMed](#)]
28. Jones, N.A.; Glyn, S.E.; Akiyama, S.; Hill, T.D.; Hill, A.J.; Weston, S.E.; Burnett, M.D.; Yamasaki, Y.; Stephens, G.J.; Whalley, B.J.; et al. Cannabidiol exerts anti-convulsant effects in animal models of temporal lobe and partial seizures. *Seizure* **2012**, *21*, 344–352. [[CrossRef](#)] [[PubMed](#)]
29. Jones, N.A.; Hill, A.J.; Smith, I.; Bevan, S.A.; Williams, C.M.; Whalley, B.J.; Stephens, G.J. Cannabidiol displays antiepileptiform and antiseizure properties in vitro and in vivo. *J. Pharmacol. Exp. Ther.* **2009**, *332*, 569–577. [[CrossRef](#)] [[PubMed](#)]
30. Campos, A.C.; Guimarães, F.S. Evidence for a potential role for TRPV1 receptors in the dorsolateral periaqueductal gray in the attenuation of the anxiolytic effects of cannabinoids. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* **2009**, *33*, 1517–1521. [[CrossRef](#)]
31. Petrocellis, L.D.; Ligresti, A.; Moriello, A.S.; Iappelli, M.; Verde, R.; Stott, C.G.; Cristino, L.; Orlando, P.; Marzo, V.D. Non-THC cannabinoids inhibit prostate carcinoma growth in vitro and in vivo: Pro-apoptotic effects and underlying mechanisms. *Br. J. Pharmacol.* **2012**, *168*, 79–102. [[CrossRef](#)]
32. Ligresti, A.; Moriello, A.S.; Starowicz, K.; Matias, I.; Pisanti, S.; Petrocellis, L.D.; Laezza, C.; Portella, G.; Bifulco, M.; Marzo, V.D. Antitumor activity of plant cannabinoids with emphasis on the effect of cannabidiol on human breast carcinoma. *J. Pharmacol. Exp. Ther.* **2006**, *318*, 1375–1387. [[CrossRef](#)]

33. Campos, A.C.; Fogaça, M.V.; Sonego, A.B.; Guimarães, F.S. Cannabidiol, neuroprotection and neuropsychiatric disorders. *Pharmacol. Res.* **2016**, *112*, 119–127. [CrossRef]
34. State of Colorado, U. Amendment 64 Implementation. 2012. Available online: <https://www.colorado.gov/pacific/sites/default/files/13%20Amendment%2064%20LEGIS.pdf> (accessed on 10 July 2021).
35. Judgement of 19 November 2020. Available online: <https://eur-lex.europa.eu/legal-content/DE/TXT/?uri=CELEX%3A62018CJ0663;CaseC-663/18.ECLI:EU:C:2020:938> (accessed on 29 August 2021).
36. World Anti Doping Agency. Summary of Major Modifications and Explanatory Notes. 2018 Prohibited List. Available online: https://www.ncaa.org/sites/default/files/2018RES_Substance_Use_Final_Report_FINAL_20180611.pdf (accessed on 10 July 2021).
37. Bergamaschi, M.M.; Queiroz, R.H.C.; Zuardi, A.W.; Crippa, J.A.S. Safety and side effects of cannabidiol, a cannabis sativa constituent. *Curr. Drug Saf.* **2011**, *6*, 237–249. [CrossRef]
38. Taylor, L.; Crockett, J.; Tayo, B.; Morrison, G. A phase 1, open-label, parallel-group, single-dose trial of the pharmacokinetics and safety of cannabidiol (CBD) in subjects with mild to severe hepatic impairment. *J. Clin. Pharmacol.* **2019**, *59*, 1110–1119. [CrossRef]
39. Taylor, L.; Gidal, B.; Blakey, G.; Tayo, B.; Morrison, G. A phase I, randomized, double-blind, placebo-controlled, single ascending dose, multiple dose, and food effect trial of the safety, tolerability and pharmacokinetics of highly purified cannabidiol in healthy subjects. *CNS Drugs* **2018**, *32*, 1053–1067. [CrossRef] [PubMed]
40. McCartney, D.; Benson, M.J.; Desbrow, B.; Irwin, C.; Suraev, A.; McGregor, I.S. Cannabidiol and sports performance: A narrative review of relevant evidence and recommendations for future research. *Sports Med.-Open* **2020**, *6*. [CrossRef] [PubMed]
41. Gamelin, F.X.; Cuvelier, G.; Mendes, A.; Aucouturier, J.; Berthoin, S.; Marzo, V.D.; Heyman, E. Cannabidiol in sport: Ergogenic or else? *Pharmacol. Res.* **2020**, *156*, 104764. [CrossRef]
42. Crippa, J.A.; Guimarães, F.S.; Campos, A.C.; Zuardi, A.W. Translational investigation of the therapeutic potential of cannabidiol (CBD): Toward a new age. *Front. Immunol.* **2018**, *9*, 86. [CrossRef]
43. Corroon, J.; Phillips, J.A. A cross-sectional study of cannabidiol users. *Cannabis Cannabinoid Res.* **2018**, *3*, 152–161. [CrossRef] [PubMed]
44. Moltke, J.; Hindocha, C. Reasons for cannabidiol use: A cross-sectional study of CBD users, focusing on self-perceived stress, anxiety, and sleep problems. *J. Cannabis Res.* **2021**, *3*, 1–12. [CrossRef] [PubMed]
45. Chagas, M.H.N.; Crippa, J.A.S.; Zuardi, A.W.; Hallak, J.E.C.; de Sousa, J.P.M.; Hirotsu, C.; Maia, L.; Tufik, S.; Andersen, M.L. Effects of acute systemic administration of cannabidiol on sleep-wake cycle in rats. *J. Psychopharmacol.* **2013**, *27*, 312–316. [CrossRef]
46. Murillo-Rodríguez, E.; Millán-Aldaco, D.; Cicconcelli, D.; Giorgetti, V.; Arankowsky-Sandoval, G.; Alcaraz-Silva, J.; Imperatori, C.; Machado, S.; Budde, H.; Tortorolo, P. Sleep-wake cycle disturbances and NeuN-altered expression in adult rats after cannabidiol treatments during adolescence. *Psychopharmacology* **2021**, *238*, 1437–1447. [CrossRef]
47. Babson, K.A.; Sottile, J.; Morabito, D. Cannabis, cannabinoids, and sleep: A review of the literature. *Curr. Psychiatry Rep.* **2017**, *19*, 1–12. [CrossRef]
48. Carlini, E.A.; Cunha, J.M. Hypnotic and antiepileptic effects of cannabidiol. *J. Clin. Pharmacol.* **1981**, *21*, 417S–427S. [CrossRef]
49. Malhotra, R.K. Sleep, recovery, and performance in sports. *Neurol. Clin.* **2017**, *35*, 547–557. [CrossRef]
50. Hatchett, A.; Armstrong, K.; Hughes, B.; Parr, B. The influence cannabidiol on delayed onset of muscle soreness. *Int. J. Phys. Educ. Sports Health* **2020**, *7*, 89–94.
51. Cochrane-Snyman, K.C.; Cruz, C.; Morales, J.; Coles, M. The effects of cannabidiol oil on noninvasive measures of muscle damage in men. *Med. Sci. Sports Exerc.* **2021**. [CrossRef] [PubMed]
52. Isenmann, E.; Veit, S.; Diel, P. Effects of cannabidiol supplementation on the skeletal muscle regeneration after intensive resistance training. *Med. Sci. Sports Exerc.* **2020**, *52*, 766. [CrossRef]
53. Junior, E.R.T.S.; de Salles, B.F.; Dias, I.; Ribeiro, A.S.; Simão, R.; Willardson, J.M. Classification and determination model of resistance training status. *Strength Cond. J.* **2021**. Online ahead of print. [CrossRef]
54. Miller, T.; Haywood, K.M.; Robertson, M.A.; Getchell, N. *NSCA's Guide to Tests and Assessments*; Human Kinetics: Champaign, IL, USA, 2012.
55. Millar, S.A.; Stone, N.L.; Yates, A.S.; O'Sullivan, S.E. A systematic review on the pharmacokinetics of cannabidiol in humans. *Front. Pharmacol.* **2018**, *9*, 1365. [CrossRef]
56. Isenmann, E.; Diel, P.; Geisler, S.; Schiffer, T. The effects of protein and carbohydrate supplementation on muscular regeneration after intense resistance training in soccer players. *Curr. Dev. Nutr.* **2020**, *4*, 1756. [CrossRef]
57. Schoenfeld, B.J.; Ogborn, D.; Krieger, J.W. Dose-response relationship between weekly resistance training volume and increases in muscle mass: A systematic review and meta-analysis. *J. Sports Sci.* **2016**, *35*, 1073–1082. [CrossRef]
58. Pauw, K.D.; Roelands, B.; Cheung, S.S.; de Geus, B.; Rietjens, G.; Meeusen, R. Guidelines to classify subject groups in sport-science research. *Int. J. Sports Physiol. Perform.* **2013**, *8*, 111–122. [CrossRef] [PubMed]