

Table S1. Significant relationships for women with obesity compared to normal-weight women.

Eating behaviors ^a	Less vs. Maintain			Less vs. More			More vs. Maintain		
	OR	CI	P-value	OR	CI	P-value	OR	CI	P-value
Healthfulness of food	0.81	0.39-1.69	0.58	1.73	0.87-3.44	0.12	2.07	1.31-3.29	p=0.002
Amount of food	0.29	0.12-0.63	p=0.002^a	0.23	0.11-0.47	p<0.0001^b	0.75	0.46-1.22	0.25
Size of meals	0.51	0.30-0.86	P=0.012	0.56	0.33-0.96	p=0.036	1.10	0.67-1.81	0.71
Snacking frequency	0.34	0.14-0.84	p=0.019	0.30	0.12-0.67	p=0.003	0.84	0.50-1.39	0.49
Amount of fat	0.45	0.24-0.85	p=0.014	0.35	0.16-0.75	p=0.007^c	0.75	0.45-1.27	0.28
Amount of fried or fast foods	0.55	0.35-0.88	p=0.012	0.79	0.42-1.47	0.45	1.30	0.71-2.38	0.39
Amount of fruit	0.34	0.12-0.93	p=0.036	0.31	0.12-0.79	p=0.015	0.81	0.52-1.27	0.36

^aPartly determined by Caucasian race (3.40 [CI 1.45-8.02], p=0.005).

^bPartly determined by higher education levels (0.46 [CI 0.22-0.95], p=0.035).

^cPartly determined by Caucasian race (2.65 [CI 1.39-5.0], p=0.003).