nutrients

MDPI

Supplementary Materials

Screening and Isolation of Potential Anti-Inflammatory Compounds from *Saxifraga atrata* via Affinity Ultrafiltration-HPLC and Multi-Target Molecular Docking Analyses

Gang Li [1], Yan Fang [1], Yonggui Ma [2], Yangzom Dawa [2], Qilan Wang [3], Jing Gan [1] and Jun Dang [3,*]

Table of Contents

Figure S1. Actual micro gel chp20p medium pressure liquid chromatography system 2

Figure S2. Schematic diagram of the principle of affinity ultrafiltration-HPLC 3

Figure S3. ESI mass spectrum of 11-O-(4'-O-methylgalloyl)-bergenin 4

Figure S4. 1H NMR Spectrum (600 MHz) of 11-O-(4'-O-methylgalloyl)-bergenin (in MeOH-d4) 5

Figure S5. 13C NMR Spectrum (151 MHz) of 11-O-(4'-O-methylgalloyl)-bergenin (in MeOH-d4) 6



Figure S1. Actual micro gel chp20p medium pressure liquid chromatography system.




Figure S2. Schematic diagram of the principle of affinity ultrafiltration-HPLC.




Figure S3. ESI mass spectrum of 11-O-(4'-O-methylgalloyl)-bergenin.




Figure S4. ^{1}H NMR Spectrum (600 MHz) of 11-O-(4'-O-methylgalloyl)-bergenin (in MeOH-d_4).




Figure S5. ^{13}C NMR Spectrum (151 MHz) of 11-O-(4'-O-methylgalloyl)-bergenin (in MeOH-d_4).