

Urinary sodium excretion and obesity markers among Bangladeshi adult population: pooled data from three cohort studies

Musarrat J. Rahman¹, Sarker M Parvez², Mahbubur Rahman², Feng J. He³, Solveig Cunningham⁴, K.M. Venkat Narayan⁴, Jaynal Abedin⁵, Abu Mohd Naser⁶

¹International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

²Environmental Interventions Unit, Infectious Disease Division, icddr,b, Dhaka, Bangladesh

³Centre for Environmental and Preventive Medicine, Wolfson Institute of Preventive Medicine, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, United Kingdom

⁴Emory Global Diabetes Research Center, Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, GA

⁵Data Science Institute, National University of Ireland Galway, Galway, Ireland

⁶Division of Epidemiology, Biostatistics and Environmental Health, School of Public Health, University of Memphis, Memphis, TN

Correspondence: Dr. Abu Mohd Naser, MBBS, PhD

Email: atitu@memphis.edu

Division of Epidemiology, Biostatistics, and Environmental Health
School of Public Health
University of Memphis

Figure S1: Cubic spline plots of 24-hour urinary sodium excretion and waist circumference, BMI, waist to hip ratio and waist to height ratio




Figure S2: Cubic spline plots of 24-hour urinary sodium excretion and body fat percentage and visceral fat percentage

