

Article

Legume Consumption and Blood Pressure Control in Individuals with Type 2 Diabetes and Hypertension: Cross-Sectional Findings from the TOSCA.IT Study

Marilena Vitale ¹, Annalisa Giosuè ¹, Sabina Sieri ², Vittorio Krogh ², Elena Massimino ¹, Angela Albarosa Rivellesse ¹, Gabriele Riccardi ¹, Olga Vaccaro ^{1,*} and Maria Masulli ¹

¹ Department of Clinical Medicine and Surgery, Federico II University of Naples, 80131 Naples, Italy; marilena.vitale@unina.it (M.V.); annalisa.giosue@unina.it (A.G.); elena.massimino@unina.it (E.M.); rivelles@unina.it (A.A.R.); riccardi@unina.it (G.R.); maria.masulli@unina.it (M.M.)

² Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei Tumori di Milano, 20121 Milan, Italy; sabina.sieri@istitutotumori.mi.it (S.S.); vittorio.krogh@istitutotumori.mi.it (V.K.)

* Correspondence: ovaccaro@unina.it; Tel.: +39-0817463665

Abstract: Background: Our aims were to evaluate the relationship of habitual legume consumption with blood pressure (BP) control in a large cohort of people with T2D and hypertension, and to investigate whether specific nutritional components of legumes or other foods may contribute to regulate BP levels. Methods: We studied 1897 participants with T2D and hypertension. Dietary habits were assessed through a validated food frequency questionnaire. Sex-specific quartiles of legume consumption were created. Results: Higher legume consumption was associated with a lower intake of energy, carbohydrates, glycaemic load, alcohol, and sodium, and a significantly greater intake of proteins, fat, monounsaturated, polyunsaturated, fibre, potassium, and polyphenols. Significantly lower systolic and diastolic BP values were observed in the highest vs. lowest quartile of legume consumption (132.9 ± 6.7 vs. 137.3 ± 7.0 mmHg, $p < 0.001$; 78.9 ± 4.1 vs. 81.0 ± 4.2 mmHg, $p = 0.002$; respectively), as well as the proportion of people meeting the treatment targets (61.3% vs. 37.4% and 71.3% vs. 52.4%, respectively, $p < 0.01$). This association was independent from other foods whose consumption is associated with the high legume intake. Conclusions: In people with T2D and hypertension, three servings of legumes per week are associated with significantly better BP control. This gives further support to current dietary guidelines in recommending the frequent consumption of legumes, as a “ready-to-use” dietary strategy to achieve optimal BP control.

Keywords: legume consumption; pulses; type 2 diabetes; hypertension; TOSCA.IT study

Citation: Vitale, M.; Giosuè, A.; Sieri, S.; Krogh, V.; Massimino, E.; Rivellesse, A.A.; Riccardi, G.; Vaccaro, O.; Masulli, M. Legume Consumption and Blood Pressure Control in Individuals with Type 2 Diabetes and Hypertension: Cross-Sectional Findings from the TOSCA.IT Study. *Nutrients* **2023**, *15*, 2895. <https://doi.org/10.3390/nu15132895>

Academic Editor: Carol Johnston

Received: 1 June 2023

Revised: 19 June 2023

Accepted: 21 June 2023

Published: 26 June 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Hypertension is a major risk factor for coronary artery disease, stroke, heart failure, peripheral vascular disease, and chronic kidney disease [1]. To date, hypertension affects between 16 and 37% of the population globally [2] and it is approximately twice as common in persons with type 2 diabetes [3,4], reaching a prevalence of about 80–90%.

Data from observational studies show that high blood pressure levels cause a significant increase in the risk of vascular complications in individuals with diabetes and predispose to a higher incidence of chronic kidney disease, ischemic cerebrovascular disease, retinopathy, and cardiovascular mortality [5,6,7,8]. There is solid evidence that blood pressure reduction in people with diabetes largely improves micro- and macrovascular outcomes. Data from the UK Prospective Diabetes Study showed that each 10 mmHg decrease in systolic blood pressure was associated with 12% reduction in risk to develop complications related to diabetes, 11% reduction in the risk of myocardial infarction, and

13% reduction in microvascular complications [9]. Therefore, the effective management of high blood pressure levels is a major target in individuals with diabetes.

Pharmacological treatment is required in the management of most patients, but lifestyle modification, including a better quality of diet, and weight management, are key components to control both blood glucose and blood pressure. Previous studies have shown that the consumption of specific foods, such as fruit and vegetables, dairy products, and whole grains, or the adherence to specific dietary patterns, such as the Mediterranean diet, or the low glycaemic index diet, can improve blood pressure control in diabetes patients [10,11,12]. The Dietary Approaches to Stop Hypertension (DASH) study, conducted more than 20 years ago [13,14], showed that a diet rich in plant-based foods, wholegrains, low-fat dairy products, and low sodium intake, is effective in the prevention and management of hypertension [15].

Among plant-based foods, legumes contain various components linked to lower blood pressure, including dietary fibre, bioactive peptides, and flavonoid polyphenols [16,17], but their relationship with blood pressure control in people with diabetes is little investigated and with discordant results [18]. A recent intervention trial of Jenkins et al., has shown a positive effect of legume consumption on systolic blood pressure in subjects with type 2 diabetes following a low-GI, legume rich diet, compared with a high wheat fibre diet [17]. However, the large daily amount of legumes used in this study (1 cup of legumes per day, i.e., 200 g/day) does not allow to extrapolate the results to real life conditions. Furthermore, another clinical trial aimed at determining the effects on cardio-metabolic risk factors of red meat replacement with legumes in people with type 2 diabetes, found no difference in systolic and diastolic blood pressures after 8 weeks of either legume-free diet or legume-based diet although improvements in other markers of metabolic health were observed [19].

Therefore, the aims of the present study were to evaluate the association between habitual legume consumption and systolic and diastolic blood pressure control in a large cohort of people with type 2 diabetes and hypertension, and to investigate whether specific nutritional components of legumes, or other foods associated with a legume rich dietary pattern, may contribute to modulate blood pressure levels in real life conditions.

2. Materials and Methods

2.1. Participants

We analysed the baseline data from the TOSCA.IT trial (URL: clinicaltrials.gov (accessed on 26 June 2023) NCT00700856), a randomized clinical trial designed to assess the cardiovascular effects of two different hypoglycaemic drugs. The TOSCA.IT study complied with the Declaration of Helsinki and was granted the approval of the Ethics Committee of the coordinating centre and of each participating centre; written informed consent was obtained from all participants. Details on the study protocol, inclusion and exclusion criteria have been published elsewhere [20,21]. Briefly, 3028 people with type 2 diabetes and with serum creatinine <1.5 mg/dL were recruited in 60 diabetes clinics distributed all over Italy. For the present analysis, we selected 1897 individuals with diagnosis of hypertension and with valid dietary data. Hypertension was defined as use of anti-hypertensive medications and/or systolic blood pressure (SBP) values ≥ 140 mmHg and/or diastolic blood pressure (DBP) values ≥ 90 mmHg [22].

For the purposes of the present analyses, we used data collected at baseline, prior to randomization to the study treatments.

2.2. Measurements

Height and weight were measured, and body mass index (BMI) was calculated as kg/m². Waist circumference was measured according to a standard protocol. Subjects were classified as current smokers when they reported having smoked at all over the previous 12 months or more.

Fasting blood sample were analysed in a central laboratory. High-performance liquid chromatography (HPLC) was used to measure HbA1c. Lipid profile (i.e., total and HDL-cholesterol, and triglycerides) were measured by standard methods. The LDL-cholesterol was calculated by the Friedewald equation for people with triglycerides values <400 mg/dL [23]. The study included patients inadequately controlled with metformin only (stable treatment for the last two months with metformin in monotherapy at 2 g/day) and HbA1c $\geq 7.0\%$ and $\leq 9.0\%$ [20]. Blood pressure was measured in a seated position after five minutes rest with a mercury sphygmomanometer. Three measurements were taken at 1 min intervals. The average value was considered in the analysis.

2.3. Dietary Assessment

Dietary habits during the past 12 months were assessed by the validated EPIC food frequency questionnaire (FFQ) [24] for a total of 248 questions concerning 188 different food items that were classified into 74 predefined food groups. The respondent indicates the absolute frequency of consumption of each item (per day, week, month, or year) and the quantity by selection of pictures showing a small, medium, and large portion size, with additional quantifiers (e.g., “smaller than the small portion” or “between the small and medium portion”) or by selection of a predefined standard portion when no image was available. Incomplete questionnaires and questionnaires with implausible data (i.e., energy intake less than 800 or greater than 5000 Kcal/day) were excluded from the analyses.

Through the use of a specifically designed software developed ad hoc by the Epidemiology and Prevention Unit, Fondazione IRCCS, Istituto Nazionale dei Tumori, Milan [25], questionnaire dietary data were converted into frequencies of consumption and average daily quantities and then linked to Italian food tables to obtain estimates of daily intake of macro- and micro-nutrients and energy. Polyphenol intake was assessed using a combination of USDA and Phenol-Explorer databases [26,27,28,29]. Use of drugs was assessed by interview.

2.4. Statistical Analysis

Data are given as numbers and percentages, or mean values (M) and standard deviation (SD) for continuous variables, as appropriate.

To correct for the different energy intake between subjects, the intake of legumes was expressed as g/1000 Kcal/day. The sex specific quartiles of legume consumption were created to assess the relation of legume consumption with blood pressure, anthropometric variables, glucose control, and plasma lipids.

To compare variables across the quartiles, the analysis of variance with a test for a linear trend was used for the continuous variables, and the χ^2 test was used to compare proportions.


Multivariate linear regression analysis was used to estimate the separate and combined association of legumes and other dietary factors with the variables of interest (i.e., systolic, and diastolic blood pressure) and results were expressed as regression coefficients (β). The explored dietary factors were cereals, wholegrains, potatoes, vegetables, fruits, nuts, fish, meat, eggs, dairy products, milk and yogurt, olive oil, animal fat, alcoholic beverages, sweeteners beverages, coffee and tea, and cakes and pastries. Multivariate linear regression analyses were performed to explore the impact of nutrients or bioactive compounds of legumes (i.e., vegetable proteins from food sources, fibre, sodium, potassium, and polyphenols) on systolic and diastolic blood pressure.

The statistical analyses were performed with IBM version 28.0. Statistical tests were two-sided, and p values of less than 0.05 were considered to indicate statistical significance.

3. Results

The study population consisted of 1079 men (56.9%) and 818 women (43.1%) with a history of type 2 diabetes 8.7 years (± 5.7 years) and hypertension of about 7.2 years (± 6.4 years). The mean age was 62.8 years (± 6.4 years), and BMI 30.8 ± 4.5 Kg/m². Most participants were taking anti-hypertensive medications (95.6%) and 22% of total population had micro-albuminuria (Supplemental Table S1).

The mean values of systolic and diastolic blood pressure, together with the percentage of people on target for systolic (<130 mmHg) and diastolic (<80 mmHg) blood pressure across the sex-specific quartiles of legume consumption are reported in Figure 1. Significantly lower levels of systolic and diastolic blood pressure were observed in the highest quartile of legume consumption (Q4) compared to the lowest (Q1) (132.9 ± 6.7 vs. 137.3 ± 7.0 mmHg, $p < 0.001$; 78.9 ± 4.1 vs. 81.0 ± 4.2 mmHg, $p = 0.002$; respectively). Accordingly, the percentage of people on target for systolic and diastolic blood pressure significantly increased across the quartiles of legumes intake (from 37.4% to 61.3% for SBP and from 52.4% to 71.3% for BDP, respectively ($p < 0.01$)) (Figure 1).






Figure 1. Systolic (A) and diastolic (B) blood pressure, expressed as absolute values and proportion of people on target (systolic blood pressure < 130 mmHg (C), diastolic blood pressure < 80 mmHg (D)), according to the sex-specific quartiles of legume intake in people with type 2 diabetes and hypertension.

No significant differences were observed across quartiles regarding age, diabetes duration, BMI, waist circumference, HbA1c, total cholesterol, LDL- and HDL-cholesterol, triglycerides, C-reactive protein, and percentage of people with micro-albuminuria (Supplemental Table S2). The same findings were observed after the exclusion of patients with micro-albuminuria.

The specific food pattern associated with legume consumption is shown in Table 1. A higher legume consumption was significantly associated with a greater intake of wholegrain cereals, vegetables, fruits (all types), fish, eggs, and olive oil ($p < 0.05$), and a lower intake of refined cereals, animal fat (i.e., butter, cream, etc.), alcoholic beverages, and cake and pastries ($p < 0.05$). No significant differences were detected for potatoes, nuts, meat, dairy products, milk and yogurt, sweetened beverages, and coffee and tea consumption (Table 1).

Table 1. Food intake (g/1000 Kcal/day) in people with type 2 diabetes and hypertension according to the sex-specific quartiles of legume intake (g/1000 kcal/day).

	Quartile 1 (2.9 ± 1.7)	Quartile 2 (8.1 ± 1.8)	Quartile 3 (14.1 ± 2.7)	Quartile 4 (28.3 ± 10.6)	<i>p</i> -Value
Cereals (pasta, rice, bread)	98.5 ± 40.7	98.1 ± 35.9	93.9 ± 33.9	86.2 ± 33.6	<0.001
Wholegrain cereals (bread)	6.7 ± 15.7	6.5 ± 13.8	6.9 ± 15.6	9.7 ± 18.3	0.006
Potatoes	9.2 ± 12.3	10.6 ± 14.9	11.0 ± 9.3	11.0 ± 12.6	0.074
Vegetables (salad, cooked vegetable)	87.1 ± 55.6	86.7 ± 41.8	98.1 ± 47.7	109.9 ± 49.1	<0.001
Fruits (all type)	148.7 ± 89.4	150.8 ± 88.1	156.9 ± 90.5	185.8 ± 106.9	<0.001
Nuts	0.67 ± 2.12	0.50 ± 0.86	0.51 ± 0.93	0.57 ± 1.43	0.238
Fish	19.2 ± 17.7	21.1 ± 14.7	23.1 ± 16.5	27.9 ± 20.8	<0.001
Meat (red, white, and processed)	53.5 ± 28.8	51.8 ± 23.9	53.0 ± 25.4	53.1 ± 25.2	0.776
Eggs	9.4 ± 7.8	10.2 ± 6.5	10.7 ± 7.0	11.2 ± 7.9	0.002
Dairy Products	20.7 ± 14.8	20.6 ± 13.8	20.0 ± 12.6	19.5 ± 12.7	0.491
Milk and Yogurt	108.4 ± 105.4	104.7 ± 101.6	110.8 ± 100.0	117.0 ± 110.6	0.320
Olive Oil	12.2 ± 6.3	13.0 ± 5.2	14.2 ± 5.5	15.8 ± 6.3	<0.001

Animal fat (butter, cream, etc.)	1.72 ± 1.78	1.60 ± 1.58	1.50 ± 1.46	1.24 ± 1.30	<0.001
Alcoholic beverages (wine, beer)	72.0 ± 98.0	64.1 ± 89.8	61.4 ± 81.2	51.3 ± 75.4	0.003
Sweetener beverages	29.9 ± 64.0	26.5 ± 56.7	24.6 ± 46.9	25.4 ± 48.4	0.466
Coffee and tea	80.8 ± 75.9	73.3 ± 59.1	76.9 ± 55.7	83.2 ± 72.5	0.105
Cake and pastries	20.9 ± 21.6	21.9 ± 18.8	18.3 ± 17.7	15.4 ± 14.3	<0.001

M ± SD.

As compared to subjects with lower legume intakes (Q1), participants reporting higher legume consumption (Q4) had significantly lower intake of energy, total carbohydrates and starch, glycaemic load, alcohol, and sodium ($p < 0.001$) (Table 2). Significantly greater intake of total proteins, especially from vegetable food sources, total fat, especially monounsaturated and polyunsaturated fat, fibre, potassium, and polyphenols, were also associated with higher legume consumption ($p < 0.05$). The intake of saturated fat, cholesterol and added sugar, and the glycaemic index of the diet did not change across the quartiles (Table 2).

Table 2. Energy and nutrient composition of the habitual diet in people with type 2 diabetes and hypertension according to the sex-specific quartiles of legume intake (g/1000 kcal/day).

	Quartile 1 (2.9 ± 1.7)	Quartile 2 (8.1 ± 1.8)	Quartile 3 (14.1 ± 2.7)	Quartile 4 (28.3 ± 10.6)	<i>p</i> -Value
Energy (Kcal/day)	1908 ± 821	1996 ± 730	1929 ± 795	1655 ± 607	<0.001
Proteins (% of TE)	18.1 ± 2.7	18.1 ± 2.4	18.3 ± 2.5	18.6 ± 2.7	0.006
from animal food sources (% of TE)	12.6 ± 3.3	12.5 ± 3.1	12.7 ± 3.1	12.9 ± 3.3	0.452
from vegetable food sources (% of TE)	5.5 ± 1.2	5.6 ± 1.2	5.6 ± 1.1	5.8 ± 1.0	0.006
Total fat (% of TE)	36.0 ± 6.5	36.5 ± 5.6	37.1 ± 5.9	37.7 ± 6.2	0.001
SAFA (% of TE)	12.3 ± 2.7	12.4 ± 2.5	12.3 ± 2.5	12.0 ± 2.5	0.187
MUFA (% of TE)	17.2 ± 4.0	17.6 ± 3.4	18.2 ± 3.6	18.8 ± 4.0	<0.001
PUFA (% of TE)	4.3 ± 1.1	4.3 ± 1.0	4.5 ± 1.2	4.7 ± 1.2	<0.001
Cholesterol (mg/1000 Kcal)	177.3 ± 56.2	183.2 ± 50.0	182.5 ± 49.2	184.5 ± 54.3	0.162
Carbohydrates (% of TE)	45.7 ± 8.0	45.3 ± 7.0	44.5 ± 7.2	43.6 ± 7.4	<0.001
Starch (% of TE)	29.3 ± 9.0	29.0 ± 8.1	28.2 ± 7.8	26.4 ± 7.7	<0.001
Soluble carbohydrates (% of TE)	16.4 ± 6.0	16.3 ± 5.6	16.2 ± 5.4	17.2 ± 5.5	0.030
Added Sugar (% of TE)	2.6 ± 3.8	2.4 ± 3.5	2.3 ± 2.7	2.3 ± 3.2	0.263
Fibre (g/1000 Kcal)	9.5 ± 2.4	9.9 ± 2.3	10.8 ± 2.4	12.7 ± 2.8	<0.001
Glycaemic Index (n)	51.7 ± 3.6	51.8 ± 3.1	51.6 ± 3.4	51.5 ± 3.6	0.729
Glycaemic Load (%)	117.5 ± 64.0	120.0 ± 51.4	113.8 ± 57.2	98.7 ± 41.6	<0.001
Alcohol (g/day)	12.3 ± 17.3	11.6 ± 18.1	10.4 ± 14.7	7.8 ± 12.6	<0.001
Sodium (mg/day)	2158 ± 1186	2250 ± 1108	2136 ± 1076	1767 ± 851	<0.001
Potassium (mg/day)	2883 ± 1093	3068 ± 1029	3089 ± 1124	2913 ± 1010	0.003
Polyphenols (mg/1000 Kcal/day)	345 ± 162	348 ± 139	371 ± 144	417 ± 168	<0.001

M ± SD. TE: Total Energy; SAFA: Saturated Fatty Acids; MUFA: Monounsaturated Fatty Acids; PUFA: Polyunsaturated Fatty Acids.

In multivariate analysis, systolic and diastolic blood pressure values were significantly and inversely associated with legume consumption ($\beta = -0.087$, $p < 0.001$; $\beta = -0.061$, $p = 0.014$, respectively) independently of any other dietary item; none of the other foods consumed by the study participants was associated with blood pressure (Table 3).

Table 3. Multivariate linear regression model assessing the relationship of individual foods with systolic and diastolic blood pressure.

	Systolic Blood Pressure		Diastolic Blood Pressure	
	β -Coefficient	<i>p</i> -Value	β -Coefficient	<i>p</i> -Value
Cereals (pasta, rice, bread)	−0.046	0.302	0.017	0.703
Wholegrain cereals (bread)	−0.004	0.863	−0.019	0.422
Potatoes	−0.004	0.853	0.016	0.508
Legumes	−0.087	<0.001	−0.061	0.014
Vegetables (salad, cooked vegetables)	−0.005	0.889	0.050	0.196
Fruits (all type)	−0.023	0.422	−0.003	0.905
Nuts	0.018	0.455	−0.013	0.572
Fish	0.036	0.122	0.025	0.551
Meat (red, white, and processed)	0.060	0.239	0.050	0.330
Eggs	0.010	0.688	0.005	0.844
Dairy products	0.043	0.127	0.016	0.566
Milk and yogurt	0.008	0.247	−0.039	0.242
Olive oil	0.008	0.848	−0.044	0.309
Animal fat (butter, cream, etc.)	0.044	0.079	0.052	0.057
Alcoholic beverages (wine, beer)	0.022	0.477	0.055	0.080
Sweetener beverages	0.024	0.342	0.049	0.062
Coffee and tea	0.027	0.265	−0.001	0.964
Cake and pastries	−0.012	0.733	0.035	0.305

In addition, in order to evaluate which nutrients linked legume consumption to systolic and diastolic blood pressure values, separate linear regression models were used. As reported in Table 4, the only dietary component associated with systolic and diastolic blood pressure values was dietary fibre ($\beta = -0.139$, $p = 0.007$; $\beta = -0.111$, $p = 0.031$, respectively). This association disappeared when legumes were included in the model and was replaced by a significant inverse association between systolic and diastolic blood pressure values with legume consumption, thus suggesting that dietary fibre plays a significant role in modulating the relationship between legume consumption and blood pressure control.

Table 4. Multivariate linear regression model assessing the relationship of nutrients and bioactive compounds with systolic and diastolic blood pressure.

	Systolic Blood Pressure		Diastolic Blood Pressure	
	β -Coefficient	<i>p</i> -Value	β -Coefficient	<i>p</i> -Value
Model 1				
Proteins from vegetable food sources	−0.004	0.946	0.065	0.233
Fibre	−0.139	0.007	−0.111	0.031
Sodium	0.021	0.628	0.017	0.700
Potassium	0.087	0.073	0.060	0.143
Polyphenols	−0.035	0.175	−0.031	0.223
Model 2				
Legumes	−0.089	<0.001	−0.082	0.002
Proteins from vegetable food sources	−0.052	0.357	0.021	0.706
Fibre	−0.059	0.287	−0.038	0.493
Sodium	0.011	0.797	0.008	0.859
Potassium	0.061	0.091	0.037	0.378
Polyphenols	−0.031	0.228	−0.028	0.280

4. Discussion

This study investigated, for the first time, the association between legume consumption and blood pressure in a large cohort of people with type 2 diabetes, in the context of real-life conditions, where the amount of legumes intake was that habitually consumed by the study population and substantially lower than in the intervention trials conducted so far [17,19,30]. Notably, in this cohort the highest quartile of legume consumption corresponded to an intake of 28.2 g/1000 Kcal/day that means 70 g (about 1/2 cup) 3 times a week, an intake fully consistent with the current nutritional recommendation for people with diabetes [31,32]. The results clearly showed that higher intakes of legumes are associated with lower levels of systolic and diastolic blood pressure, and with a greater proportion of people meeting the treatment target for blood pressure. Furthermore, multivariate analyses show that this association is independent from other foods whose consumption is associated with a legume rich dietary pattern, such as cereals, wholegrain foods, vegetables, and fruits, and may be, at least in part, due to the high fibre content of legumes.

Legume consumption in relation to blood pressure has been studied in people with or without type 2 diabetes in intervention studies exploring the beneficial effects of high legume intakes. Jenkins et al. [17] reported a beneficial effect of legume consumption on systolic blood pressure among individuals with type 2 who consumed about 1 cup of legumes per day (190 g/day) for 3 months within the context of a low-GI legume rich diet compared to those following a high wheat fibre diet. Another clinical trial aimed at evaluating the effects on cardio-metabolic risk factors of red meat replacement with legumes in people with type 2 diabetes, found no difference in systolic and diastolic blood pressures after 8 weeks of either a legume-free diet or a legume-based diet [19]. In this substitution study, half a cup of cooked legumes (70 g/day) was substituted for one serving of red meat. Moreover, in a meta-analysis of eight isocaloric dietary pulse intervention trials including 554 participants with and without type 2 diabetes or hypertension a significant improvement of blood pressure was achieved by a median consumption of 162 g/day of pulses [30].

A feature shared by all these studies is the high amounts of legumes used within dietary interventions which is unfeasible for the long term in real life conditions. Yet, our study confirms the association of legumes with lower systolic and diastolic blood pressure for a lesser amount of consumption (70 g three times a week), which is more feasible and in line with that recommended for the general population and suggested by the dietary guidelines for people with type 2 diabetes.

Prior population studies in non-diabetic people have also reported that an increased legume intake is associated with a reduced risk of developing hypertension. A meta-analysis of five observational studies reported a ~5% risk reduction to develop hypertension for about 70 g/day of legume consumption [33]. A subsequent work on 83,284 incident cases of hypertension confirmed a protective association for the highest compared to the lowest levels of legume intake [34]. The 1999–2002 National Health and Nutrition Examination Survey (NHANES) has shown that US adults who consumed approximately 1/2 cup per day (1 serving) of cooked dry beans or peas have lower odds of elevated blood pressure and a 1.7 mm/Hg lower mean systolic blood pressure than non-consumers [35]. Recent data from the EPIC Norfolk cohort also showed that legume consumption between 55–70 g/day was associated with a lower subsequent risk of hypertension [36].

All these previous observational studies were performed in people without diabetes; conversely, we performed our study in a large cohort of individuals with type 2 diabetes; it is of interest that our population was almost entirely under anti-hypertensive pharmacological treatment. This means that legume consumption can represent an additional tool—besides medications—to lower blood pressure, allowing a larger proportion of people with type 2 diabetes to reach the target.

Among several legume components potentially involved in the beneficial relationship observed with blood pressure in people with type 2 diabetes, like proteins, fibre, sodium, potassium, and polyphenols, only fibre showed a significant association with lower levels

of systolic and diastolic blood pressure according to our results. This can be due, at least in part, to the demonstrated role of fibre in reducing inflammation [37,38] and improving the endothelial function [39]. When the same analysis was performed adding as covariate legume consumption, only legumes were significantly and inversely associated with systolic and diastolic blood pressure, while the relationship with dietary fibre disappeared, thus demonstrating that the high fibre content of legumes plays an important role in the improvement of blood pressure control associated with consumption of this food group.

However, the possible contribution of other macro- and micro-nutrients present in legumes in explaining the association between consumption of this food group with a better blood pressure control should also be considered. In this respect, the vasodilatory effect of arginine in legume proteins via converting into nitric oxide has been demonstrated in animal and in vitro models and proposed as a potential mechanism driving the relationship observed [40,41]. Other bioactive peptides obtained from legume sources have been shown by in vitro studies to have high antioxidant capacity and the potential to inhibit the dipeptidyl peptidase-IV and the Angiotensin Converting Enzyme [42]. The inhibition of these two enzymes is the therapeutic target of drugs used for type 2 diabetes mellitus and for hypertension, while the antioxidant activity can prevent the development of several chronic diseases related to oxidative stress. However, more studies are needed to understand the bioavailability of these molecules, since the peptides released from beans can be inactivated during further digestion by peptidases in the intestinal epithelial cells.

The health benefits of polyphenols from legumes in relation to blood pressure control were summarized in a recent comprehensive review by Ganesan et al. [43] demonstrating that bioactive components, particularly phenolic compounds, play an important role in modulating vascular integrity and inflammatory markers, two factors associated with improvement of blood pressure.

There are several strengths of this study. This is the first large cohort of people with type 2 diabetes in which the links between legume intake and blood pressure were investigated in real-life conditions. Furthermore, the study was able to control for medications, metabolic parameters, and dietary components, which can play a significant modulating effect on blood pressure. Finally, the use of a central laboratory for biochemical analyses ensures good quality standards for the measurements.

Our study also has limitations, the most relevant being the cross-sectional study design which does not enable the establishing of a cause–effect relationship. Residual confounding cannot be ruled out, even though our findings are independent of (1) other dietary items and their components, such as mono- and poly-unsaturated fatty acids, glycaemic load, sodium, and alcohol intake; and (2) the main potential confounders, such as age, BMI, and smoking habit, were controlled. Furthermore, we did not analyse the role of different types of legumes consumed, as legume intake was aggregated into total amount. In this regard, we cannot exclude that different types of legumes may have different effects due to their distinct nutritional characteristics. Finally, we cannot exclude that individuals eating a healthier diet and consuming less alcohol may be more adherent to their medication.

5. Conclusions

Our findings in people with type 2 diabetes showed that the association of legume consumption with lower systolic and diastolic blood pressure values is already present for an intake of approximately three servings per week, which is feasible and in line with the current dietary recommendations for people with diabetes. This gives further support to current dietary guidelines in recommending the use of legumes (beans, peas, chickpeas, and lentils) for people with type 2 diabetes.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/nu15132895/s1>. Table S1. General characteristics of the study participants ($N = 1897$); Table S2. General characteristics and metabolic profile of the study participants according to the sex-specific quartiles of legume intake (g/1000 kcal /day)).

Author Contributions: Conceptualization, M.V., G.R., O.V. and M.M.; methodology, M.V.; formal analysis, M.V.; data curation, M.V., S.S., V.K. and M.M.; writing—original draft preparation, M.V., A.G. and E.M.; writing—review and editing, A.A.R., G.R., O.V., and M.M.; supervision, G.R. and O.V.; funding acquisition, O.V. All authors have read and agreed to the published version of the manuscript.

Funding: The study was supported by the Italian Medicines Agency (AIFA) within the Independent Drug Research Program—contract number FARM6T9CET—and by Diabete Ricerca, the non-profit Research Foundation of the Italian Diabetes Society. The funding agency played no role in the study design; in the collection, analysis, and interpretation of data; in the writing of the manuscript; or in the decision to submit the manuscript for publication.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the FEDERICO II UNIVERSITY Ethics Committee (protocol code: 123/08; date of approval: 21 January 2008).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Not applicable.

Acknowledgments: The participation of the patients in the study is gratefully acknowledged. We thank all the investigators and the dietitians in the TOSCA.IT centres for their excellent cooperation. We are also indebted to the administrative personnel of the Italian Diabetes Society (SID) for their support.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Lackland, D.T.; Weber, M.A. Global Burden of Cardiovascular Disease and Stroke: Hypertension at the Core. *Can. J. Cardiol.* **2015**, *31*, 569–571. <https://doi.org/10.1016/j.cjca.2015.01.009>.
2. Al-Makki, A.; DiPette, D.; Whelton, P.K.; Murad, M.H.; Mustafa, R.A.; Acharya, S.; Beheiry, H.M.; Champagne, B.; Connell, K.; Cooney, M.T.; et al. Hypertension Pharmacological Treatment in Adults: A World Health Organization Guideline Executive Summary. *Hypertension* **2022**, *79*, 293–301. <https://doi.org/10.1161/hypertensionaha.121.18192>.
3. Kabakov, E.; Norymberg, C.; Osher, E.; Koffler, M.; Tordjman, K.; Greenman, Y.; Stern, N. Prevalence of hypertension in type 2 diabetes mellitus: Impact of the tightening definition of high blood pressure and association with confounding risk factors. *J. Cardiometa. Syndr.* **2006**, *1*, 95–101. <https://doi.org/10.1111/j.1559-4564.2006.05513.x>.
4. Klein, R.; Klein, B.E.K.; Lee, K.E.; Cruickshanks, K.J.; Moss, S.E. The incidence of hypertension in insulin-dependent diabetes. *Arch. Intern. Med.* **1996**, *156*, 622–627. <https://doi.org/10.1001/archinte.156.6.622>.
5. Haffner, S.M.; Lehto, S.; Rönkä, T.; Pyörälä, K.; Laakso, M. Mortality from Coronary Heart Disease in Subjects with Type 2 Diabetes and in Nondiabetic Subjects with and without Prior Myocardial Infarction. *N. Engl. J. Med.* **1998**, *339*, 229–234. <https://doi.org/10.1056/nejm199807233390404>.
6. Adlerberth, A.M.; Rosengren, A.; Wilhelmsen, L.J. Diabetes and long-term risk of mortality from coronary and other causes in middle-aged Swedish men. A general population study. *Diabetes Care* **1998**, *21*, 539–545. doi: 10.2337/diacare.21.4.539.
7. Stamler, J.; Vaccaro, O.; Neaton, J.D.; Wentworth, D. Diabetes, Other Risk Factors, and 12-Yr Cardiovascular Mortality for Men Screened in the Multiple Risk Factor Intervention Trial. *Diabetes Care* **1993**, *16*, 434–444. <https://doi.org/10.2337/diacare.16.2.434>.
8. Adler, A.I.; Stratton, I.M.; Neil, H.A.W.; Yudkin, J.S.; Matthews, D.R.; Cull, C.A.; Wright, A.D.; Turner, R.C.; Holman, R.R. Association of systolic blood pressure with macrovascular and microvascular complications of type 2 diabetes (UKPDS 36): Prospective observational study. *Br. Med. J.* **2000**, *321*, 412–419. <https://doi.org/10.1136/bmj.321.7258.412>.
9. Turner, R.; Holman, R.; Stratton, I.; Cull, C.; Frighi, V.; Manley, S.; Matthews, D.; Neil, A.; McElroy, H.; Kohner, E.; et al. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. *Br. Med. J.* **1998**, *317*, 703–713. <https://doi.org/10.1136/bmj.317.7160.703>.
10. De Pergola, G.; D’alessandro, A. Influence of Mediterranean Diet on Blood Pressure. *Nutrients* **2018**, *10*, 1700.
11. Filippou, C.D.; Thomopoulos, C.G.; Kouremeti, M.M.; Sotiropoulou, L.I.; Nihoyannopoulos, P.I.; Tousoulis, D.M.; Tsioufis, C.P. Mediterranean diet and blood pressure reduction in adults with and without hypertension: A systematic review and meta-analysis of randomized controlled trials. *Clin. Nutr.* **2021**, *40*, 3191–3200. <https://doi.org/10.1016/j.clnu.2021.01.030>.
12. Lonnie, M.; Johnstone, A.M. The public health rationale for promoting plant protein as an important part of a sustainable and healthy diet. *Nutr. Bull.* **2020**, *45*, 281–293. <https://doi.org/10.1111/mbu.12453>.
13. Vogt, T.M.; Appel, L.J.; Obarzanek, E.; Moore, T.J.; Vollmer, W.M.; Svetkey, L.P.; Sacks, F.M.; Bray, G.A.; Cutler, J.A.; Windhauser, M.M.; et al. Dietary Approaches to Stop Hypertension: Rationale, Design, and Methods. *J. Am. Diet. Assoc.* **1999**, *99*, S12–S18. [https://doi.org/10.1016/s0002-8223\(99\)00411-3](https://doi.org/10.1016/s0002-8223(99)00411-3).

14. Sacks, F.M.; Svetkey, L.P.; Vollmer, W.M.; Appel, L.J.; Bray, G.A.; Harsha, D.; Obarzanek, E.; Conlin, P.R.; Miller, E.R.; Simons-Morton, D.G.; et al. Effects on Blood Pressure of Reduced Dietary Sodium and the Dietary Approaches to Stop Hypertension (DASH) Diet. *N. Engl. J. Med.* **2001**, *344*, 3–10. <https://doi.org/10.1056/nejm200101043440101>.
15. Ozemek, C.; Laddu, D.R.; Arena, R.; Lavie, C.J. The role of diet for prevention and management of hypertension. *Curr. Opin. Cardiol.* **2018**, *33*, 388–393. <https://doi.org/10.1097/hco.0000000000000532>.
16. Willett, W.; Rockström, J.; Loken, B.; Springmann, M.; Lang, T.; Vermeulen, S.; Garnett, T.; Tilman, D.; DeClerck, F.; Wood, A.; et al. Food in the Anthropocene: The EAT–Lancet Commission on healthy diets from sustainable food systems. *Lancet* **2019**, *393*, 447–492. [https://doi.org/10.1016/s0140-6736\(18\)31788-4](https://doi.org/10.1016/s0140-6736(18)31788-4).
17. Jenkins, D.J.A.; Kendall, C.W.C.; Augustin, L.S.A.; Mitchell, S.; Sahye-Pudaruth, S.; Mejia, S.B.; Chiavaroli, L.; Mirrahimi, A.; Ireland, C.; Bashyam, B.; et al. Effect of Legumes as Part of a Low Glycemic Index Diet on Glycemic Control and Cardiovascular Risk Factors in Type 2 Diabetes Mellitus: A Randomized Controlled Trial. *Arch. Intern. Med.* **2012**, *172*, 1653–1660. <https://doi.org/10.1001/2013.jamainternmed.70>.
18. Simpson, H.C.R.; Lousley, S.; Geekie, M.; Simpson, R.W.; Carter, R.D.; Hockaday, T.D.R.; Mann, J.I. A High Carbohydrate Leguminous Fibre Diet Improves All Aspects of Diabetic Control. *Lancet* **1981**, *317*, 1–5. [https://doi.org/10.1016/s0140-6736\(81\)90112-4](https://doi.org/10.1016/s0140-6736(81)90112-4).
19. Hosseinpour-Niazi, S.; Mirmiran, P.; Hedayati, M.; Azizi, F. Substitution of red meat with legumes in the therapeutic lifestyle change diet based on dietary advice improves cardiometabolic risk factors in overweight type 2 diabetes patients: A cross-over randomized clinical trial. *Eur. J. Clin. Nutr.* **2015**, *69*, 592–597. <https://doi.org/10.1038/ejcn.2014.228>.
20. Vaccaro, O.; Masulli, M.; Bonora, E.; Del Prato, S.; Giorda, C.B.; Maggioni, A.P.; Mocarelli, P.; Nicolucci, A.; Rivellesse, A.A.; Squatrito, S.; et al. Addition of Either Pioglitazone or a Sulfonylurea in Type 2 Diabetic Patients Inadequately Controlled with Metformin Alone: Impact on Cardiovascular Events: A Randomized Controlled Trial. *Nutr. Metab. Cardiovasc. Dis.* **2012**, *22*, 997–1006. <https://doi.org/10.1016/j.numecd.2012.09.003>.
21. Vaccaro, O.; Masulli, M.; Nicolucci, A.; Bonora, E.; Del Prato, S.; Maggioni, A.P.; Rivellesse, A.A.; Squatrito, S.; Giorda, C.B.; Sesti, G.; et al. Effects on the Incidence of Cardiovascular Events of the Addition of Pioglitazone versus Sulfonylureas in Patients with Type 2 Diabetes Inadequately Controlled with Metformin (TOSCA.IT): A Randomised, Multicentre Trial. *Lancet Diabetes Endocrinol.* **2017**, *5*, 887–897. [https://doi.org/10.1016/s2213-8587\(17\)30317-0](https://doi.org/10.1016/s2213-8587(17)30317-0).
22. Unger, T.; Borghi, C.; Charchar, F.; Khan, N.A.; Poulter, N.R.; Prabhakaran, D.; Ramirez, A.; Schlaich, M.; Stergiou, G.S.; Tomaszewski, M.; et al. Hypertension: New Guidelines from the International Society of Hypertension. *Am. Fam. Physician* **2021**, *103*, 763–765.
23. Friedewald, W.T.; Levy, R.I.; Fredrickson, D.S. Estimation of the Concentration of Low-Density Lipoprotein Cholesterol in Plasma, Without Use of the Preparative Ultracentrifuge. *Clin. Chem.* **1972**, *18*, 499–502. <https://doi.org/10.1093/clinchem/18.6.499>.
24. Pisani, P.; Faggiano, F.; Krogh, V.; Palli, D.; Vineis, P.; Berrino, F. Relative Validity and Reproducibility of a Food Frequency Dietary Questionnaire for Use in the Italian EPIC Centres. *Int. J. Epidemiol.* **1997**, *26*, 152–160. https://doi.org/10.1093/ije/26.suppl_1.s152.
25. Pala, V.; Sieri, S.; Palli, D.; Salvini, S.; Berrino, F.; Bellegotti, M.; Frasca, G.; Tumino, R.; Sacerdote, C.; Fiorini, L.; et al. Diet in the Italian Epic Cohorts: Presentation of Data and Methodological Issues. *Tumori J.* **2003**, *89*, 594–607. <https://doi.org/10.1177/030089160308900603>.
26. Bhagwat, S.; Haytowitz, D.B. *USDA Database for the Isoflavone Content of Selected Foods*; U.S. Department of Agriculture, Agricultural Research Service: Washington, DC, USA, 2015.
27. Haytowitz, D.; Wu, X.; Bhagwat, S. *USDA Database for the Flavonoid Content of Selected Foods, Release 3.3*; USDA Agricultural Research Service: Washington, DC, USA, 2018.
28. Rothwell, J.A.; Pérez-Jiménez, J.; Neveu, V.; Medina-Remón, A.; M'Hiri, N.; García-Lobato, P.; Manach, C.; Knox, C.; Eisner, R.; Wishart, D.S.; et al. Phenol-Explorer 3.0: A major update of the Phenol-Explorer database to incorporate data on the effects of food processing on polyphenol content. *Database* **2013**, *2013*, bat070. <https://doi.org/10.1093/database/bat070>.
29. Neveu, V.; Pérez-Jiménez, J.; Vos, F.; Crespy, V.; du Chaffaut, L.; Mennen, L.; Knox, C.; Eisner, R.; Cruz, J.; Wishart, D.; et al. Phenol-Explorer: An online comprehensive database on polyphenol contents in foods. *Database* **2010**, *2010*, bap024. <https://doi.org/10.1093/database/bap024>.
30. Jayalath, V.H.; de Souza, R.J.; Sievenpiper, J.L.; Ha, V.; Chiavaroli, L.; Mirrahimi, A.; Di Buono, M.; Bernstein, A.M.; Leiter, L.A.; Kris-Etherton, P.M.; et al. Effect of Dietary Pulses on Blood Pressure: A Systematic Review and Meta-analysis of Controlled Feeding Trials. *Am. J. Hypertens.* **2014**, *27*, 56–64. <https://doi.org/10.1093/ajh/hpt155>.
31. Ley, S.H.; Hamdy, O.; Mohan, V.; Hu, F.B. Prevention and management of type 2 diabetes: Dietary components and nutritional strategies. *Lancet* **2014**, *383*, 1999–2007. [https://doi.org/10.1016/s0140-6736\(14\)60613-9](https://doi.org/10.1016/s0140-6736(14)60613-9).
32. American Diabetes Association Professional Practice Committee; Draznin, B.; Aroda, V.R.; Bakris, G.; Benson, G.; Brown, F.M.; Freeman, R.; Green, J.; Huang, E.; Isaacs, D.; et al. 8. Obesity and Weight Management for the Prevention and Treatment of Type 2 Diabetes: Standards of Medical Care in Diabetes—2022. *Diabetes Care* **2022**, *45*, S113–S124. <https://doi.org/10.2337/dc22-s008>.
33. Schwingshackl, L.; Schwedhelm, C.; Hoffmann, G.; Knüppel, S.; Iqbal, K.; Andriolo, V.; Bechthold, A.; Schlesinger, S.; Boeing, H. Food Groups and Risk of Hypertension: A Systematic Review and Dose-Response Meta-Analysis of Prospective Studies. *Adv. Nutr.* **2017**, *8*, 793–803. <https://doi.org/10.3945/an.117.017178>.
34. Vigiouliou, E.; Glenn, A.J.; Nishi, S.K.; Chiavaroli, L.; Seider, M.; Khan, T.; Bonaccio, M.; Iacoviello, L.; Mejia, S.B.; Jenkins, D.J.A.; et al. Associations between Dietary Pulses Alone or with Other Legumes and Cardiometabolic Disease Outcomes: An Umbrella

- Review and Updated Systematic Review and Meta-analysis of Prospective Cohort Studies. *Adv. Nutr.* **2019**, *10*, S308–S319. <https://doi.org/10.1093/advances/nmz113>.
35. Papanikolaou, Y.; Fulgoni, V.L. Bean Consumption Is Associated with Greater Nutrient Intake, Reduced Systolic Blood Pressure, Lower Body Weight, and a Smaller Waist Circumference in Adults: Results from the National Health and Nutrition Examination Survey 1999–2002. *J. Am. Coll. Nutr.* **2008**, *27*, 569–576. <https://doi.org/10.1080/07315724.2008.10719740>.
 36. Hartley, M.; Fyfe, C.L.; Wareham, N.J.; Khaw, K.-T.; Johnstone, A.M.; Myint, P.K. Association between Legume Consumption and Risk of Hypertension in the European Prospective Investigation into Cancer and Nutrition (EPIC)-Norfolk Cohort. *Nutrients* **2022**, *14*, 3363. <https://doi.org/10.3390/nu14163363>.
 37. Ma, Y.; Griffith, J.A.; Chasan-Taber, L.; Olendzki, B.C.; Jackson, E.; Stanek, E.J.; Li, W.; Pagoto, S.L.; Hafner, A.R.; Ockene, I.S. Association between dietary fiber and serum C-reactive protein. *Am. J. Clin. Nutr.* **2006**, *83*, 760–766. <https://doi.org/10.1093/ajcn/83.4.760>.
 38. Krishnamurthy, V.M.R.; Wei, G.; Baird, B.C.; Murtaugh, M.; Chonchol, M.B.; Raphael, K.L.; Greene, T.; Beddhu, S. High dietary fiber intake is associated with decreased inflammation and all-cause mortality in patients with chronic kidney disease. *Kidney Int.* **2012**, *81*, 300–306. <https://doi.org/10.1038/ki.2011.355>.
 39. Brock, D.W.; Davis, C.K.; Irving, B.A.; Rodriguez, J.; Barrett, E.J.; Weltman, A.; Taylor, A.G.; Gaesser, G.A. A High-Carbohydrate, High-Fiber Meal Improves Endothelial Function in Adults with the Metabolic Syndrome. *Diabetes Care* **2006**, *29*, 2313–2315. <https://doi.org/10.2337/dc06-0917>.
 40. Chuman, H.; Sugimoto, T.; Nao-I, N. Vasodilatory effect of L-arginine on isolated rabbit and human posterior ciliary arteries in vitro and increased optic disc blood flow in vivo. *Graefe's Arch. Clin. Exp. Ophthalmol.* **2017**, *255*, 2381–2388. <https://doi.org/10.1007/s00417-017-3824-5>.
 41. Zhou, C.; Wu, Q.; Ye, Z.; Liu, M.; Zhang, Z.; Zhang, Y.; Li, H.; He, P.; Li, Q.; Liu, C.; et al. Inverse Association Between Variety of Proteins with Appropriate Quantity from Different Food Sources and New-Onset Hypertension. *Hypertension* **2022**, *79*, 1017–1027. <https://doi.org/10.1161/hypertensionaha.121.18222>.
 42. Garcia, B.D.F.; de Barros, M.; Rocha, T.D.S. Bioactive peptides from beans with the potential to decrease the risk of developing noncommunicable chronic diseases. *Crit. Rev. Food Sci. Nutr.* **2021**, *61*, 2003–2021. <https://doi.org/10.1080/10408398.2020.1768047>.
 43. Ganesan, K.; Xu, B. Polyphenol-Rich Dry Common Beans (*Phaseolus vulgaris* L.) and Their Health Benefits. *Int. J. Mol. Sci.* **2017**, *18*, 2331. <https://doi.org/10.3390/ijms18112331>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.