

Article

Combined Aerobic Training and Mediterranean Diet Is Not Associated with a Lower Prevalence of Sarcopenia in Italian Older Adults

Hélio José Coelho-Júnior ^{1,*}, Riccardo Calvani ^{1,2}, Anna Picca ^{2,3}, Stefano Cacciatore ¹, Matteo Tosato ², Francesco Landi ^{1,2} and Emanuele Marzetti ^{1,2,*}

¹ Department of Geriatrics, Orthopedics, and Rheumatology, Università Cattolica del Sacro Cuore, 00168 Rome, Italy; riccardo.calvani@unicatt.it (R.C.); francesco.landi@unicatt.it (F.L.)

² Fondazione Policlinico Universitario “A. Gemelli”, IRCCS, 00168 Rome, Italy; picca@lum.it (A.P.); matteo.tosato@policlinicogemelli.it (M.T.)

³ Department of Medicine and Surgery, LUM University, 70100 Casamassima, Italy

* Correspondence: coelhojunior@hotmail.com.br (H.J.C.-J.); emanuele.marzetti@policlinicogemelli.it (E.M.); Tel.: +39-(06)-3015-4859 (H.J.C.-J. and E.M.)

Abstract: Previous studies found a lower prevalence of sarcopenia in older adults engaged in regular aerobic training (AT) or with greater adherence to a Mediterranean (MED) diet. However, the effect of their combination on sarcopenia indices is unknown. The present study tested the association between AT plus a MED diet and the presence of sarcopenia and its defining elements in a sample of Italian older adults enrolled in the Longevity Check-up 7+ (Lookup 7+) project. Analyses were conducted in participants 65+ years, with a body mass index of at least 18.5 kg/m², engaged in regular AT, and without missing information for the variables of interest. MED diet adherence was evaluated via a modified version of the MEDI-LITE score and categorized as low, moderate, or high. The presence of sarcopenia was established by handgrip strength and appendicular skeletal muscle mass (ASM) values below sex-specific cut-points recommended by the European Working Group on Sarcopenia in Older People 2. Data from 491 older adults were analyzed for the present study. The mean age was 72.7 ± 5.7 years, and 185 (37.7%) were women. MED diet adherence was low in 59 (12.0%) participants, moderate in 283 (57.6%), and high in 149 (30.3%). Sarcopenia was identified in 26 participants (5.3%), with no differences across MED diet adherence groups. The results of binary logistic regression showed no significant associations between AT plus adherence to a MED diet and dynapenia, low ASM, or sarcopenia. The findings of the present study indicate that the combination of AT with a MED diet is not associated with a lower probability of sarcopenia or its defining elements in Italian older adults enrolled in Lookup 7+. Further research is warranted to establish whether exercise frequency, volume, intensity, and length of engagement in AT impact the association between MED diet and sarcopenia.

Keywords: muscle strength; frailty; exercise training; nutrition; diet patterns; elderly

Citation: Coelho-Júnior, H.J.; Calvani, R.; Picca, A.; Cacciatore, S.; Tosato, M.; Landi, F.; Marzetti, E. Combined Aerobic Training and Mediterranean Diet Is Not Associated with a Lower Prevalence of Sarcopenia in Italian Older Adults. *Nutrients* **2023**, *15*, 2963. <https://doi.org/10.3390/nu15132963>

Academic Editor: Motoyuki Iemitsu

Received: 9 June 2023

Accepted: 28 June 2023

Published: 29 June 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Sarcopenia is a neuromuscular disease that involves a supraphysiologic loss of muscle mass and strength [1,2], while a reduced physical function is an indicator of severity [1,2]. The prevalence of sarcopenia increases with age and exceeds 20% in those 65+ years [3]. Sarcopenia is associated with numerous negative health outcomes, including falls, hospitalization, admission to residential care, and death [4]. The progression of sarcopenia is also frequently associated with the occurrence of other health conditions (e.g., malnutrition, frailty, high blood pressure) [4–7].

Exercise training, a combination of organized motor tasks that aims to develop or maintain one or more attributes of physical fitness [8], is indicated as a first-line intervention to counteract sarcopenia development and progression [9–11]. Resistance training, a form of exercise in which muscles contract to support or move a load [8], has been shown to enhance sarcopenia-related parameters in older persons [12,13]. A recent network meta-analysis found that older adults with sarcopenia who engaged in resistance training programs had greater gains in muscle strength and function compared with other exercise modalities [14].

Aerobic training (AT), a form of exercise where muscle contractions are maintained for extended periods of time [8], has also been tested against sarcopenia. Although promising findings have recently been reported [15], pooled analyses indicated limited effects of this type of intervention on sarcopenia-related parameters [16]. The subject deserves further exploration, given that AT is an effective strategy for the prevention and management of cardiovascular and metabolic diseases [17–20]. Furthermore, land- and water-based AT programs can be integrated into ludic activities (e.g., water volleyball) [21], which might foster adherence and participant well-being.

The Mediterranean (MED) diet refers to a dietary pattern passed down through many generations resulting from centuries of cultural interactions among inhabitants of the MED region [22,23]. The traditional MED diet pyramid involves a high intake of plant foods (e.g., fruits, vegetables, legumes, grains, nuts), moderate amounts of eggs, seafood, and dairy products, and occasional consumption of meat [22,23]. The cardioprotective properties of the MED diet have long been recognized [24,25]. More recently, a higher MED diet adherence has been associated with better physical performance in older adults [26]. In addition, the MED diet has been negatively associated with the presence of sarcopenia [27]. These premises suggest that older adults who practice AT and have moderate or high adherence to a MED diet might show a lower prevalence of sarcopenia relative to physically active peers with low MED diet adherence.

The Longevity Check-Up 7+ (Lookup 7+) project is an ongoing study that started in June 2015. The objective of Lookup 7+ is to promote the adoption of healthy lifestyles by the general Italian population to lower the prevalence of risk factors for chronic diseases. The Lookup 7+ database offered the possibility to explore, in a “real world” context, several health parameters relevant to clinicians and researchers, including the prevalence of cardiometabolic risk factors [28] and their association with physical function [29] and anthropometric measures [30,31]. The adherence to physical activity recommendations and the practice of different types of exercise training have also been examined [32].

Two prior studies by our group explored the relationship between physical exercise or adherence to a MED diet and sarcopenia in older adults enrolled in Lookup 7+ [15,27]. In one study, we investigated the association between adherence to specific exercise modalities and the presence of sarcopenia or severe sarcopenia [15]. Sarcopenia was operationalized based on the cooccurrence of low appendicular skeletal muscle mass (ASM) and low muscle strength of upper or lower extremities estimated by isometric handgrip (IHG) strength and five-time sit-to-stand tests, respectively. Information on self-reported mobility limitations was used to define the severity of sarcopenia. Analyses were conducted in 3289 participants (mean age: 72.7 years, standard deviation (SD) \pm 5.7 years; 55.2% women) and showed that engagement in AT was negatively associated with sarcopenia in both men and women, with sex-specific patterns [15]. In another study, we explored whether the level of adherence to a MED diet was associated with probable sarcopenia, operationalized as an IHG strength below sex-specific cut-points recommended by the European Working Group on Sarcopenia in Older People 2 (EWGSOP2) [2,27]. We found that a high MED diet adherence was indeed associated with lower odds of probable sarcopenia in 2963 older adults (mean age 72.8 years, SD \pm 5.7 years; 54.4% women).

Altogether, these observations suggest that the negative association between AT and sarcopenia might be amplified in older adults with a high MED diet adherence. This possibility was explored in the present study by analyzing associations between the practice

of AT plus different levels of adherence to a MED diet and low IHG strength, low ASM, and sarcopenia in older adults enrolled in Lookup 7+.

2. Materials and Methods

Data analyzed for the present study were obtained from the Lookup 7+ database. Participants were recruited in unconventional contexts, including exhibitions, shopping centers, and social gatherings, or during prevention campaigns organized by our university. This recruitment strategy was adopted to include relatively unselected participants outside of the usual contexts for medical care or scientific research. The manuscript was written in accordance with the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines for observational studies [33].

2.1. Participants

From 1 June 2015 to 31 March 2023, 13,515 adults living in the community were enrolled in the Lookup 7+ project. Exclusion criteria were self-reported pregnancy, difficulty or incapacity to perform the physical function tests required by the protocol, and refusal or inability to provide written informed consent. For the present study, analyses were performed using information from participants ($n = 491$) who were 65 or older, had a body mass index (BMI) of at least 18.5 kg/m², practiced regular AT, and had no missing data for the variables of interest.

2.2. Data Collection

Data on dietary and exercise habits and other lifestyle behaviors were collected using a structured interview. Smoking status was divided into current (defined as having smoked 100 or more cigarettes in their lifetime and still doing so) and non-current smoker categories. A standard stadiometer and an analog scale were used to measure standing height and body weight, respectively. The BMI was determined as the ratio between body weight (kg) and the square of height (m²). Regular physical activity was defined as engaging in leisure activities at least twice a week, for a minimum of 30 minutes each time, in the prior year [32]. Regular engagement in AT was operationalized as the practice of aerobic activities (e.g., running, swimming) at least twice weekly for a minimum of 30 minutes per session during the previous year [32,34]. The interview did not include questions about exercise frequency, volume, intensity, or length of engagement in AT.

2.3. Mediterranean Diet Adherence

Nutrition data were gathered through the administration of a food frequency questionnaire. A modified version of the MEDI-LITE score was used to evaluate adherence to a MED diet, as previously described [27]. The instrument is based on the frequency of consumption of nine food groups: (1) fruit, (2) vegetables, (3) cereals, (4) legumes, (5) fish and seafood, (6) meat and derivatives, (7) dairy products, (8) alcohol, (9) olive oil. Food items typically included in a MED diet (i.e., fruit, vegetables, grains, legumes, and fish) received a score from 2 (highest intake) to 0 (lowest consumption). Meat, meat derivatives, and dairy products, which are not included in a typical MED diet, were given 0 points for the highest intake, 1 point for intermediate consumption, and 2 points for the lowest consumption. For olive oil, a score of 2 points was assigned to daily consumption, 1 point to frequent use, and 0 points to infrequent use. Information on alcohol intake was not collected, and the item was not scored. Thus, the maximum MEDI-LITE score was 16 rather than 18. A high MED diet adherence was operationalized as a MEDI-LITE score of 12 or greater. Moderate MED diet adherence was operationalized as a score between 9 and 11, while low adherence was defined as a score of 8 or lower.

2.4. Operationalization of Sarcopenia

Sarcopenia was operationalized as the coexistence of dynapenia and low ASM [1,2]. IHG was used as a measure of muscle strength. Participants took the IHG test while seated in a chair with their shoulders in a neutral position. The thumb was up, the hand was neutral, and the elbow was flexed at 90 degrees close to the body on the arm that was being assessed. The maximum contraction was measured using a portable hydraulic dynamometer (North Coast Hydraulic Hand Dynamometer; North Coast Medical, Inc., Morgan Hill, CA, USA). One familiarization trial was performed before the test. For the analysis, the greater reading (in kg) from two IHG measurements was considered. ASM was estimated from the calf circumference of the dominant leg. The measurement was made at the largest circumference between the ankle and knee joints with participants sitting on a chair. ASM was calculated according to the formula [35]:

$$\text{ASM} = -10.427 + (\text{calf circumference (cm)} \times 0.768) - (\text{age} \times 0.029) + (\text{sex (male} = 1, \text{female} = 0)) \times 7.523$$

Sex-specific cut-points by EWGSOP2 were used to categorize muscle strength and ASM as low [1,2].

2.5. Statistical Analysis

The Shapiro–Wilk test was employed to confirm that the data distribution was normal. Categorical variables are given as exact numbers and percentages, while continuous variables are reported as mean \pm SD. One-way analysis of variance (ANOVA) was used to assess differences among MED diet adherence groups. Differences in categorical variables were examined by χ^2 statistics. Binary logistic regression was conducted to test the association between AT plus moderate or high MED diet adherence and sarcopenia status and related parameters. The final model was adjusted for age, sex, BMI, walking activity, and smoking status. All tests were two-sided, with significance set at 5% ($p < 0.05$). Confidence intervals (CIs) that contained the number of 1 were not statistically significant. All analyses were performed using the SPSS software (version 23.0, SPSS Inc., Chicago, IL, USA).

3. Results

The main characteristics of study participants according to categories of MED diet adherence are listed in Table 1. The mean age of the 491 participants practicing AT was 72.7 ± 5.7 years, and 185 (37.7%) were women. Mean BMI values indicated that participants were in the high-normal range. The average IHG strength values were higher than the EWGSOP2 cut points for sarcopenia, while ASM was borderline. The distribution of participants with sarcopenia was not significantly different across MED diet adherence groups.

Table 1. Main characteristics of study participants according to categories of adherence to a Mediterranean diet ($n = 491$).

	AT Plus Low MED Diet Adherence ($n = 59$)	AT Plus Moderate MED Diet Adherence ($n = 283$)	AT Plus High MED Diet Adherence ($n = 149$)	p Value
Age, years	71.3 ± 4.8	71.3 ± 5.3	72.2 ± 5.2	0.842
Sex, female	9, 1.8	49, 10.0	17, 3.5	0.199
BMI, kg/m ²	26.1 ± 3.5	25.4 ± 3.4	24.9 ± 3.5	0.585
IHG strength, kg	30.2 ± 8.5	31.4 ± 10.9	30.9 ± 10.3	1.000
ASM, kg/m ²	19.7 ± 5.3	19.5 ± 4.8	18.5 ± 4.8	0.344
Smoking, yes	21, 4.3	99, 20.2	65, 13.2	0.268
Walking, yes	56, 11.5	275, 56.4	144, 29.5	0.917

Dynapenia, yes	6, 1.2	30, 6.2	10, 2.1	0.422
Low ASM loss, yes	22, 4.5	90, 18.5	60, 12.3	0.169
Sarcopenia, yes	3, 0.9	15, 4.7	8, 2.5	0.970

Continuous data are shown as mean \pm standard deviation, while binary data are reported as number, %. Abbreviations: ASM, appendicular skeletal muscle; BMI, body mass index; IHG, isometric handgrip.

Table 2 shows the results of the binary logistic regression for the relationship between AT plus MED adherence and sarcopenia indices. No significant associations were observed between MED adherence and dynapenia, low ASM, or sarcopenia. **Table 2.** Association between adherence to a Mediterranean diet plus aerobic training and sarcopenia indices.

	Unadjusted OR* (95% CI)	Adjusted OR (95% CI)		Unadjusted OR* (95% CI)	Adjusted OR (95% CI)		Unadjusted OR* (95% CI)	Adjusted OR (95% CI)
Dynapenia								
AT plus Low MED diet adherence	1.00 (Reference)	1.00 (Reference)	AT plus Low MED diet adherence	1.00 (Reference)	1.00 (Reference)	AT plus Moderate MED diet adherence	1.000 (Reference)	1.00 (Reference)
AT plus Moderate MED diet adherence	0.96 (0.38, 2.44)	1.01 (0.38, 2.66)	AT plus High MED diet adherence	1.58 (0.54, 4.56)	2.43 (0.75, 7.86)	AT plus High MED diet adherence	1.63 (0.77, 3.43)	1.99 (0.77, 3.43)
Low appendicular skeletal muscle mass								
AT plus Low MED diet adherence	1.00 (Reference)	1.00 (Reference)	AT plus Low MED diet adherence	1.00 (Reference)	1.00 (Reference)	AT plus Moderate MED diet adherence	1.00 (Reference)	1.00 (Reference)
AT plus Moderate MED diet adherence	1.26 (0.70, 2.26)	1.64 (0.69, 3.89)	AT plus High MED diet adherence	0.85 (0.45, 1.58)	1.52 (0.64, 3.62)	AT plus High MED diet adherence	0.67 (0.44, 1.02)	0.93 (0.51, 1.70)
Sarcopenia								
AT plus Low MED diet adherence	1.00 (Reference)	1.00 (Reference)	AT plus Low MED diet adherence	1.00 (Reference)	1.00 (Reference)	AT plus Moderate MED diet adherence	1.00 (Reference)	1.00 (Reference)
AT plus Moderate MED diet adherence	0.93 (0.25–3.42)	0.97 (0.19, 4.77)	AT plus High MED diet adherence	1.04 (0.26, 4.19)	0.29 (0.03, 2.78)	AT plus High MED diet adherence	1.11 (0.45, 2.73)	0.50 (0.14, 1.72)

* The model was adjusted model was adjusted for age, sex, body mass index, walking activity, and smoking status. Abbreviations: AT, aerobic training; CI, confidence interval, MED, Mediterranean, OR, odds ratio.

4. Discussion

In the present study, we used the Lookup 7+ database to explore whether the combination between AT and moderate/high adherence to a MED diet would be associated with lower odds of sarcopenia than AT with a low MED diet adherence. Our findings revealed no significant relationships.

Studies have investigated the effects of combining AT and various dietary patterns on sarcopenia-related parameters with conflicting results. Markofski et al. [36] observed greater improvements in leg isokinetic extension and muscle quality in apparently healthy, independent-living older adults after a 24-week moderate-intensity AT program combined with essential amino acid supplementation compared with exercise or supplementation alone. No differences between groups were observed for lean mass. In contrast, Li et al. [37] reported that an eight-week low-to-moderate AT program combined with isolated soy protein supplementation increased lean mass, but not physical performance in young women. Fairbairn et al. [38] did not observe significant improvements in

mobility in older women who adhered to a 24-week AT program plus docosahexaenoic acid supplementation. The results of a recent meta-analysis indicated that protein supplementation did not convey further gains in muscle strength or physical performance in endurance-trained older adults, although greater increases in muscle mass were observed [39].

The effects of a combined AT and MED diet on musculoskeletal parameters have been sparsely investigated, and, to the best of our knowledge, this is the first study that examined the association with sarcopenia. In one of the few investigations in the field, Pineda-Juárez et al. [40] reported that AT plus a MED diet had no effects on IHG strength in women with rheumatoid arthritis.

Our findings are somewhat unexpected, owing to the numerous studies that reported significant associations between adherence to a MED diet and measures of physical performance and muscle strength [26]. A possible explanation for our results lies in the fact that high adherence to a MED diet involves a moderate intake of high-protein food sources, such as fish (two to six servings a week), poultry, dairy products, and eggs (two to four servings a week), and a sparse ingestion of meat (≤ 2 servings a week) [22,41].

Protein intake stimulates muscle protein synthesis by providing amino acids to support muscle growth [42–44]. Essential amino acids that are not produced by the human body and need to be obtained through dietary protein are necessary to activate muscle anabolism [44,45]. Most of these anabolic effects are mediated by the action of branched-chain amino acids (i.e., isoleucine, leucine, and valine), especially leucine [44,46]. Indeed, several studies reported that leucine supplementation may improve muscle mass, strength, and function [47].

Meat, fish, eggs, and dairy products are known as “myoprotective whole foods” [48] due to their protein content and potential effects on sarcopenia-related parameters [49–52]. The ingestion of approximately 110 g of beef increased muscle protein synthesis by 51% relative to the basal state [53]. Furthermore, a small observational study reported that beef meat consumption was positively associated with muscle mass in older adults [54]. A pooled analysis found significant improvements in lower-limb muscle strength following beef protein supplementation [55]. Further to this point, Marcos-Pardo et al. [56] suggested that a limited consumption of red and processed meat could explain the lack of associations between high adherence to a MED diet, according to the Prevención con Dieta Mediterránea (PREDIMED) score, and sarcopenia-related parameters. The authors also noted that eating more than one serving of red meat, hamburger, sausage, or cold cuts each day was associated with preservation of muscle strength and mass in European middle-aged and older adults [56].

Interesting results have also been reported by studies that examined dairy products. Whey and casein are two milk proteins that differ in amino acid composition as well as digestion and absorption kinetics [57–60], with anabolic effects on muscle tissue. Whey protein has a high leucine content [57–60] and stimulates muscle protein synthesis to a greater extent than casein. Indeed, supplementation with whey protein increases lean body mass, mainly if combined with exercise training [61].

Regarding fish, rats fed fish-based protein had significant improvements in muscle diameter in both fast and slow fiber types [62,63]. Data from the Seniors-ENRICA study indicates that older adults with a greater consumption of fish or dairy products were at a lower risk of functional impairment than those with lower intakes [64]. Moreover, a high consumption of eggs was associated with a reduced prevalence of sarcopenia in older Korean adults [52]. Notably, a low consumption of the foods mentioned above might also contribute to the development of sarcopenia through an insufficient intake of other nutrients (e.g., vitamin D and magnesium) [65].

An opposing view is that the MED diet allows protein requirements to be met through the consumption of plant-based foods. The MED diet focuses on plant food sources instead of animal-based foods [22], reflected by the fact that vegetable foods are located at the basis of the MED diet pyramid [22]. An increasing number of studies have

investigated the association between plant-based diets and neuromuscular aspects. Coelho-Junior et al. [66] reported that, in well-functioning older adults, a high intake of plant-based protein was associated with faster walking speed than animal-derived protein. Yeung and Woo [67] added to these findings by reporting that a high vegetable protein intake was associated with lower declines in walking speed in older Chinese adults. Montiel-Rojas et al. [68] observed that favoring vegetable protein over animal-derived protein reduced the risk of sarcopenia in older European adults.

Such findings led experts in the field to propose that a high consumption of plant-based protein might properly stimulate muscle protein synthesis [69]. However, protein of plant origin is expected to provide a limited amount of essential amino acids, including leucine, and is characterized by a lower digestibility compared with protein from animal sources [70]. An experimental study reported that the ingestion of soy protein promoted lower muscle protein synthesis than whey protein in older men under resting conditions and after exercise, regardless of protein dose [71]. Furthermore, findings of the LifeAge Study indicated that the consumption of three or more portions of legumes per week was not associated with a better performance on IHG, sit-to-stand, gait speed, Timed “Up and Go!”, or Short Physical Performance Battery (SPPB) tests, greater muscle mass, or lower prevalence of sarcopenia in European middle-aged and older adults [56].

Altogether, the available evidence suggests that MED diet protein content might not be sufficient to improve sarcopenia parameters in older adults. Although some results in favor of vegetable protein have been observed in older adults [66–68], exercise-trained individuals may require larger amounts of protein than their physically inactive peers [72]. According to Camera [73], a high protein intake is not only required to properly stimulate muscle protein synthesis in aerobically trained individuals but is imperative to promote adequate recovery and performance, avoiding overtraining syndrome.

Another important aspect that might have influenced our results is lifestyle. The MED diet is a concept used to reflect the typical dietary habits observed in inhabitants of the MED basin, including Greece and South Italy [73]. This dietary pattern was mainly followed by people from rural societies and was associated with other potentially healthy behaviors [22,74]. Some authors have suggested that diet alone may not be sufficient for managing the complexity of chronic conditions, such as sarcopenia, and that the benefits of a MED diet might be conveyed by a combination of multiple healthy lifestyle factors [75,76]. Information on other lifestyle habits (e.g., social engagement, hydration, sleep time and quality, food preparation methods) were not collected in Lookup 7+ and should be analyzed in future studies.

The investigation of dietary patterns other than the MED diet is necessary to identify the optimal nutritional approach that amplifies the neuromuscular benefits of AT in older adults. However, the selection of possible candidates is not an easy task to accomplish. Results of a recent systematic review and meta-analysis indicated that healthy dietary patterns, created from established indexes or using a factor or cluster analysis, were not associated with better performance on IHG, sit-to-stand, balance, or SPPB tests, or with the prevalence of sarcopenia [77]. Although a higher adherence to healthy dietary patterns was linked to a lower reduction in gait speed, limitations in the studies included in the meta-analysis prevented the authors from conducting deeper analyses to identify the dietary pattern that conveyed the greatest benefits.

Healthy Eating Index (HEI), a set of widely accepted metrics that estimate diet quality according to the amount of culturally neutral foods (e.g., fruits and vegetables, whole and refined grains, added sugar) ingested per 1000 kcal [78], have been examined in relation to sarcopenia-related parameters. Parsons et al. [79] analyzed the database of the British Regional Heart Study (BRHS) to investigate the association between adherence to an HEI, composed of 86 food and drink items, and the development of mobility limitations (i.e., difficulty in walking 400 yards) in older men. Results indicated that men with greater adherence to the HEI at baseline were less likely to develop mobility limitations over 15 years of follow-up. In contrast, participants with a high-fat/low-fiber dietary

pattern at baseline showed a greater risk of incident mobility limitations at follow-up [79]. On the other hand, Chan et al. [80] did not find significant associations between the adherence to an HEI and the incidence of sarcopenia, defined according to the Asian Working Group for Sarcopenia (AWGS) criteria [2], during four years of follow-up in older Chinese adults living in the community. A posteriori analysis indicated that men who followed a “vegetables-fruits” or a “snacks-drinks-milk products” dietary pattern had a lower risk of incident sarcopenia than those with high adherence to other nutritional regimens [80].

Other studies have focused on region-specific dietary patterns. The Nordic Diet Score (NDS) is a type of HEI that involves foods commonly consumed in Nordic countries (i.e., Denmark, Norway, Sweden, Finland, and Iceland), such as berries, root and cruciferous vegetables, rye bread, and a high fish intake [78]. Perala et al. [81] examined the association between adherence to an NDS and the 10-year risk of dynapenia, assessed according to IHG and isometric leg strength tests, in 1094 Finnish older adults. The authors observed that each unit increase in the NDS was related to greater leg and IHG strength in women but not men [81].

Our study has limitations that deserve acknowledgment. First, participants were community-dwelling older Caucasian adults who attended specific events, and extrapolations to other cohorts should be considered with caution. Second, ASM was estimated using an anthropometric measure (i.e., calf circumference). Third, the MED diet was operationalized using an adapted version of the MEDI-LITE score, and the possibility that results could be different using other validated questionnaires (e.g., MedDiet score, PREDIMED score) cannot be ruled out. Fourth, information on chronic diseases (e.g., osteoarthritis) or medications (e.g., corticosteroids) that could impact musculoskeletal health was not collected. Fifth, exercise training variables (e.g., frequency, volume, intensity, length of engagement in AT) were not collected or controlled for. This limitation, albeit not negligible, is intrinsic to the design of the project and the unconventional settings where the survey is conducted. The collection of detailed information on lifestyle habits and medical history would substantially increase the duration of the assessments, which may deter many candidates from participating. Sixth, the use of IHG as an assessment tool for dynapenia has received some criticism [82], owing to its epidemiological and biomechanical aspects. Other tests, such as the sit-to-stand, seem to be a better proxy of muscle strength [82]. Hence, our results need to be confirmed in investigations using other measures of sarcopenia parameters. Seventh, the cross-sectional design of the study precludes the possibility of drawing cause-and-effect implications.

5. Conclusions

The findings of the present study indicate that the combination of AT and moderate or high adherence to a MED diet is not associated with a lower probability of sarcopenia in older Italian adults. Future investigations are warranted to explore whether other dietary regimens (e.g., high-protein diets) in combination with AT training offer therapeutic gain against sarcopenia. Further research is also necessary to establish whether exercise frequency, volume, intensity, and length of engagement in AT impact the association between MED diet and sarcopenia.

Author Contributions: Conceptualization, F.L., H.J.C.-J., E.M., R.C., A.P. and M.T.; Data curation, H.J.C.-J., E.M., R.C. and M.T.; Formal analysis, F.L., H.J.C.-J., S.C., E.M., R.C. and A.P.; Funding acquisition, F.L. and E.M.; Investigation, F.L., S.C., E.M., R.C., A.P. and M.T.; Methodology, F.L., H.J.C.-J., S.C., E.M. and A.P.; Resources, F.L.; Supervision, F.L. and E.M.; Writing—original draft, F.L., H.J.C.-J., S.C., E.M., R.C., A.P. and M.T.; Writing—review & editing, F.L., H.J.C.-J., E.M. and A.P. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the Università Cattolica del Sacro Cuore [D1.2020 and D1.2022], the Italian Ministry of Health [Ricerca Corrente 2023], the nonprofit research foundation “Centro Studi Achille e Linda Lorenzon” [N/A], and the European Commission—NextGeneration

EU (Age-IT spoke 3). The views and opinions expressed are only those of the authors and do not necessarily reflect those of the European Union or the European Commission. Neither the European Union nor the European Commission can be held responsible for them.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki and approved by the Institutional Review Board of Università Cattolica del Sacro Cuore (protocol #: A.1220/CE/2011; date of approval: 3 March 2014).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Data are available upon reasonable request from the corresponding authors.

Acknowledgments: We would like to thank Maria Banfi for her constructive comments.

Conflicts of Interest: The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Cruz-Jentoft, A.J.; Bahat, G.; Bauer, J.; Boirie, Y.; Bruyère, O.; Cederholm, T.; Cooper, C.; Landi, F.; Rolland, Y.; Sayer, A.A.; et al. Sarcopenia: Revised European Consensus on Definition and Diagnosis. *Age Ageing* **2019**, *48*, 16–31. <https://doi.org/10.1093/ageing/afy169>.
2. Chen, L.K.; Woo, J.; Assantachai, P.; Auyeung, T.W.; Chou, M.Y.; Iijima, K.; Jang, H.C.; Kang, L.; Kim, M.; Kim, S.; et al. Asian Working Group for Sarcopenia: 2019 Consensus Update on Sarcopenia Diagnosis and Treatment. *J. Am. Med. Dir. Assoc.* **2020**, *21*, 300–307.e2. <https://doi.org/10.1016/j.jamda.2019.12.012>.
3. Petermann-Rocha, F.; Balntzi, V.; Gray, S.R.; Lara, J.; Ho, F.K.; Pell, J.P.; Celis-Morales, C. Global Prevalence of Sarcopenia and Severe Sarcopenia: A Systematic Review and Meta-Analysis. *J. Cachexia Sarcopenia Muscle* **2022**, *13*, 86–99. <https://doi.org/10.1002/jcsm.12783>.
4. Chew, S.T.H.; Tey, S.L.; Yalawar, M.; Liu, Z.; Baggs, G.; How, C.H.; Cheong, M.; Chow, W.L.; Low, Y.L.; Huynh, D.T.T.; et al. Prevalence and Associated Factors of Sarcopenia in Community-Dwelling Older Adults at Risk of Malnutrition. *BMC Geriatr.* **2022**, *22*, 997. <https://doi.org/10.1186/s12877-022-03704-1>.
5. Picca, A.; Calvani, R.; Coelho-Júnior, H.J.; Landi, F.; Marzetti, E. Anorexia of Aging: Metabolic Changes and Biomarker Discovery. *Clin. Interv. Aging* **2022**, *17*, 1761–1767. <https://doi.org/10.2147/CIA.S325008>.
6. Picca, A.; Coelho-Junior, H.J.; Calvani, R.; Marzetti, E.; Vetrano, D.L. Biomarkers Shared by Frailty and Sarcopenia in Older Adults: A Systematic Review and Meta-Analysis. *Ageing Res. Rev.* **2022**, *73*, 101530. <https://doi.org/10.1016/j.arr.2021.101530>.
7. Han, K.; Park, Y.-M.; Kwon, H.-S.; Ko, S.-H.; Lee, S.-H.; Yim, H.W.; Lee, W.-C.; Park, Y.G.; Kim, M.K.; Park, Y.-M. Sarcopenia as a Determinant of Blood Pressure in Older Koreans: Findings from the Korea National Health and Nutrition Examination Surveys (KNHANES) 2008–2010. *PLoS ONE* **2014**, *9*, e86902. <https://doi.org/10.1371/journal.pone.0086902>.
8. Garber, C.E.; Blissmer, B.; Deschenes, M.R.; Franklin, B.A.; Lamonte, M.J.; Lee, I.-M.; Nieman, D.C.; Swain, D.P.; American College of Sports Medicine. American College of Sports Medicine Position Stand. Quantity and Quality of Exercise for Developing and Maintaining Cardiorespiratory, Musculoskeletal, and Neuromotor Fitness in Apparently Healthy Adults: Guidance for Prescribing Exercise. *Med. Sci. Sports Exerc.* **2011**, *43*, 1334–1359. <https://doi.org/10.1249/MSS.0b013e318213febf>.
9. Makanae, Y.; Fujita, S. Role of Exercise and Nutrition in the Prevention of Sarcopenia. *J. Nutr. Sci. Vitam.* **2015**, *61*, S125–S127. <https://doi.org/10.3177/JNSV.61.S125>.
10. Cruz-Jentoft, A.J.; Landi, F.; Schneider, S.M.; Zúñiga, C.; Arai, H.; Boirie, Y.; Chen, L.K.; Fielding, R.A.; Martin, F.C.; Michel, J.; et al. Prevalence of and Interventions for Sarcopenia in Ageing Adults: A Systematic Review. Report of the International Sarcopenia Initiative (EWGSOP and IWGS). *Age Ageing* **2014**, *43*, 48–759. <https://doi.org/10.1093/AGEING/AFU115>.
11. Marzetti, E.; Calvani, R.; Tosato, M.; Cesari, M.; Di Bari, M.; Cherubini, A.; Broccatelli, M.; Saveria, G.; D’Elia, M.; Pahor, M.; et al. Physical Activity and Exercise as Countermeasures to Physical Frailty and Sarcopenia. *Aging Clin. Exp. Res.* **2017**, *29*, 35–42. <https://doi.org/10.1007/s40520-016-0705-4>.
12. Lopez, P.; Pinto, R.S.; Radaelli, R.; Rech, A.; Grazioli, R.; Izquierdo, M.; Cadore, E.L. Benefits of Resistance Training in Physically Frail Elderly: A Systematic Review. *Aging Clin. Exp. Res.* **2018**, *30*, 889–899. <https://doi.org/10.1007/s40520-017-0863-z>.
13. Coelho-Júnior, H.J.; Uchida, M.C. Effects of Low-Speed and High-Speed Resistance Training Programs on Frailty Status, Physical Performance, Cognitive Function, and Blood Pressure in Pre frail and Frail Older Adults. *Front. Med.* **2021**, *8*, 702436. <https://doi.org/10.3389/fmed.2021.702436>.
14. Shen, Y.; Shi, Q.; Nong, K.; Li, S.; Yue, J.; Huang, J.; Dong, B.; Beauchamp, M.; Hao, Q. Exercise for Sarcopenia in Older People: A Systematic Review and Network Meta-Analysis. *J. Cachexia Sarcopenia Muscle* **2023**, *14*, 1199–1211. <https://doi.org/10.1002/JCSM.13225>.
15. Coelho-Júnior, H.J.; Calvani, R.; Picca, A.; Tosato, M.; Landi, F.; Marzetti, E. Engagement in Aerobic Exercise Is Associated with a Reduced Prevalence of Sarcopenia and Severe Sarcopenia in Italian Older Adults. *J. Pers. Med.* **2023**, *13*, 655. <https://doi.org/10.3390/JPM13040655>.

16. Grgic, J.; Mcllvenna, L.C.; Fyfe, J.J.; Sabol, F.; Bishop, D.J.; Schoenfeld, B.J.; Pedisic, Z. Does Aerobic Training Promote the Same Skeletal Muscle Hypertrophy as Resistance Training? A Systematic Review and Meta-Analysis. *Sports Med.* **2019**, *49*, 233–254. <https://doi.org/10.1007/s40279-018-1008-z>.
17. Sharman, J.E.; Smart, N.A.; Coombes, J.S.; Stowasser, M. Exercise and Sport Science Australia Position Stand Update on Exercise and Hypertension. *J. Hum. Hypertens.* **2019**, *33*, 837–843. <https://doi.org/10.1038/S41371-019-0266-Z>.
18. Cornelissen, V.A.; Smart, N.A. Exercise Training for Blood Pressure: A Systematic Review and Meta-Analysis. *J. Am. Heart Assoc.* **2013**, *2*, e004473. <https://doi.org/10.1161/JAHA.112.004473>.
19. Earnest, C.P.; Johannsen, N.M.; Swift, D.L.; Gillison, F.B.; Mikus, C.R.; Lucia, A.; Kramer, K.; Lavie, C.J.; Church, T.S. Aerobic and Strength Training in Concomitant Metabolic Syndrome and Type 2 Diabetes. *Med. Sci. Sports Exerc.* **2014**, *46*, 1293–1301. <https://doi.org/10.1249/MSS.0000000000000242>.
20. Cattadori, G.; Segurini, C.; Picozzi, A.; Padeletti, L.; Anzà, C. Exercise and Heart Failure: An Update. *ESC Heart Fail.* **2018**, *5*, 222–232. <https://doi.org/10.1002/EHF2.12225>.
21. Sousa, N.; Mendes, R.; Abrantes, C.; Sampaio, J.; Oliveira, J. Long-Term Effects of Aerobic Training versus Combined Aerobic and Resistance Training in Modifying Cardiovascular Disease Risk Factors in Healthy Elderly Men. *Geriatr. Gerontol. Int.* **2013**, *13*, 928–935. <https://doi.org/10.1111/GGI.12033>.
22. Bach-Faig, A.; Berry, E.M.; Lairon, D.; Reguant, J.; Trichopoulou, A.; Dernini, S.; Medina, F.X.; Battino, M.; Belahsen, R.; Miranda, G.; et al. Mediterranean Diet Pyramid Today. Science and Cultural Updates. *Public Health Nutr.* **2011**, *14*, 2274–2284. <https://doi.org/10.1017/S1368980011002515>.
23. Diolintzi, A.; Panagiotakos, D.B.; Sidossis, L.S. From Mediterranean Diet to Mediterranean Lifestyle: A Narrative Review. *Public Health Nutr.* **2019**, *22*, 2703–2713. <https://doi.org/10.1017/S1368980019000612>.
24. Sofi, F.; Cesari, F.; Abbate, R.; Gensini, G.F.; Casini, A. Adherence to Mediterranean Diet and Health Status: Meta-Analysis. *BMJ* **2008**, *337*, 673–675. <https://doi.org/10.1136/bmj.a1344>.
25. Sofi, F.; Macchi, C.; Abbate, R.; Gensini, G.F.; Casini, A. Mediterranean Diet and Health Status: An Updated Meta-Analysis and a Proposal for a Literature-Based Adherence Score. *Public Health Nutr.* **2014**, *17*, 2769–2782. <https://doi.org/10.1017/S1368980013003169>.
26. Coelho-Júnior, H.J.; Trichopoulou, A.; Panza, F. Cross-Sectional and Longitudinal Associations between Adherence to Mediterranean Diet with Physical Performance and Cognitive Function in Older Adults: A Systematic Review and Meta-Analysis. *Ageing Res. Rev.* **2021**, *70*, 101395. <https://doi.org/10.1016/j.arr.2021.101395>.
27. Cacciatore, S.; Calvani, R.; Marzetti, E.; Picca, A.; Coelho-Júnior, H.J.; Martone, A.M.; Massaro, C.; Tosato, M.; Landi, F. Low Adherence to Mediterranean Diet Is Associated with Probable Sarcopenia in Community-Dwelling Older Adults: Results from the Longevity Check-Up (Lookup) 7+ Project. *Nutrients* **2023**, *15*, 1026. <https://doi.org/10.3390/NU15041026>.
28. Marzetti, E.; Calvani, R.; Picca, A.; Sisto, A.; Tosato, M.; Martone, A.M.; Ortolani, E.; Salini, S.; Pafundi, T.; Santoliquido, A.; et al. Prevalence of Dyslipidaemia and Awareness of Blood Cholesterol Levels among Community-Living People: Results from the Longevity Check-up 7+ (Lookup 7+) Cross-Sectional Survey. *BMJ Open* **2018**, *8*, e021627. <https://doi.org/10.1136/BMJOPEN-2018-021627>.
29. Landi, F.; Calvani, R.; Picca, A.; Tosato, M.; D'Angelo, E.; Martone, A.M.; Serafini, E.; Ortolani, E.; Saveria, G.; Salini, S.; et al. Relationship between Cardiovascular Health Metrics and Physical Performance in Community-Living People: Results from the Longevity Check-up (Lookup) 7+ Project. *Sci. Rep.* **2018**, *8*, 16353. <https://doi.org/10.1038/s41598-018-34746-4>.
30. Landi, F.; Calvani, R.; Picca, A.; Tosato, M.; Martone, A.M.; Ortolani, E.; Salini, S.; Pafundi, T.; Saveria, G.; Pantanelli, C.; et al. Cardiovascular Health Metrics, Muscle Mass and Function among Italian Community-Dwellers: The Lookup 7+ Project. *Eur. J. Public Health* **2018**, *28*, 766–772. <https://doi.org/10.1093/eurpub/cky034>.
31. Landi, F.; Calvani, R.; Picca, A.; Tosato, M.; Martone, A.M.; Ortolani, E.; Sisto, A.; D'angelo, E.; Serafini, E.; Desideri, G.; et al. Body Mass Index Is Strongly Associated with Hypertension: Results from the Longevity Check-up 7+ Study. *Nutrients* **2018**, *10*, 1976. <https://doi.org/10.3390/NU10121976>.
32. Landi, F.; Calvani, R.; Picca, A.; Tosato, M.; Martone, A.M.; D'Angelo, E.; Serafini, E.; Bernabei, R.; Marzetti, E. Impact of Habitual Physical Activity and Type of Exercise on Physical Performance across Ages in Community-Living People. *PLoS ONE* **2018**, *13*, e0191820. <https://doi.org/10.1371/JOURNAL.PONE.0191820>.
33. von Elm, E.; Altman, D.G.; Egger, M.; Pocock, S.J.; Gøtzsche, P.C.; Vandenbroucke, J.P.; STROBE Initiative The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for Reporting Observational Studies. *Int. J. Surg.* **2014**, *12*, 1495–1499. <https://doi.org/10.1016/j.ijsu.2014.07.013>.
34. Klitgaard, H.; Mantoni, M.; Schiaffino, S.; Ausoni, S.; Gorza, L.; Laurent-Winter, C.; Schnohr, P.; Saltin, B. Function, Morphology and Protein Expression of Ageing Skeletal Muscle: A Cross-Sectional Study of Elderly Men with Different Training Backgrounds. *Acta Physiol. Scand* **1990**, *140*, 41–54. <https://doi.org/10.1111/J.1748-1716.1990.TB08974.X>.
35. Santos, L.P.; Gonzalez, M.C.; Orlandi, S.P.; Bielemann, R.M.; Barbosa-Silva, T.G.; Heymsfield, S.B.; COCONUT Study Group. New Prediction Equations to Estimate Appendicular Skeletal Muscle Mass Using Calf Circumference: Results From NHANES 1999–2006. *JPEN J. Parenter. Enteral. Nutr.* **2019**, *43*, 998–1007. <https://doi.org/10.1002/jpen.1605>.
36. Markofski, M.M.; Jennings, K.; Timmerman, K.L.; Dickinson, J.M.; Fry, C.S.; Borack, M.S.; Reidy, P.T.; Deer, R.R.; Randolph, A.; Rasmussen, B.B.; et al. Effect of Aerobic Exercise Training and Essential Amino Acid Supplementation for 24 Weeks on Physical Function, Body Composition, and Muscle Metabolism in Healthy, Independent Older Adults: A Randomized Clinical Trial. *J. Gerontol. A Biol. Sci. Med. Sci.* **2019**, *74*, 1598–1604. <https://doi.org/10.1093/GERONA/GLY109>.

37. Li, F.; Hsueh, Y.T.; Hsu, Y.J.; Lee, M.C.; Chang, C.H.; Ho, C.S.; Huang, C.C. Effects of Isolated Soy Protein Supplementation Combined with Aerobic Exercise Training on Improving Body Composition, Anthropometric Characteristics and Cardiopulmonary Endurance in Women: A Pilot Study. *Int. J. Env. Res. Public Health* **2021**, *18*, 11798. <https://doi.org/10.3390/IJERPH182211798>.
38. Fairbairn, P.; Tsofliou, F.; Johnson, A.; Dyall, S.C. Effects of a High-DHA Multi-Nutrient Supplement and Exercise on Mobility and Cognition in Older Women (MOBILE): A Randomised Semi-Blinded Placebo-Controlled Study. *Br. J. Nutr.* **2020**, *124*, 146–155. <https://doi.org/10.1017/S0007114520000719>.
39. Lin, Y.N.; Tseng, T.T.; Knuijman, P.; Chan, W.P.; Wu, S.H.; Tsai, C.L.; Hsu, C.Y. Protein Supplementation Increases Adaptations to Endurance Training: A Systematic Review and Meta-Analysis. *Clin. Nutr.* **2021**, *40*, 3123–3132. <https://doi.org/10.1016/J.CLNU.2020.12.012>.
40. Pineda-Juárez, J.A.; Lozada-Mellado, M.; Hinojosa-Azaola, A.; García-Morales, J.M.; Ogata-Medel, M.; Llorente, L.; Alcocer-Varela, J.; Orea-Tejeda, A.; Martín-Nares, E.; Castillo-Martínez, L. Changes in Hand Grip Strength and Body Weight after a Dynamic Exercise Program and Mediterranean Diet in Women with Rheumatoid Arthritis: A Randomized Clinical Trial. *Physiother. Theory Prac.* **2022**, *38*, 504–512. <https://doi.org/10.1080/09593985.2020.1777605>.
41. Trichopoulou, A.; Vasilopoulou, E. Mediterranean Diet and Longevity. *Br. J. Nutr.* **2000**, *84* (Suppl. 2), 205–209. <https://doi.org/10.1079/096582197388554>.
42. Phillips, S.M. The Science of Muscle Hypertrophy: Making Dietary Protein Count. *Proc. Nutr. Soc.* **2011**, *70*, 100–103. <https://doi.org/10.1017/S002966511000399X>.
43. Mitchell, W.K.; Wilkinson, D.J.; Phillips, B.E.; Lund, J.N.; Smith, K.; Atherton, P.J. Human Skeletal Muscle Protein Metabolism Responses to Amino Acid Nutrition. *Adv. Nutr.* **2016**, *7*, 828S–838S. <https://doi.org/10.3945/an.115.011650>.
44. Coelho-Junior, H.J.; Marzetti, E.; Picca, A.; Cesari, M.; Uchida, M.C.; Calvani, R. Protein Intake and Frailty: A Matter of Quantity, Quality, and Timing. *Nutrients* **2020**, *12*, 2915. <https://doi.org/10.3390/NU12102915>.
45. Smith, K.; Reynolds, N.; Downie, S.; Patel, A.; Rennie, M.J. Effects of Flooding Amino Acids on Incorporation of Labeled Amino Acids into Human Muscle Protein. *Am. J. Physiol. Endocrinol. Metab.* **1998**, *275*, E73–E78. <https://doi.org/10.1152/ajpendo.1998.275.1.e73>.
46. Borack, M.S.; Volpi, E. Efficacy and Safety of Leucine Supplementation in the Elderly. *J. Nutr.* **2016**, *146*, 2625S–2629S. <https://doi.org/10.3945/JN.116.230771>.
47. Lee, S.Y.; Lee, H.J.; Lim, J.-Y. Effects of Leucine-Rich Protein Supplements in Older Adults with Sarcopenia: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Arch. Gerontol. Geriatr.* **2022**, *102*, 104758. <https://doi.org/10.1016/j.archger.2022.104758>.
48. Granic, A.; Dismore, L.; Hurst, C.; Robinson, S.M.; Sayer, A.A. Myoprotective Whole Foods, Muscle Health and Sarcopenia: A Systematic Review of Observational and Intervention Studies in Older Adults. *Nutrients* **2020**, *12*, 2257. <https://doi.org/10.3390/NU12082257>.
49. Gorissen, S.H.M.; Rémond, D.; van Loon, L.J.C. The Muscle Protein Synthetic Response to Food Ingestion. *Meat Sci.* **2015**, *109*, 96–100. <https://doi.org/10.1016/J.MEATSCI.2015.05.009>.
50. Symons, T.B.; Sheffield-Moore, M.; Wolfe, R.R.; Paddon-Jones, D. A Moderate Serving of High-Quality Protein Maximally Stimulates Skeletal Muscle Protein Synthesis in Young and Elderly Subjects. *J. Am. Diet. Assoc.* **2009**, *109*, 1582–1586. <https://doi.org/10.1016/J.JADA.2009.06.369>.
51. Morris, M.S.; Jacques, P.F. Total Protein, Animal Protein and Physical Activity in Relation to Muscle Mass in Middle-Aged and Older Americans. *Br. J. Nutr.* **2013**, *109*, 1294–1303. <https://doi.org/10.1017/S0007114512003133>.
52. Robinson, S.M.; Jameson, K.A.; Batelaan, S.F.; Martin, H.J.; Syddall, H.E.; Dennison, E.M.; Cooper, C.; Sayer, A.A. Hertfordshire Cohort Study Group. Diet and Its Relationship With Grip Strength in Community-Dwelling Older Men and Women: The Hertfordshire Cohort Study. *J. Am. Geriatr. Soc.* **2008**, *56*, 84–90. <https://doi.org/10.1111/j.1532-5415.2007.01478.x>.
53. Symons, T.B.; Schutzler, S.E.; Cocke, T.L.; Chinkes, D.L.; Wolfe, R.R.; Paddon-Jones, D. Aging Does Not Impair the Anabolic Response to a Protein-Rich Meal. *Am. J. Clin. Nutr.* **2007**, *86*, 451–456. <https://doi.org/10.1093/AJCN/86.2.451>.
54. Asp, M.L.; Richardson, J.R.; Collene, A.L.; Droll, K.R.; Belury, M.A. Dietary Protein and Beef Consumption Predict for Markers of Muscle Mass and Nutrition Status in Older Adults. *J. Nutr. Health Aging* **2012**, *16*, 784–790. <https://doi.org/10.1007/s12603-012-0064-6>.
55. Valenzuela, P.L.; Mata, F.; Morales, J.S.; Castillo-García, A.; Lucia, A. Does Beef Protein Supplementation Improve Body Composition and Exercise Performance? A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Nutrients* **2019**, *11*, 1429. <https://doi.org/10.3390/NU11061429>.
56. Marcos-Pardo, P.J.; González-Gálvez, N.; López-Vivancos, A.; Espeso-García, A.; Martínez-Aranda, L.M.; Gea-García, G.M.; Orquín-Castrillón, F.J.; Carbonell-Baeza, A.; Jiménez-García, J.D.; Velázquez-Díaz, D.; et al. Sarcopenia, Diet, Physical Activity and Obesity in European Middle-Aged and Older Adults: The LifeAge Study. *Nutrients* **2020**, *13*, 8. <https://doi.org/10.3390/NU13010008>.
57. Tipton, K.D.; Elliott, T.A.; Cree, M.G.; Wolf, S.E.; Sanford, A.P.; Wolfe, R.R. Ingestion of Casein and Whey Proteins Result in Muscle Anabolism after Resistance Exercise. *Med. Sci. Sports Exerc.* **2004**, *36*, 2073–2081. <https://doi.org/10.1249/01.MSS.0000147582.99810.C5>.

58. Tang, J.E.; Moore, D.R.; Kujbida, G.W.; Tarnopolsky, M.A.; Phillips, S.M. Ingestion of Whey Hydrolysate, Casein, or Soy Protein Isolate: Effects on Mixed Muscle Protein Synthesis at Rest and Following Resistance Exercise in Young Men. *J. Appl. Physiol.* **2009**, *107*, 987–992. <https://doi.org/10.1152/JAPPLPHYSIOL.00076.2009>.
59. Sepandi, M.; Samadi, M.; Shirvani, H.; Alimohamadi, Y.; Taghdir, M.; Goudarzi, F.; Akbarzadeh, I. Effect of Whey Protein Supplementation on Weight and Body Composition Indicators: A Meta-Analysis of Randomized Clinical Trials. *Clin. Nutr. ESPEN* **2022**, *50*, 74–83. <https://doi.org/10.1016/J.CLNESP.2022.05.020>.
60. Pennings, B.; Boirie, Y.; Senden, J.M.G.; Gijzen, A.P.; Kuipers, H.; Van Loon, L.J.C. Whey Protein Stimulates Postprandial Muscle Protein Accretion More Effectively than Do Casein and Casein Hydrolysate in Older Men. *Am. J. Clin. Nutr.* **2011**, *93*, 997–1005. <https://doi.org/10.3945/ajcn.110.008102>.
61. Calvani, R.; Picca, A.; Coelho-Júnior, H.J.; Tosato, M.; Marzetti, E.; Landi, F. Diet for the Prevention and Management of Sarcopenia. *Metabolism* **2023**, *146*, 155637. <https://doi.org/10.1016/j.metabol.2023.155637>.
62. Morisasa, M.; Goto-Inoue, N.; Sato, T.; Machida, K.; Fujitani, M.; Kishida, T.; Uchida, K.; Mori, T. Investigation of the Lipid Changes That Occur in Hypertrophic Muscle Due to Fish Protein-Feeding Using Mass Spectrometry Imaging. *J. Oleo Sci.* **2019**, *68*, 141–148. <https://doi.org/10.5650/JOS.ESS18193>.
63. Mizushige, T.; Kawabata, F.; Uozumi, K.; Tsuji, T.; Kishida, T.; Ebihara, K. Fast-Twitch Muscle Hypertrophy Partly Induces Lipid Accumulation Inhibition with Alaska Pollack Protein Intake in Rats. *Biomed. Res.* **2010**, *31*, 347–352. <https://doi.org/10.2220/BIMEDRES.31.347>.
64. Struijk, E.A.; Banegas, J.R.; Rodríguez-Artalejo, F.; Lopez-Garcia, E. Consumption of Meat in Relation to Physical Functioning in the Seniors-ENRICA Cohort. *BMC Med.* **2018**, *16*, 50. <https://doi.org/10.1186/s12916-018-1036-4>.
65. Yokoyama, Y.; Kitamura, A.; Seino, S.; Kim, H.; Obuchi, S.; Kawai, H.; Hirano, H.; Watanabe, Y.; Motokawa, K.; Narita, M.; et al. Association of Nutrient-Derived Dietary Patterns with Sarcopenia and Its Components in Community-Dwelling Older Japanese: A Cross-Sectional Study. *Nutr. J.* **2021**, *20*, 7. <https://doi.org/10.1186/s12937-021-00665-w>.
66. Coelho-Junior, H.J.; Calvani, R.; Gonçalves, I.O.; Rodrigues, B.; Picca, A.; Landi, F.; Bernabei, R.; Uchida, M.C.; Marzetti, E. High Relative Consumption of Vegetable Protein Is Associated with Faster Walking Speed in Well-Functioning Older Adults. *Aging Clin. Exp. Res.* **2019**, *31*, 837–844. <https://doi.org/10.1007/s40520-019-01216-4>.
67. Yeung, S.S.Y.; Woo, J. Association of Plant Protein Intake with Change in Physical Performance in Chinese Community-Dwelling Older Adults. *Nutrients* **2022**, *14*, 4534. <https://doi.org/10.3390/nu14214534>.
68. Montiel-Rojas, D.; Nilsson, A.; Santoro, A.; Bazzocchi, A.; de Groot, L.C.P.G.M.; Feskens, E.J.M.; Berendsen, A.A.M.; Madej, D.; Kaluza, J.; Pietruszka, B.; et al. Fighting Sarcopenia in Ageing European Adults: The Importance of the Amount and Source of Dietary Proteins. *Nutrients* **2020**, *12*, 3601. <https://doi.org/10.3390/nu12123601>.
69. Nichele, S.; Phillips, S.M.; Boaventura, B.C.B. Plant-Based Food Patterns to Stimulate Muscle Protein Synthesis and Support Muscle Mass in Humans: A Narrative Review. *Appl. Physiol. Nutr. Metab.* **2022**, *47*, 700–710. <https://doi.org/10.1139/apnm-2021-0806>.
70. Domić, J.; Grootswagers, P.; Van Loon, L.J.C.; De Groot, L.C.P.G.M. Perspective: Vegan Diets for Older Adults? A Perspective On the Potential Impact On Muscle Mass and Strength. *Adv. Nutr.* **2022**, *13*, 712–725. <https://doi.org/10.1093/ADVANCES/NMAC009>.
71. Yang, Y.; Churchward-Venne, T.A.; Burd, N.A.; Breen, L.; Tarnopolsky, M.A.; Phillips, S.M. Myofibrillar Protein Synthesis Following Ingestion of Soy Protein Isolate at Rest and after Resistance Exercise in Elderly Men. *Nutr. Metab.* **2012**, *9*, 57. <https://doi.org/10.1186/1743-7075-9-57>.
72. Phillips, S.M. Dietary Protein Requirements and Adaptive Advantages in Athletes. *Br. J. Nutr.* **2012**, *108* (Suppl. 2), S158–S167. <https://doi.org/10.1017/S0007114512002516>.
73. Camera, D.M. Evaluating the Effects of Increased Protein Intake on Muscle Strength, Hypertrophy and Power Adaptations with Concurrent Training: A Narrative Review. *Sports Med.* **2022**, *52*, 441–461. <https://doi.org/10.1007/S40279-021-01585-9>.
74. Trichopoulou, A.; Martínez-González, M.A.; Tong, T.Y.N.; Forouhi, N.G.; Khandelwal, S.; Prabhakaran, D.; Mozaffarian, D.; de Lorgeril, M. Definitions and Potential Health Benefits of the Mediterranean Diet: Views from Experts around the World. *BMC Med.* **2014**, *12*, 112. <https://doi.org/10.1186/1741-7015-12-112>.
75. Echeverría, G.; Tiboni, O.; Berkowitz, L.; Pinto, V.; Samith, B.; von Schultendorff, A.; Pedrals, N.; Bitran, M.; Ruini, C.; Ryff, C.D.; et al. Mediterranean Lifestyle to Promote Physical, Mental, and Environmental Health: The Case of Chile. *Int. J. Env. Res. Public Health* **2020**, *17* (Suppl. 3), 200. <https://doi.org/10.3390/IJERPH17228482>.
76. Hershey, M.S.; Martínez-González, M.Á.; Álvarez-Álvarez, I.; Martínez Hernández, J.A.; Ruiz-Canela, M. The Mediterranean Diet and Physical Activity: Better Together than Apart for the Prevention of Premature Mortality. *Br. J. Nutr.* **2022**, *128*, 1413–1424. <https://doi.org/10.1017/S0007114521002877>.
77. Van Elswyk, M.E.; Teo, L.; Lau, C.S.; Shanahan, C.J. Dietary Patterns and the Risk of Sarcopenia: A Systematic Review and Meta-Analysis. *Curr. Dev. Nutr.* **2022**, *6*, nza001. <https://doi.org/10.1093/CDN/NZAC001>.
78. Kennedy, E.T.; Ohls, J.; Carlson, S.; Fleming, K. The Healthy Eating Index: Design and Applications. *J. Am. Diet. Assoc.* **1995**, *95*, 1103–1108. [https://doi.org/10.1016/S0002-8223\(95\)00300-2](https://doi.org/10.1016/S0002-8223(95)00300-2).
79. Parsons, T.J.; Papachristou, E.; Atkins, J.L.; Papacosta, O.; Ash, S.; Lennon, L.T.; Whincup, P.H.; Ramsay, S.E.; Wannamethee, S.G. Healthier Diet Quality and Dietary Patterns Are Associated with Lower Risk of Mobility Limitation in Older Men. *Eur. J. Nutr.* **2019**, *58*, 2335–2343. <https://doi.org/10.1007/S00394-018-1786-Y>.

80. Chan, R.; Leung, J.; Woo, J. A Prospective Cohort Study to Examine the Association Between Dietary Patterns and Sarcopenia in Chinese Community-Dwelling Older People in Hong Kong. *J. Am. Med. Dir. Assoc.* **2016**, *17*, 336–342. <https://doi.org/10.1016/J.JAMDA.2015.12.004>.
81. Perälä, M.M.; Von Bonsdorff, M.B.; Männistö, S.; Salonen, M.K.; Simonen, M.; Kanerva, N.; Rantanen, T.; Pohjolainen, P.; Eriksson, J.G. The Healthy Nordic Diet Predicts Muscle Strength 10 Years Later in Old Women, but Not Old Men. *Age Ageing* **2017**, *46*, 588–594. <https://doi.org/10.1093/AGEING/AFX034>.
82. Coelho-Júnior, H.J.; Calvani, R.; Picca, A.; Marzetti, E. Are Sit-to-Stand and Isometric Handgrip Tests Comparable Assessment Tools to Identify Dynapenia in Sarcopenic People? *Arch. Gerontol. Geriatr.* **2023**, *114*, 105059. <https://doi.org/10.1016/J.ARCHGER.2023.105059>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.