

**Supplementary Table S1:** Spearman correlation coefficients between MNA score, MNA sub-scores and body composition/function patterns in patients with remission (subsample of the present study).

	Pattern 1 (MAC, FFMI, Max HGS)	Pattern 2 (Gait speed)	Pattern 3 (CC, MAC, age)	Pattern 4 (Chair-standing test, OLST)	Pattern 5 (Alb, TSF)
MNA Total score	0.215  p=0.312	0.110  p=0.608	<b>0.417</b>  <b>p=0.043</b>	0.066  p=0.761	0.178  p=0.405
MNA Food intake	0.306  p=0.147	0.092  p=0.670	0.332  p=0.113	0.366  p=0.078	<b>0.455</b>  <b>p=0.025</b>
MNA weight loss	<b>-0.404</b>  <b>p=0.050</b>	0.291  p=0.168	-0.048  p=0.822	-0.162  p=0.451	0.210  p=0.325
MNA mobility	NA	NA	NA	NA	NA
MNA acute stress	0.044  p=0.840	0.106  p=0.623	0.367  p=0.078	0.180  p=0.399	0.193  p=0.367
MNA neurological problems	0.256  p=0.227	-0.045  p=0.834	0.346  p=0.097	-0.166  p=0.439	-0.166  p=0.439
MNA BMI	<b>0.700</b>  <b>p&lt;0.001</b>	-0.112  p=0.602	0.347  p=0.097	-0.045  p=0.836	0.116  p=0.588

MAC: Mid arm circumference; FFMI: Fat free mass index; HGS: hand-grip strength; CC: Calf circumference; OLST: One-Legged Stance Test; Alb: Albumin; TSF: Triceps Skinfold; MNA: Mini Nutritional Assessment; BMI: Body Mass Index

NA: Not applicable. It is noted that no variability was detected for MNA-mobility and thus no correlation coefficient could be calculated.