

Supplementary Table S1. The proportion of women with UIC \geq or $<$ 100 $\mu\text{g/L}$

	UIC, n (%)		<i>p</i>
	$< 100 \mu\text{g/L}$ (n=37)	$\geq 100 \mu\text{g/L}$ (n=34)	
Age, years	34.3 \pm 3.9	33.8 \pm 4.4	0.70
Educational status			0.64
University or below (n = 58)	31 (53.4)	27 (46.6)	
Master or above (n = 13)	6 (46.2)	7 (53.8)	
Infant birth body weight, g	3062.5 \pm 563.3	3052.8 \pm 309.4	0.65
Parity			0.44
First (n = 41)	23 (56.1)	18 (43.9)	
Second or above (n = 30)	14 (46.7)	16 (53.3)	
Miscarriage history			0.14
Yes (n = 27)	11 (40.7)	16 (59.3)	
No (n = 44)	26 (59.1)	18 (40.9)	
Post-partum nourishment diet			0.18
Yes (n = 13)	9 (69.2)	4 (30.8)	
No (n = 58)	28 (48.3)	30 (51.7)	
Exclusive breastfeeding			0.94
Yes (n = 17)	9 (52.9)	8 (47.1)	
No (n = 54)	28 (51.9)	26 (48.1)	
Salt intake			0.32
Iodized (n = 39)	19 (48.7)	20 (51.3)	
Non-iodized (n = 14)	6 (42.9)	8 (57.1)	
Unknown (n = 18)	12 (66.7)	6 (33.3)	
Seaweed intake			0.02
Yes (n = 42)	17 (40.5)	25 (59.5)	
No (n = 29)	20 (69.0)	9 (31.0)	
Fish intake			0.44
Yes (n = 59)	32 (54.2)	27 (45.8)	
No (n = 12)	5 (41.7)	7 (58.3)	
Seafood intake			0.68
Yes (n = 42)	21 (50.0)	21 (50.0)	
No (n = 29)	16 (55.2)	13 (44.8)	
Dairy intake			0.18
Yes (n = 63)	31 (49.2)	32 (50.8)	
No (n = 8)	6 (75.0)	2 (25.0)	
Multivitamin intake			0.42
Yes (n = 34)	16 (47.1)	18 (52.9)	
No (n = 37)	21 (56.8)	16 (43.2)	

Age and infant birth body weight was presented by mean and standard deviation whereas other variables were presented by number and percentage.