

**Table S1.** Adjusted associations \* between food items with the highest factor loading of each dietary pattern identified during pregnancy and postpartum BMI modification \*\* in women with pregnancy complicated by gestational diabetes mellitus. LINDA-Brasil Study.

	BMI at six months postpartum			BMI at twelve months postpartum		
	RR	[95% CI]	P-value	RR	[95% CI]	P-value
<b>Pregnancy</b>						
<b>Risk pattern</b>						
<b>Fried foods</b>						
<b>Model 1</b>						
0-1/week						
2-4/week	0.93	[0.75 - 1.15]	0.511	0.96	[0.77- 1.19]	0.733
5-7/week	1.61	[1.25 - 2.07]	<0.001	1.47	[1.08- 2.02]	0.015
<b>Model 2</b>						
0-1/week						
2-4/week	1.01	[0.82 - 1.25]	0.899	1.04	[0.84- 1.29]	0.709
5-7/week	1.83	[1.42 - 2.37]	<0.001	1.48	[1.06- 2.07]	0.022
<b>Model 3</b>						
0-1/week						
2-4/week	1.01	[0.82 - 1.25]	0.928	1.04	[0.84- 1.29]	0.694
5-7/week	1.83	[1.43 - 2.34]	<0.001	1.48	[1.06- 2.07]	0.023
<b>Cookies and sweets</b>						
<b>Model 1</b>						
0-1/week						
2-4/week	1.01	[0.84- 1.22]	0.870	0.89	[0.73- 1.09]	0.276
5-7/week	1.03	[0.81- 1.31]	0.804	0.87	[0.66- 1.13]	0.292
<b>Model 2</b>						
0-1/week						
2-4/week	1.08	[0.90 - 1.30]	0.389	0.94	[0.77- 1.15]	0.546
5-7/week	1.16	[0.91- 1.48]	0.236	0.92	[0.70- 1.21]	0.550
<b>Model 3</b>						
0-1/week						
2-4/week	1.07	[0.89- 1.28]	0.477	0.93	[0.76- 1.13]	0.460
5-7/week	1.15	[0.90- 1.47]	0.254	0.92	[0.70- 1.20]	0.536
<b>Sweetened beverages</b>						
<b>Model 1</b>						
0-1/week						
2-4/week	1.05	[0.86- 1.29]	0.618	0.84	[0.66- 1.06]	0.151
5-7/week	1.24	[1.03- 1.49]	0.022	1.24	[1.03- 1.48]	0.019
<b>Model 2</b>						
0-1/week						
2-4/week	1.09	[0.89- 1.33]	0.395	0.85	[0.68- 1.08]	0.194
5-7/week	1.33	[1.10- 1.60]	0.003	1.30	[1.08- 1.55]	0.005
<b>Model 3</b>						
0-1/week						
2-4/week	1.09	[0.89- 1.33]	0.400	0.86	[0.68- 1.09]	0.217
5-7/week	1.31	[1.09- 1.58]	0.004	1.29	[1.08- 1.55]	0.005
<b>Processed meat</b>						

**Model 1**

0-1/week								
2-4/week	1.06	[0.88-	1.27]	0.529	0.97	[0.80-	1.17]	0.746
5-7/week	1.20	[0.97-	1.48]	0.074	1.07	[0.85-	1.33]	0.581

**Model 2**

0-1/week						
2-4/week	1.05	[0.87- 1.26]	0.571	0.94	[0.78- 1.14]	0.554
5-7/week	1.20	[0.97- 1.48]	0.094	1.07	[0.85- 1.34]	0.531

**Model 3**

0-1/week						
2-4/week	1.04	[0.87- 1.25]	0.625	0.94	[0.77- 1.13]	0.506
5-7/week	1.18	[0.95- 1.45]	0.122	1.07	[0.85- 1.33]	0.564

**Healthy pattern**

## Red meat without visible fat

**Model 1**

0-1/week						
2-4/week	0.84	[0.70- 1.00]	0.005	0.91	[0.75- 1.10]	0.346
5-7/week	1.15	[0.94- 1.40]	0.170	1.16	[0.94- 1.43]	0.153

**Model 2**

0-1/week						
2-4/week	0.83	[0.69- 1.00]	0.054	0.89	[0.74- 1.08]	0.240
5-7/week	1.16	[0.95- 1.41]	0.149	1.15	[0.94- 1.42]	0.166

**Model 3**

0-1/week						
2-4/week	0.84	[0.70- 1.00]	0.057	0.89	[0.74- 1.08]	0.247
5-7/week	1.15	[0.95- 1.41]	0.156	1.15	[0.94- 1.42]	0.158

## Raw salad

**Model 1**

0-1/week						
2-4/week	0.89	[0.70- 1.12]	0.322	0.74	[0.57- 0.95]	0.019
5-7/week	0.82	[0.65- 1.03]	0.094	0.87	[0.69- 1.09]	0.239

**Model 2**

0-1/week						
2-4/week	0.85	[0.67- 1.08]	0.179	0.73	[0.57- 0.94]	0.014
5-7/week	0.77	[0.61- 0.97]	0.027	0.84	[0.67- 1.05]	0.126

**Model 3**

0-1/week						
2-4/week	0.85	[0.67- 1.07]	0.176	0.73	[0.56- 0.93]	0.012
5-7/week	0.77	[0.62- 0.97]	0.029	0.84	[0.67- 1.05]	0.129

## Cooked salad

**Model 1**

0-1/week						
2-4/week	1.00	[0.84- 1.20]	0.987	0.92	[0.76- 1.11]	0.387
5-7/week	0.78	[0.61- 0.99]	0.048	0.77	[0.60- 0.98]	0.039

**Model 2**

0-1/week						
2-4/week	0.99	[0.82- 1.18]	0.909	0.91	[0.75- 1.09]	0.315
5-7/week	0.75	[0.59- 0.96]	0.026	0.75	[0.59- 0.95]	0.020

**Model 3**

0-1/week						
2-4/week	0.99	[0.83- 1.19]	0.992	0.91	[0.75- 1.09]	0.298
5-7/week	0.76	[0.59- 0.97]	0.029	0.75	[0.59- 0.95]	0.019

**DASH**

## Low fat dairy

**Model 1**

0-1/week

2-4/week 0.94 [0.70- 1.28] 0.714 1.08 [0.83- 1.40] 0.564

5-7/week 1.12 [0.95- 1.32] 0.158 0.99 [0.84- 1.18] 0.964

**Model 2**

0-1/week

2-4/week 0.93 [0.68- 1.26] 0.642 1.09 [0.83- 1.42] 0.533

5-7/week 1.09 [0.92- 1.28] 0.309 0.97 [0.81- 1.15] 0.722

**Model 3**

0-1/week

2-4/week 0.95 [0.71- 1.30] 0.769 0.98 [0.78- 1.24] 0.915

5-7/week 1.08 [0.92- 1.27] 0.337 0.95 [0.56- 1.57] 0.847

\* Poisson regressions with robust variance adjusted for the outcome; \*\* increase in BMI between 2 months and 6 months to 12 months postpartum and frequency of exposure of dietary items. Model 1: Adjusted for age, skin color, income, city, smoking during pregnancy and number of previous pregnancies. Model 2: Adjusted for model 1 + pre-pregnancy BMI and Gestational weight gain. Model 3: Adjusted for model 2 + Breastfeeding in the follow-up period.

**Table S2.** Adjusted associations \* between food items with the highest factor loading of each dietary pattern identified six months after delivery and postpartum BMI modification \*\* in women with pregnancy complicated by gestational diabetes mellitus. LINDA-Brasil Study.

	BMI at six months postpartum			BMI at twelve months postpartum		
	RR	[95% CI]	P-value	RR	[95% CI]	P-value
Postpartum						
<b>Risk pattern</b>						
<b>Sweetened beverages</b>						
<b>Model 1</b>						
0-1/week						
2-4/week	1.14	[0.93- 1.40]	0.213	1.06	[0.85- 1.32]	0.610
5-7/week	1.09	[0.89- 1.34]	0.386	1.24	[1.01- 1.52]	0.015
<b>Model 2</b>						
0-1/week						
2-4/week	1.18	[0.96- 1.44]	0.109	1.08	[0.86- 1.35]	0.516
5-7/week	1.16	[0.95- 1.42]	0.147	1.27	[1.04- 1.55]	0.017
<b>Model 3</b>						
0-1/week						
2-4/week	1.17	[0.95- 1.43]	0.136	1.06	[0.85- 1.33]	0.583
5-7/week	1.16	[0.95- 1.41]	0.154	1.27	[1.04- 1.55]	0.019
<b>Fried foods</b>						
<b>Model 1</b>						
0-1/week						
2-4/week	1.08	[0.89- 1.31]	0.428	0.92	[0.73- 1.16]	0.487
5-7/week	1.63	[1.15- 2.31]	0.006	1.42	[0.90- 2.24]	0.126
<b>Model 2</b>						
0-1/week						
2-4/week	1.07	[0.88- 1.29]	0.507	0.87	[0.70- 1.09]	0.234
5-7/week	1.85	[1.39- 2.47]	<0.001	1.50	[1.01- 2.23]	0.043
<b>Model 3</b>						
0-1/week						

2-4/week	1.05	[0.86- 1.27]	0.645	0.86	[0.68- 1.08]	0.189
5-7/week	1.87	[1.40; 2.50)	<0.001	1.52	[1.00- 2.29]	0.047
<b>Processed meat</b>						
<b>Model 1</b>						
0-1/week						
2-4/week	1.12	[0.95- 1.32]	0.185	0.99	[0.82- 1.19]	0.912
5-7/week	0.95	[0.72- 1.25]	0.712	1.18	[0.93- 1.50]	0.179
<b>Model 2</b>						
0-1/week						
2-4/week	1.12	[0.95- 1.33]	0.173	0.98	[0.81- 1.17]	0.810
5-7/week	0.93	[0.71- 1.21]	0.580	1.15	[0.91- 1.47]	0.242
<b>Model 3</b>						
0-1/week						
2-4/week	1.11	[0.94- 1.32]	0.195	0.97	[0.81- 1.17]	0.774
5-7/week	0.90	[0.70- 1.18]	0.451	1.13	[0.89- 1.44]	0.307
<b>Cookies and sweets</b>						
<b>Model 1</b>						
0-1/week						
2-4/week	1.11	[0.93- 1.32]	0.227	1.09	[0.92- 1.30]	0.313
5-7/week	1.36	[1.08- 1.70]	0.008	0.98	[0.74- 1.31]	0.910
<b>Model 2</b>						
0-1/week						
2-4/week	1.16	[0.98- 1.37]	0.082	1.18	[0.94- 1.33]	0.209
5-7/week	1.44	[1.14- 1.82]	0.002	0.97	(0.73- 1.29]	0.850
<b>Model 3</b>						
0-1/week						
2-4/week	1.16	[0.98- 1.37]	0.091	1.11	[0.94- 1.33]	0.211
5-7/week	1.44	[1.14- 1.82]	0.002	0.97	[0.73- 1.30]	0.860
<b>Mixed pattern</b>						
<b>Natural fruit juice</b>						
<b>Model 1</b>						
0-1/week						
2-4/week	1.11	[0.91- 1.36]	0.290	0.98	[0.79- 1.22]	0.879
5-7/week	0.92	[0.71- 1.19]	0.547	0.87	[0.66- 1.15]	0.339
<b>Model 2</b>						
0-1/week						
2-4/week	1.09	[0.90- 1.33]	0.375	0.96	[0.78- 1.19]	0.742
5-7/week	0.90	[0.70- 1.17]	0.444	0.87	[0.66- 1.15]	0.343
<b>Model 3</b>						
0-1/week						
2-4/week	1.08	[0.89- 1.32]	0.417	0.96	[0.77- 1.18]	0.687
5-7/week	0.91	[0.70; 1.17)	0.463	0.87	[0.66- 1.15]	0.329
<b>Salty snacks</b>						
<b>Model 1</b>						
0-1/week						
2-4/week	1.00	[0.84- 1.19]	0.946	1.07	[0.90- 1.28]	0.454
5-7/week	1.00	[0.80- 1.26]	0.994	1.20	[0.96- 1.49]	0.098
<b>Model 2</b>						
0-1/week						
2-4/week	0.98	[0.83- 1.17]	0.880	1.06	[0.89- 1.27]	0.506
5-7/week	0.97	[0.77- 1.22]	0.782	1.13	[0.90- 1.40]	0.284

**Model 3**

0-1/week							
2-4/week	0.98	[0.83- 1.16]	0.866	1.07	[0.89- 1.27]	0.464	
5-7/week	0.94	[0.75- 1.19]	0.638	1.11	[0.89- 1.39]	0.348	

## Fruits

**Model 1**

0-1/week							
2-4/week	0.89	[0.73- 1.08]	0.242	0.81	[0.66- 0.98]	0.032	
5-7/week	0.87	[0.72- 1.08]	0.176	0.76	[0.63- 0.93]	0.0078	

**Model 2**

0-1/week							
2-4/week	0.86	[0.71- 1.05]	0.147	0.79	[0.65- 0.96]	0.020	
5-7/week	0.84	[0.69- 1.03]	0.090	0.75	[0.61- 0.91]	0.004	

**Model 3**

0-1/week							
2-4/week	0.87	[0.71- 1.06]	0.166	0.79	[0.65- 0.96]	0.021	
5-7/week	0.86	[0.72- 1.05]	0.128	0.76	[0.62- 0.92]	0.006	

## Chicken

**Model 1**

0-1/week							
2-4/week	0.82	[0.66- 1.03]	0.092	1.01	[0.76- 1.35]	0.912	
5-7/week	0.75	[0.58- 0.98]	0.037	1.00	[0.73- 1.36]	0.996	

**Model 2**

0-1/week							
2-4/week	0.78	[0.62- 0.97]	0.031	0.97	[0.74- 1.28]	0.844	
5-7/week	0.70	[0.54- 0.91]	0.008	0.96	[0.70- 1.30]	0.781	

**Model 3**

0-1/week							
2-4/week	0.79	[0.63- 0.98]	0.037	0.97	[0.74- 1.28]	0.857	
5-7/week	0.70	[0.53- 0.91]	0.008	0.95	[0.70- 1.29]	0.744	

**DASH Type**

## Cooked salad

**Model 1**

0-1/week							
2-4/week	0.94	[0.78- 1.12]	0.473	0.83	[0.69- 0.98]	0.037	
5-7/week	0.97	[0.76- 1.23]	0.784	0.81	[0.63- 1.05]	0.115	

**Model 2**

0-1/week							
2-4/week	0.91	[0.77- 1.09]	0.315	0.82	[0.69- 0.98]	0.030	
5-7/week	0.96	[0.76- 1.22]	0.770	0.83	[0.65- 1.07]	0.149	

**Model 3**

0-1/week							
2-4/week	0.92	[0.77- 1.09]	0.333	0.83	[0.69- 0.98]	0.034	
5-7/week	0.99	[0.78- 1.25]	0.928	0.85	[0.66- 1.09]	0.209	

## Raw salad

**Model 1**

0-1/week							
2-4/week	0.90	[0.74- 1.11]	0.333	0.85	[0.69- 1.03]	0.099	
5-7/week	0.90	[0.73- 1.10]	0.327	0.73	[0.59- 0.89]	0.003	

**Model 2**

0-1/week

2-4/week	0.85	[0.70- 1.05]	0.131	0.80	[0.66- 0.97]	0.026
5-7/week	0.85	[0.69- 1.04]	0.118	0.68	[0.55- 0.84]	<0.001
<b>Model 3</b>						
0-1/week						
2-4/week	0.85	[0.69- 1.05]	0.132	0.80	[0.65- 0.97]	0.024
5-7/week	0.84	[0.68- 1.04]	0.115	0.68	[0.55- 0.84]	<0.001
Low fat dairy						
<b>Model 1</b>						
0-1/week						
2-4/week	1.07	[0.85- 1.34]	0.566	0.99	[0.78- 1.26]	0.960
5-7/week	0.85	[0.46- 1.54]	0.605	0.97	[0.62- 1.54]	0.916
<b>Model 2</b>						
0-1/week						
2-4/week	1.07	[0.85- 1.34]	0.980	0.98	[0.78- 1.24]	0.910
5-7/week	0.78	[0.40- 1.52]	0.476	0.93	[0.56- 1.54]	0.790
<b>Model 3</b>						
0-1/week						
2-4/week	1.07	[0.85- 1.33]	0.555	0.99	[0.78- 1.24]	0.915
5-7/week	0.81	[0.42- 1.58]	0.538	0.95	[0.58- 1.57]	0.847

\* Poisson regressions with robust variance adjusted for the outcome increase in. \*\* BMI between 2 months and 6 months to 12 months post-partum and frequency of exposure of dietary items. Model 1: Adjusted for age, skin color, income, city, smoking during pregnancy and number of previous pregnancies. Model 2: Adjusted for model 1 + pre-pregnancy BMI and Gestational weight gain. Model 3: Adjusted for model 2 + + Breastfeeding in the follow-up period.