


**Figure S1. Analytic Sample Flow Diagram**



<b>Table S1. Baseline Characteristics of Non-Pregnant Partner by Enrollment</b>			
	<b>Pregnancy (n=131)</b>	<b>Postpartum (n=26)</b>	<b>p-value</b>
<b>Age (Mean±SD)</b>	30.8±4.0	33.0±4.5	<b>0.007</b>
<b>Body Mass Index (BMI; kg/m<sup>2</sup>; Mean±SD)</b>	28.5±5.2	28.7±4.5	0.834
Normal Weight, N(%)	35 (26.7)	7 (29.3)	0.954
Overweight, N(%)	54 (40.6)	10 (39.0)	
Obesity, N(%)	42 (32.7)	9 (31.7)	
<b>Race</b>			0.284
White/Caucasian, N(%)	117 (89.0)	22 (85.4)	
Black/African American, N(%)	14 (11.0)	3 (12.2)	
Asian or more than One Race, N(%)	0 (0.0)	1 (2.4)	
Unknown or Not Reported, N(%)	0 (0.0)	0 (0.0)	
<b>Highest Education Completed</b>			0.178
<College Graduate, N(%)	56 (42.7)	15 (59.0)	
≥College Graduate, N(%)	72 (55.2)	11 (41.0)	
Unknown/Not Reported, N(%)	3 (2.1)	0 (0.0)	
<b>Individual Income</b>			0.229
<\$40,000 USD, N(%)	42 (31.7)	15 (59.0)	
≥\$40,000 USD, N(%)	83 (63.4)	11 (41.0)	
Unknown/Not Reported, N(%)	6 (4.9)	0 (0.0)	
<b>The Eating Inventory</b>			
Cognitive Restraint, M±SD	6.6±4.8	7.0±4.5	0.625
Low, N(%)	102 (78.2)	22 (82.9)	0.528
High, N(%)	29 (21.8)	4 (17.1)	
Dietary Disinhibition, M±SD	5.4±3.1	5.0±4.0	0.534
Low, N(%)	113 (86.1)	22 (82.9)	0.625
High, N(%)	18 (13.9)	4 (17.1)	
Perceived Hunger, M±SD	4.8±3.2	4.5±2.9	0.677
Low, N(%)	104 (79.2)	22 (82.9)	0.614
High, N(%)	27 (20.8)	4 (17.1)	
Reported as N (%) if not otherwise noted.			

<b>Table S2. Baseline Characteristics of Pregnant Persons and Non-Pregnant Partners</b>			
	<b>Pregnant Person (n=137)</b>	<b>Non-Pregnant Partner (n=137)</b>	<b>p-value</b>
<b>Age (Mean±SD)</b>	30.0±3.7	31.4±4.3	0.07
<b>Body Mass Index (BMI; kg/m<sup>2</sup>; Mean±SD)</b>	26.1±4.3	28.5±5.3	<0.0001
Normal Weight, N(%)	62 (45.3)	39 (28.5)	
Overweight, N(%)	44 (32.1)	55 (40.1)	
Obesity, N(%)	31 (22.6)	43 (31.4)	
<b>Marital Status*</b>			
Married or Cohabiting, N(%)	125 (91.2)	-	
Single, Divorced, or Unknown/Not Reported, N(%)	12 (8.8)	-	
<b>Race</b>			0.70
White/Caucasian, N(%)	112 (81.7)	110 (80.3)	
Black/African American, N(%)	12 (8.8)	16 (11.7)	
Asian or more than One Race, N(%)	2 (1.5)	0 (0.0)	
Unknown or Not Reported, N(%)	11 (8.0)	11 (8.0)	
<b>Highest Education Completed</b>			<0.01
<College Graduate, N(%)	46 (33.6)	54 (39.4)	
≥College Graduate, N(%)	81 (59.1)	64 (46.7)	
Unknown/Not Reported, N(%)	10 (7.3)	19 (13.9)	
<b>Individual Income</b>			<0.01
<\$40,000 USD, N(%)	67 (48.9)	42 (30.7)	
≥\$40,000 USD, N(%)	40 (29.2)	80 (58.4)	
Unknown/Not Reported, N(%)	30 (21.9)	15 (10.9)	
Reported as N (%) if not otherwise noted. *Marital Status only reported by Pregnant Persons			

<b>Table S3. Gestational Weight Gain and The Eating Inventory Overall and by Body Mass Index</b>				
<b>Pregnant Person</b>	Overall (137)	Normal Weight (62)	Overweight (44)	Obese (31)
<b>Gestational Weight Gain (Total; kg)</b>	11.5±4.4	13.1±3.0*	11.4±4.9*	8.6±4.6
Inadequate, N(%)	51 (37.2)	38 (61.3) <sup>‡</sup>	7 (15.9)	6 (19.4)
Recommended, N(%)	62 (45.3)	24 (38.7) <sup>‡</sup>	23 (52.3)	15 (48.4)
Excess, N(%)	24 (17.5)	0 (0.0) <sup>‡</sup>	14 (31.8)	10 (32.3)
<b>The Eating Inventory</b>				
Cognitive Restraint, M±SD	9.9±4.7 <sup>‡</sup>	9.0±4.7 <sup>‡</sup>	10.5±4.8	10.6±4.3
Low, N(%)	73 (53.3) <sup>‡</sup>	36 (58.1) <sup>‡</sup>	20 (45.5)	17 (54.8)
High, N(%)	64 (46.7) <sup>‡</sup>	26 (41.9) <sup>‡</sup>	24 (54.5)	14 (45.2)
Dietary Disinhibition, M±SD	4.6±3.1	3.0±2.1 <sup>‡</sup>	5.5±3.0	6.3±3.7
Low, N(%)	122 (89.1)	61 (98.4) <sup>‡</sup>	37 (84.1)	24 (77.4)
High, N(%)	15 (10.9)	1 (1.6) <sup>‡</sup>	15 (15.9)	7 (22.6)
Perceived Hunger, M±SD	4.6±2.6	4.0±2.3 <sup>‡</sup>	5.3±2.7	4.9±2.8
Low, N(%)	119 (86.9)	58 (93.6) <sup>‡</sup>	34 (77.3)	27 (87.1)
High, N(%)	18 (13.1)	4 (6.5) <sup>‡</sup>	10 (22.7)	7 (12.9)
<b>Non-Pregnant Partner</b>	Overall (137)	Normal Weight (62)	Overweight (44)	Obese (31)
<b>The Eating Inventory</b>				
Cognitive Restraint, M±SD	6.7±4.6 <sup>‡</sup>	6.8±4.8	6.9±4.3	6.2±4.9
Low, N(%)	110 (80.3) <sup>‡</sup>	51 (82.3)	35 (79.5)	24 (77.4)
High, N(%)	27 (19.7) <sup>‡</sup>	11 (17.7)	9 (20.5)	7 (22.6)
Dietary Disinhibition, M±SD	5.2±3.4	5.1±3.2	4.8±3.2	5.9±3.9
Low, N(%)	120 (87.6)	56 (90.3)	39 (88.6)	25 (80.7)
High, N(%)	17 (12.4)	6 (9.7)	5 (11.4)	6 (19.3)
Perceived Hunger, M±SD	4.7±3.2	4.3±3.2	4.7±2.9	5.4±3.5
Low, N(%)	111 (81.0)	53 (85.5)	35 (79.5)	23 (74.2)
High, N(%)	29 (19.0)	9 (14.5)	9 (20.5)	8 (25.8)
<b>Couple (Average)</b>	Overall (137)	Normal Weight (62)	Overweight (44)	Obese (31)
<b>The Eating Inventory</b>				
Cognitive Restraint, M±SD	8.3±3.4	7.9±3.5	8.7±3.3	8.4±3.2
Low, N(%)	100 (73.0)	48 (77.4)	30 (68.2)	22 (71.0)
High, N(%)	37 (27.0)	14 (22.6)	14 (31.8)	9 (29.0)
Dietary Disinhibition, M±SD	4.9±2.4	4.1±2.0	5.2±2.2	6.1±2.7
Low, N(%)	125 (91.2)	60 (96.8)	40 (90.9)	25 (80.6)
High, N(%)	12 (8.8)	2 (3.2)	4 (9.1)	6 (19.4)
Perceived Hunger, M±SD	4.6±2.0	4.1±1.8	5.0±2.0	5.1±2.3
Low, N(%)	130 (94.9)	60 (96.8)	42 (95.5)	28 (90.3)
High, N(%)	7 (5.1)	2 (3.2)	2 (4.5)	3 (9.7)
Parent average was calculated the average between pregnant person and non-pregnant partner Eating Inventory. *p<0.05 for significance compared to those having obesity; <sup>‡</sup> p<0.05 for significance compared to those having overweight or obesity; <sup>‡</sup> p<0.05 for significance compared between maternal and paternal Eating Inventory constructs				