

Figure S1. Schematic diagram depicting the experimental groups involved in the study.




Table S1. Sequences of primer sets used for qRT-PCR experiments.

Gene	Primer sequence (5' → 3')	
	Forward Primer	Reverse Primer
<i>C3</i>	CCAGCTCCCCATTAGCTCTG	CCAGCTCCCCATTAGCTCTG
<i>C5</i>	GAACAAACCTACGTCATTTACAGC	GTCAACAGTGCCGCGTTTT
<i>VEGF</i>	GCACATAGGAGAGATGAGCTTCC	CTCCGCTCTGAACAAGGCT
<i>TNF-α</i>	GCCTCTTCTCATTCTGCTT	CTCCTCCACTTGGTGGTTTG
<i>IL-1β</i>	TCCTTG TGCAAGTGTCTGAAGC	ATGAGTGATACTGCCTGCCTGA
<i>IL-6</i>	TCTGCAAGAGACTTCCATCCAGT	TCTGCAACTGCATCATCGTTGT
<i>IL-8</i>	ACTTTCAGAGACAGCAGAGC	GTGGTCCACTCTCAATCACT
<i>ICAM-1</i>	CGCTGTGCTTTGAGAACTGTG	ATACACGGTGATGGTAGCGGA
<i>iNOS</i>	GGCAAACCCAAGGTCTACGTT	TCGCTCAAGTTCAGCTTGGT
<i>GFAP</i>	CGGAGACGCATCACCTCTG	AGGGAGTGGAGGAGTCATTCTG
<i>CD68</i>	TGTCTGATCTTGCTAGGACCG	GAGAGTAACGGCCTTTTTGTGA
<i>F4/80</i>	TGACTCACCTTGTGGTCCTAA	CTTCCCAGAATCCAGTCTTTCC
<i>Rpl13a</i>	CACTCTGGAGGAGAAACGGAAGG	GCAGGCATGAGGCAAACAGTC