

Review

Microcystin Contamination and Toxicity: Implications for Agriculture and Public Health

Rajesh Melaram *, Amanda R. Newton and Jennifer Chafin

Department of Science, Eastern Florida State College, Melbourne, FL 32935, USA;
newtona@easternflorida.edu (A.R.N.); chafin.jennifer@titans.easternflorida.edu (J.C.)
* Correspondence: melaramr@easternflorida.edu

Abstract: Microcystins are natural hepatotoxic metabolites secreted by cyanobacteria in aquatic ecosystems. When present at elevated concentrations, microcystins can affect water quality aesthetics; contaminate drinking water reservoirs and recreational waters; disrupt normal ecosystem functioning; and cause health hazards to animals, plants, and humans. Animal and human exposures to microcystins generally result from ingesting contaminated drinking water or physically contacting tainted water. Much research has identified a multitude of liver problems from oral exposure to microcystins, varying from hepatocellular damage to primary liver cancer. Provisional guidelines for microcystins in drinking and recreational water have been established to prevent toxic exposures and protect public health. With increasing occurrences of eutrophication in freshwater systems, microcystin contamination in groundwater and surface waters is growing, posing threats to aquatic and terrestrial plants and agricultural soils used for crop production. These microcystins are often transferred to crops via irrigation with local sources of water, such as bloom-forming lakes and ponds. Microcystins can survive in high quantities in various parts of plants (roots, stems, and leaves) due to their high chemical stability and low molecular weight, increasing health risks for consumers of agricultural products. Studies have indicated potential health risks associated with contaminated fruits and vegetables sourced from irrigated water containing microcystins. This review considers the exposure risk to humans, plants, and the environment due to the presence of microcystins in local water reservoirs used for drinking and irrigation. Additional studies are needed to understand the specific health impacts associated with the consumption of microcystin-contaminated agricultural plants.

Keywords: microcystin; contamination; toxicity; agriculture; health

Key Contribution: The review highlights the contamination and toxicity due to microcystins in agricultural plants, while emphasizing their associated risks with human health.



Citation: Melaram, R.; Newton, A.R.; Chafin, J. Microcystin Contamination and Toxicity: Implications for Agriculture and Public Health. *Toxins* **2022**, *14*, 350. <https://doi.org/10.3390/toxins14050350>

Received: 27 March 2022

Accepted: 12 May 2022

Published: 17 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Eutrophication is an environmental problem characterized by excessive algal and plant growth in aquatic ecosystems. Such a phenomenon largely occurs from agricultural nutrient runoff and continues to worsen due to global climate change, human population growth, and industrialization [1,2]. The symptoms of eutrophication have increased throughout coastal waters, drinking water reservoirs, estuaries, inland lakes, and slow-moving rivers, resulting in hypoxia, poor water quality, and noxious harmful algal blooms [2–4]. Harmful algal blooms, such as cyanobacterial blooms, commonly occur in eutrophic waters and may cause unsightly blue-green hues and release toxic secondary metabolites into the surrounding environment. Multiple environmental factors can promote cyanobacterial bloom formation and cyanotoxin production including environmental pollutants, light intensity, nutrients, salinity, pH, temperature, trace metals, and ultraviolet radiation [5].

Microcystins are the most prevalent and studied cyanotoxins within freshwater cyanobacterial blooms [6]. These biotoxins contain a rare 3-amino-9-methoxy-2,6,8-trimethyl-10-phenyl-4,6-decadienoic acid (ADDA) with two variable L amino acid residues [7]. The literature confirms over 300 variants of microcystin [8], with microcystin-leucine arginine (MC-LR) as the most important one due to its global distribution, ubiquity, and toxicity [9]. Multiple bloom-forming species (toxigenic and non-toxigenic) can produce MC-LR although *Microcystis* remains its primary manufacturer during cyanobacterial bloom incidents [10,11]. The presence of MC-LR in the environment constitutes an ecological and human health risk as concentration levels may exceed provisional limits in drinking (1 µg/L) and recreational waters (10 µg/L) [12].

The consumption of contaminated drinking water, aerosol inhalation, and dermal contact represent the main pathways of exposure to microcystins [13]. Additional exposure pathways, though less common, include the ingestion of contaminated algae dietary supplements, aquatic organisms, and fruits and vegetables [14]. Many studies have identified potential associations between microcystin exposure and liver disease. Hepatotoxic microcystins were found to accumulate in the tissue of dialysis patients in a Brazilian treatment center, many of whom died of subsequent liver failure [15]. Fishermen who frequented freshwater systems were also found to have well above the World Health Organization's recommended amount of microcystins within their bloodstream (2.2–3.9 ng/mL) [16]. Freshwater microcystins in drinking water sources were identified as one potential risk factor for primary liver cancer in endemic China [17]. Similarly, in central Serbia, freshwater toxic cyanobacterial blooms releasing microcystins correlated with primary liver cancer incidence and mortality [18].

Microcystins are potent hepatotoxins and inhibit serine and threonine phosphatases in living organisms including animals and plants [19,20]. MC-LR is the most common and toxic variant of the microcystins; it traverses the cell membrane into hepatocytes through organic anion-transporting polypeptides (OATPs) [21]. MC-LR uptake is mediated by OATP1B1, OATP1B3, OATP1A2, and OATP1B2, and many of these OATPs are expressed in various organs (brain, kidney, small and large intestines, stomach), indicating the possible extensive toxicity of MC-LR [22]. MC-LR in mice has been demonstrated to cause alterations in mitochondrial membrane permeability and apoptosis [23], and prolonged exposure can suppress hematopoiesis function, leading to normocytic anemia [24].

Eutrophic waters containing hazardous microcystin concentrations has caused great concern for the agricultural industry. Repeated use of microcystin-contaminated water can alter the nutritional quality of crops, contaminate agricultural soils and plants, inhibit plant growth, and reduce crop production and yield [25]. For example, plant exposure to microcystins has been found to negatively impact tissue organization, cytoskeleton and membrane integrity, and the ability to uptake nutrients and manufacture sugars [25]. Rice seedlings exposed to microcystins of 100 µg/L were significantly affected in their growth and height, adding to the dry weights of leaves and roots [26]. A study examined the effects of microcystin buildup in agricultural crops, finding cabbage, dill, and parsley plants accumulated threefold greater amounts of microcystins compared to fruiting crops. Another study observed a direct correlation between the amount of irrigation water used to water leafy vegetables and microcystin-contaminated crop soils, with leafy vegetables having 1.2–3.3 times more water used for irrigation purposes [27].

Lakes, streams, rivers, lagoons, and reservoirs account for the main sources of irrigation and drinking water worldwide. Microcystin variants have been detected in all these water reservoirs, increasing the potential likelihood of biotoxin transport into crop and drinking water systems [28]. Microcystins have also been found in groundwater samples, which account for half of the drinking water available on Earth [28]. With excess fertilizer and nutrient use over the past few decades [29], the number of cyanobacteria and their associated toxins in eutrophic waters is expected to increase, suggesting more significant and persistent toxic harmful algal blooms. Agricultural water use from eutrophic sources has significantly impacted over 160 terrestrial food crop species [30]. Leafy greens, such as

lettuce, spinach, watercress, and culinary herbs show the greatest uptake of microcystins into tissue most likely to be used in food preparation worldwide [30].

The purpose of the review is to reveal the persistence of toxic microcystins in water reservoirs and crops soils and their resultant impacts on agricultural plants and public health. The review is presented in chronological fashion to understand the interrelationships between water, soil, crops, and health in the context of microcystins. We first describe microcystin contamination in surface water, groundwater, and soil, followed by the most common input pathways that the cyanotoxins depend on for bioaccumulation in crops. Next, we cover the biosynthetic origin, mechanism of action, and toxicity of microcystins at the ecosystem and cellular levels. The review also considers the most recent works on toxic microcystins in agricultural plant parts and their potential health risks in humans, with the latter necessitating future studies.

2. Microcystins

The microcystins are perhaps the most widespread cyanotoxins in the natural environment and account for a maximum dry weight of 1% in terms of cyanobacterial mass [31]. Microcystin production is stimulated by important environmental factors such as light intensity, pH, temperature, trace metals, and ratio of total nitrogen to total phosphorus [5]. Microcystins belong to a family of monocyclic heptapeptides with ADDA and Mdha (*N*-methyldehydroalanine) binding moieties and two variable sites of amino acids. These structural components facilitate the binding of microcystin congeners to protein phosphatases in organisms while the variable sites influence toxicity [32]. MC-LR remains the priority congener in the environment and causes acute and chronic health effects in humans [12]. The environmental fate of microcystins remains unclear but influences the relationship with ecological and human health [33]. Contamination of water supplies and soil may provide insight into the environmental fate of microcystins relative to agricultural impacts and human health risks.

3. Contamination

3.1. Surface Water

The occurrence of microcystins in surface waters at hazardous levels can negatively affect ecosystem health. Microcystin concentrations generally remain below 100 µg/L in microcystin-polluted waters, although environmental conditions (nutrients, pH, and water temperature) can encourage concentrations greater than 1000 µg/L. The abundance of *Microcystis* morphospecies in waters has contributed to elevated microcystin concentrations [34–36]. *Microcystis aeruginosa* is reported as the high microcystin-producing morphospecies, followed by *Microcystis viridis* (low) and *Microcystis wesenbergii* (non-microcystin-producing) [37,38]. Microcystin contaminants are resistant to enzymatic degradation, high temperatures, pH alterations, and ultraviolet radiation owing to their heterocyclic structure [39,40]. Therefore, microcystin stability may facilitate transport into groundwater and agricultural soils in areas enduring cyanobacterial blooms.

3.2. Groundwater

After microcystins gradually accumulate in soil, they are released and migrate to lower soil horizons following precipitation events. Upon reaching deep soil layers, microcystins percolate into groundwater supplies and impair drinking water quality [41]. However, it has been reported that the use of contaminated groundwater wells for agricultural irrigation results in microcystin bioaccumulation among vegetable plants. A study in Saudi Arabia revealed microcystin levels in groundwater wells surpassing the World Health Organization's guidance value of 1.0 µg/L, along with vegetable leaves and roots containing levels up to 1.2 µg/g fresh weight [42]. Microcystins from toxic cyanobacterial blooms accounted for a high concentration of 1.07 µg/L in a groundwater well hundreds of meters from the western coast of Lake Chaohu, China, an important source of water for drinking,

washing, and irrigation [43]. These studies imply the hazardous nature of microcystins and their implications for agricultural practices, especially when they accumulate in crop soils.

3.3. Soil

Soil contamination from microcystins can disturb essential microorganisms of crop soil, which can accelerate toxin bioaccumulation in agricultural plants. Oxidative stresses caused by medial lethal concentrations of MC-LR (filter paper test = 0.149 µg/cm; acute soil test = 0.460 mg/kg) were discovered to exert profound impacts on the hatchability and survival of earthworms [44]. Frequent exposure of sediment bacteria to MC-LR was shown to inhibit the overall carbon metabolic activity of a microbial community [45]. These findings indicate the toxic effects of MC-LR on soil microorganisms, a critical biotic component involved in plant growth and development. Bioaccumulation in plants occurs after microcystin uptake, and the amount and time of absorption into edible plant tissue raises a food security concern [46,47]. Additionally, limited knowledge exists regarding the impact of microcystins on bacteria, fungi, and other essential organisms associated with plant growth. Although soil assists in microcystin uptake, the direct factors affecting its bioaccumulation in plants remain unknown [47]. Therefore, an understanding of the effects of microcystin input pathways through soils on crop bioaccumulation and toxicity, considering variations in the bioavailability and speciation of microcystins, may be beneficial.

4. Input Pathways

4.1. Irrigation with Polluted Water

Irrigation with polluted water is the primary route of introduction of cyanotoxins (microcystins) in agricultural soils. Extracellular microcystins within irrigation water may leach into soils from constant spray, increasing the bioavailability of toxin [48]. Two studies indicated that contaminated irrigation water with microcystins threaten crop quality and yield. One study confirmed high uptake values of MC-LR and microcystin-leucine phenylalanine (MC-LF) in the roots of alfalfa and wheat (1.3 mg/kg dry weight) [49]. A greenhouse experiment measured microcystin content in the roots of clover, lettuce, rape, and ryegrass, with clover presenting the highest total microcystin concentrations (1.45 mg/kg dry weight). Since microcystins can bioconcentrate in plant tissue, they can enter the animal and human food chains at concentrations above the recommended tolerable limits [50]. The tolerable daily intake value for chronic exposure to MC-LR is 0.04 µg/kg body weight, as proposed by the World Health Organization [12].

4.2. Application of Cyanobacterial Manure

Cyanobacterial fertilizer, which contains living cyanobacteria cultures that aid in harnessing solar energy, nutrients, and water resources essential to plant growth have become popular in recent years [51]. The application of cyanobacterial fertilizer in tandem with manure has the potential to improve soil quality and plant growth and reduce crop production costs in agriculture. However, its use in soils affected by cyanobacterial blooms is not recommended due to higher bioaccumulation capacity in edible plants from polluted soil [51,52]. Intracellular microcystins are contained in cyanobacterial manure, and cell lysis can facilitate the bioavailability of microcystins in soil. Furthermore, the co-existence of microcystin congeners in manure is considered a human health risk. These cyanotoxins have been associated with tumor promotion in animals and liver and colon cancer in humans [53]. Careful and proper application of cyanobacterial manure in soils may improve agricultural plant quality and protect public health from contaminated agricultural foods.

4.3. Compost

Compost, a mixture containing decaying organic matter has traditionally been used as a soil conditioner and fertilizer to aid vigorous plant growth. The agricultural use of compost has risen in the past decade to foster improved soil ecosystem health for greater

crop outputs with less dependence on chemical inputs [54]. While the negative influence of microcystin contamination is well documented for marine invertebrates, few studies have indicated its effect on soil-dwelling species. Soil nematodes were shown to have a reduction in overall lifespan and a decrease in reproduction and were less motile in the presence of a mere 1.0 µg/kg MC-LR concentration, with any higher dose largely obliterating nematode existence [55]. While this might be a positive for the control of harmful plant-feeding species, the broad-spectrum nature means it is equally problematic for beneficial species [47]. Earthworms who digest large quantities of organic matter in the soil layer are more likely to absorb high concentrations of MC-LR [44]. Interestingly, when compost levels decreased in total concentration, earthworms continued to show high levels, perhaps due to bioaccumulation.

Cyanobacteria have demonstrated promise in the removal of harmful pathogens in manure and compost; however, microcystin toxins reduce the overall activity of beneficial bacteria and fungi involved in nitrogen fixation [56]. Compost used for crop production has increased interest in the likelihood of unintentional contamination [56]. As has been the case for pesticides, leafy greens were most vulnerable to the accumulation of hazardous levels of microcystin, representing a potential health risk to consumers [57]. Higher mean concentrations of MCs, in the 12.3–22.8 µg/kg range, were found in lettuce and cabbage, as compared to root vegetables such as carrot (10.5–12.6 µg/kg) that were grown in compost-rich soil [58]. The results imply possible health concerns in agricultural areas where compost material is preferred over other soil applicators and fertilizers.

5. Toxicity

5.1. Biosynthesis

Microcystins are toxic peptides of non-protein amino acids encoded by non-ribosomal peptide synthetase (NRPS)/polyketide synthase enzymes [59]. Genetic analysis has indicated that a large NRPS gene cluster is involved in promoting microcystin biosynthesis. DNA sequence results from *Microcystis* strain K-139 identified an *mycABC* operon and revealed a microcystin synthetase gene operon (*mycD*, *mycE*, *mycF*, *mycG*) within the NRPS gene cluster [59]. NRPS enzymes use a thiotemplate mechanism to catalyze the synthesis of peptides in prokaryotes and lower eukaryotes. Many cyclic peptides are synthesized by NRPS enzymes including the antimicrobial agents, cyclosporine, penicillin, and vancomycin. NRPS products are highly diverse in chemical structure, owing to incorporated hydroxylated and methylated amino acids [60]. Covalent modifications of amino acid residues may explain the myriad of microcystin variants detected in freshwater sites, some of which present a danger to ecosystem aquatic and terrestrial agricultural plants.

5.2. Mechanism of Action

Microcystins are cyclic heptapeptides with a chemical structure of Adda-D-Glu-Mdha-D-Ala-L-X-D-MeAsp-L-Z, where X and Z refer to variable sites of amino acid residues [60]. Microcystin-leucine arginine (MC-LR) is the most prevalent and toxic of all microcystin congeners [9]. The mechanism of action of MC-LR involves the inhibition of serine and threonine phosphatases [19,20]. Microcystins exert a strong affinity toward protein phosphatase 1 (PP1) and 2A (PP2A) and, to a lesser extent, protein phosphatase 2B (PP2B) [61]. A two-step mechanism occurs between microcystin and PP1 and PP2A, wherein microcystin inactivates catalytic subunits (PP1-c and PP-2ac) through rapid binding, followed by an extended period of covalent interaction [62]. Accessibility to PP1-c's (catalytic subunit) active site is blocked when the carboxyl terminus of MeAsp interacts with Arg96 and Tyr134 [63]. When its active site is blocked, PP1 can no longer bind to its substrates, preventing the removal of phosphates from these molecules, which are critical to maintaining cellular homeostasis in biological organisms. MC-LR toxicity has been well studied, as supported by changes in liver structure and cytoskeleton deformities in animals and plants, respectively. The following section describes some recent findings on the adverse effects of toxic microcystins in the local environment as well as in plants.

5.3. Ecotoxicity

Microcystins are secondary metabolites produced by cyanobacteria in the aquatic environment. The exposure of aquatic organisms to microcystins has been shown to alter behavior, cause bioaccumulation in tissues, inhibit growth, and reduce fecundity and survivorship [64]. Toxic microcystins have resulted in episodes of fatality to aquatic biota, livestock, and wild animals worldwide [65]. The biological detoxification pathways of resident organisms are major determinants of microcystin toxicity in the ecosystem [33]. Biotransformation of microcystins into conjugate intermediates via the glutathione pathway is conserved in many aquatic organisms, which may offer protection from toxicologic effects of microcystins. The metabolic process serves to eliminate microcystins from the organism and release them back in the natural environment [33]. However, toxin release from exposed organisms can harm other aquatic organisms and humans as microcystins transfer to higher trophic levels in the food chain. Microcystins can also leach into groundwater and transfer into soils for agricultural production [41]. Thus, we present the human health risks associated with microcystins in waters sourced for irrigation and crop soils in agriculture.

5.4. Phytotoxicity

Phytotoxicity, a delay in overall seed germination, inhibition of plant growth, or other adverse effect on plant growth and development, has been attributed to the presence of microcystins [57]. Dephosphorylation is a critical process of cell cycle control and protein modification in living organisms, including higher plants. However, the inhibition of dephosphorylation can negatively affect cell function, causing hyperphosphorylation among cytokeleatin proteins and significant alterations in cytoskeletal elements [66,67]. More precisely, microcystins can associate with F-actin and microtubules, altering F-actin organization and disorienting microtubules in plant cells [68]. This directly impacts photosynthesis activity, with overall chlorophyll concentration decreasing as much as 0.80 mg/g [68,69]. Stomatal integrity is similarly impacted by the weakening cytoskeletal elements and can lead to oxidative stress including less leaf transpiration and poor gas exchange [69]. Roots also indicated adverse influence, with low concentrations (<250 µg/L) causing an overall reduction in the ability to uptake necessary nutrients. Microcystin concentrations above 250 µg /L caused total inhibition of assemblage of nutrients at the root level [69]. The metabolism of nitrogen, an essential element for plant growth, shows significant decrease when aquatic and terrestrial plants are placed in concentrations of 1.0 µg/L of microcystin. Enzymatic activity that assists in sequestering nitrogen into tissue, including glutamic transaminase and glutamate synthase, are unable to function at this low concentration level [70].

6. Agricultural Plants

Plant growth, whether in terrestrial or aquatic form, has shown sensitivities to microcystins [70–73]. The two are distinguished from each other by the traditional definition of true aquatic plant, meaning it must be submersed in water for most of its life cycle [74]. Terrestrial plants, in contrast, spend their life cycle on land and root directly into the soil [75]. Those that can withstand periods of standing water are still classified as terrestrial, land-dwelling plants [75]. Of the two, terrestrial plants have received significant attention where crops are concerned, as those are directly consumed [76]. The effects on tissue and seedling growth are presented below (Table 1).

Table 1. Physiological effects in agricultural plants from various microcystin exposure routes.

Species	Stage of Development	Route of Exposure	Microcystin Present	Concentration of Microcystin ($\mu\text{g/L}$)	Physiological Effects	Reference
<i>Brassica juncea</i> (mustard greens)	Mature Plants	Soil	MCs	150	Reduced plant height and weight	[70–74]
<i>Brassica napus</i> (rape seed)	Seeds	Soil	MCs	600–3000	Reduced germination, conductivity, and phototrophic response	[70]
<i>Dacus carota</i> (carrot)	Mature Plants, Seeds	Soil	MCs	50	Reduced biomass of leaves and taproot	[71,72]
<i>Ipomoea batatas</i> (sweet potato)	Mature Plants	Soil	MCs	150	Reduced plant height and weight	[77]
<i>Lactuca sativa</i> (lettuce)	Seeds, Seedlings	Germination, Soil	MCs, MC-LR	2–5000	Reduced radicle length	[75,76]
<i>Lepidium sativum</i> (watercress)	Seeds	Germination, Soil	MCs, MC-LR, MC-RR	500–20,000	Reduced radicle length and shoot weight	[76]
<i>Medicago sativa</i> (alfalfa)	Seeds, Seedlings	Irrigation	MCs	5–20	Reduced nodule production, reduced nodule number, reduced biomass	[77,78]
<i>Oryza sativa</i> (rice)	Seedlings	Hydroponics	MCs, MC-LR	1–3000	Reduced biomass of leaves, stems, and roots, reduced starch in grains, reduced root surface area	[78]
<i>Phaseolus vulgaris</i> (green bean)	Mature Plants, Seeds	Germination, Soil	MCs	1–15,000	Reduced chlorophyll content, delayed development, reduced bean length and number produced	[78]
<i>Spinacia oleracea</i> (spinach)	Mature Plants, Seedlings	Soil, Hydroponics	MCs, MC-LR	0.5	Reduced photosynthesis and biomass production	[79]
<i>Solanum lycopersicum</i> (tomato)	Developed Plants, Seeds	Soil	MCs	3–1000	Reduced biomass, reduced flower production, low sugar content	[76,77]
<i>Triticum aestivum</i> (wheat)	Seeds, Seedlings	Germination, Soil	MCs	0.5–20,000	Reduced germination, reduced photosynthesis, reduced biomass	[77]
<i>Zea mays</i> (corn)	Seeds, Seedlings	Germination, Hydroponics, Soil	MCs, MC-LR, MC-RR	5–3000	Reduced plant height, reduced root weight	[72]

Total concentration of microcystins includes all studied variants. Concentration depicts range covered across multiple studies on same crop species, with first number being for the earliest stage of development. MCs = total concentration of all studied microcystin variants; LR = microcystin leucine arginine; RR = microcystin arginine-arginine; YR = microcystin tyrosine arginine.

6.1. Plant Seedling Growth

Rigorous regulation of microcystin-contaminated water use for irrigation is not actively practiced worldwide. As a result, entire crop networks are susceptible to potentially

toxic bacteria, including cyanobacteria, which may directly impact the human food system. In the past decade, studies have measured the effects of varying concentrations of microcystins on the growth of agriculturally important crops, including leafy greens, herbs, root vegetables, and squash [76]. Height, biomass, leaf surface area, seedling diameter, and root development were significantly reduced across all crops (root vegetables, herbs, leafy greens, and squash) tested, with leafy greens holding the highest risk to exposure [77]. While all vegetable and fruit crops studied demonstrated negative impacts, the seedling stage appeared the most vulnerable to uptake of microcystins into developing tissue. *Rhizobia* nodules, which are essential for many growing seedlings to successfully fix and uptake nitrogen, are reduced or non-existent when grown in high concentrations of microcystins [77,78]. As a result, root surface area, length, number of roots present, and total biomass are reduced by upward of 60% in leafy green vegetables [78].

6.2. Tissue Growth

Irrigating mature crops with microcystin-contaminated water also leads to poor nitrogen fixation, reduced height, and reduction in total biomass in tissue development [70,71]. As seen with seedlings, the higher the concentration of microcystins present in irrigation water, the more at-risk tissues are for reduced photosynthesis, decreased leaf and root production, and an overall reduction in biomass [74,77–80]. In a survey of 38 crop species ranging from leafy greens to root vegetables, all reported losses of at least 30% in total leaf surface area [80]. Further analysis suggests that microcystins play a direct role in the inhibition of photosynthetic activities and can cause tissue death, increase oxidative stress, reduce membrane integrity, and impair the ability of roots to absorb nutrients [74,75,77–80]. Anomalies in tissue structure were apparent in crops grown hydroponically, with high concentration of microcystins ($\geq 10 \mu\text{g/L}$) [77]. Additional alternations, such as slowing of mitotic and metabolic processes and reduced development, increased with the increase of concentration [72,74,80].

Despite the effects of microcystins on seedling and tissue growth, they are not the sole contributing factor in plant developmental changes [74]. The impacts of microcystins are greatest at higher concentrations, with the greatest effect on the seedling stage [70,71,81,82]. Due to the structure, surface area, and gas exchange exhibited by leafy greens, microcystins can readily diffuse through stomata into plant tissue and cause negative impacts [73,77,80]. Hydroponically grown species of leafy greens including watercress, lettuce, and spinach, over those grown in soil-based cultures, were 30% more likely to have hinderance of growth [83]. A study identified root growth inhibition within *Lactuca sativa* (lettuce) from exposure to *Microcystis aeruginosa* strains [84]. Photosynthesis activity within *Lycopersicon esculentum* leaves was not affected when exposed to a pure concentration of MC-LR (100 $\mu\text{g/L}$) [85]. Additional analysis is needed to determine whether microcystin concentrations in irrigation water used for hydroponically grown lettuce and herbs pose a risk to human health.

6.3. Aquatic Plants

Microcystins can readily enter many aquatic plant species through simple diffusion, stomata of leaves in contact with water, and absorption via the root system [86]. Several cases have documented accumulated microcystin uptake in both field and laboratory settings. Submerged species show the highest concentrations present in tissue, with *Elodea canadensis* showing a concentration of 16.9 $\mu\text{g/g}$ of dry weight [87]. *Lemna minor* (duckweed) and *Wolffia arrhizal* (duckweed) demonstrated a reduction in growth over the course of a 5-day exposure study [88]. Clearly, method of uptake, plant species, concentration, and type of microcystins present, and length of exposure play a significant role in determining accumulation (Table 2).

While microcystin accumulation proves harmful to essential processes such as photosynthesis, overall height, and weight of the mature plant, aquatic species can detoxify the bacteria through the action of a natural antioxidant, glutathione S-transferases (GSTs)

(Table 2). High levels of GSTs found in contaminated plant tissue suggests that microcystin presence may trigger their production [89]. Laboratory studies of several different species show similar results, signaling the likelihood of microcystins contributing to the enhancement of GST concentrations in exposed aquatic plants. In a timed-exposure study of cyanobacteria and microcystins, Lemnaceae sp. (duckweed), experienced increased concentrations of GSTs present within a 24 h timespan [90]. In another timed-exposure study, similar effects were seen in *Vallisneria natans* [91]. Perhaps most impressive, bioaccumulation of microcystin in *Lemna minor* after 5 days of exposure showed resistance to a concentration level (2.5 µg/L) that had earlier been detrimental [88]. More studies are needed to explore the possible implications of GST for resistance against microcystins.

Table 2. Accumulation of microcystins in aquatic plants.

Species	Environment	Microcystin Toxins	Concentration of Microcystin (µg/g Dry Weight)	Mode of Uptake	Plant Response	References
<i>Alternanthera philoxeroides</i> (alligator weed)	Submerged	MCs	0.19–3.95	Diffusion	Detoxification via tGSH production	[85]
<i>Ceratophyllum demersum</i> (hornwort)	Submerged	MC-LR	0.05	Root absorption, diffusion	Detoxification via tGSH production	[89]
<i>Elodea canadensis</i> (American waterweed)	Floating	MC-LR, RR, YR	15,000	Root absorption, leaf contact with the water surface	Detoxification via tGSH production	[87]
<i>Hydrilla verticillate</i> (water thyme)	Submerged	MCs	2–5000	Diffusion, root absorption	Detoxification via tGSH production	[88]
<i>Lemna gibba</i> (duckweed)	Floating	MC-LR, RR	2.44	Root absorption, leaf contact with the water surface	Detoxification via tGSH production	[88]
<i>Phragmites australis</i> (common reed)	Shore	MC-LR, RR, YR	2.7	Root absorption, stem, or leaf contact with water surface	Reduction in photosynthesis, reduction in biomass	[88]
<i>Polygonum portoricensis</i> (smooth smartweed)	Submerged	MC-LR, RR, YR	4850–7000	Root absorption, diffusion	Detoxification via tGSH production	[86]
<i>Trapa natans</i> (water chestnut)	Floating	MCs	7.0	Reduced chlorophyll content, delayed development, reduced bean length and number produced	Detoxification via tGSH production	[86]
<i>Typha</i> sp. (cattail)	Shore	MC-LR, YR	16,000–17,000	Root absorption, stem, or leaf contact with water surface	Reduction in photosynthesis, reduction in biomass	[88]
<i>Vallisneria natans</i>	Submerged	MC-RR	0.3–14.8	Diffusion	Detoxification via tGSH production	[91]

Total concentration of microcystins includes all studied variants; LR = microcystin leucine arginine; RR = microcystin arginine-arginine; YR = microcystin tyrosine arginine; tGSH: glutathione S-transferase.

7. Human Health Risks

Epidemiological research has indicated potential health effects of microcystins from human consumption of contaminated drinking water and aquatic foods [57]. A recent work posits environmental microcystins as an emergent risk factor in endemic regions plagued by hepatocellular carcinoma [92]. It is possible that these regions continually rely on bloom-forming water reservoirs for drinking water, food, and irrigation. Nevertheless, little remains known about the human health risks of microcystins in agricultural products. Given the discussed high levels of microcystin accumulation in leafy vegetables, including lettuces and spinach when grown through use of contaminated irrigation, the possibility of health risk is present [57,73]. Since leafy greens are now popular to grow hydroponically, their sensitivity to microcystins and other bacteria uptake needs to be studied and taken into serious consideration from a health standpoint. This also raises questions and concerns regarding the potential health risks to agricultural workers who are directly in contact with irrigation water and thus exposed to high microcystin concentrations [73,74].

8. Implications for Future Research

The overall findings of this review depict a clear presence of the microcystin in drinking water and crop systems, raising concerns of health risk and food safety. This review indicates a need for further research in areas experiencing anthropogenic eutrophication of water reservoirs. The increase in microcystins in enriched water sources used for irrigation and consumption raises concern over the health risks posed to nearby communities. Studies examining soil- and hydroponic-based agricultural systems have indicated plant sequestering of microcystins into plant tissue, with leafy greens showing the highest numbers. More research is needed, particularly on a larger scale, to further examine the relationship of microcystins with hydroponic, in soil, and laboratory-based conditions. At present, few studies have explored the presence of microcystins in media amendments such as manure and compost, which could also add to the number of toxic microcystins present in agricultural plants. Although drinking water contamination from microcystin appears well studied, the food safety risks associated with the irrigation of agricultural crops, both in small- and large-scale agriculture, could benefit from additional review. Several studies indicate the possibility of microcystins eliciting a response to release glutathione S-transferases to detoxify potential negative effects of the toxin. More research in this arena could provide more insight into how this could be employed in large-scale detoxification of water used in environmental and agricultural contexts.

Author Contributions: R.M. proposed the research topic and contributed significantly to background material on microcystin and research on microcystin contamination and toxicity. A.R.N. integrated substantive research on the effects of microcystin exposure in agricultural and aquatic plant species. J.C. researched several areas including but not limited to eutrophication, microcystin occurrence, and microcystin uptake in plants. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Withers, P.J.A.; Neal, C.; Jarvie, H.P.; Doody, D.G. Agriculture and eutrophication: Where do we go from here? *Sustainability* **2014**, *6*, 5853–5875. [[CrossRef](#)]
2. Rabalais, N.N.; Turner, R.E.; Díaz, R.J.; Justić, D. Global change and eutrophication of coastal waters. *ICES J. Mar. Sci.* **2009**, *66*, 1528–1537. [[CrossRef](#)]

3. Duncan, E.; Kleinman, P.J.; Sharpley, A.N. *Eutrophication of Lakes and Rivers*; John Wiley & Sons: Chichester, UK, 2012.
4. Padedda, B.M.; Sechi, N.; Lai, G.G.; Mariani, M.A.; Pulina, S.; Sarria, M.; Satta, C.T.; Viridis, T.; Buscarinu, P.; Lugliè, A. Consequences of eutrophication in the management of water resources in Mediterranean reservoirs: A case study of Lake Cedrino (Sardinia, Italy). *Glob. Ecol. Conserv.* **2017**, *12*, 21–35. [[CrossRef](#)]
5. Neilan, B.A.; Pearson, L.A.; Muenchhoff, J.; Moffitt, M.C.; Dittmann, E. Environmental conditions that influence toxin biosynthesis in cyanobacteria. *Environ. Microbiol.* **2013**, *15*, 1239–1253. [[CrossRef](#)] [[PubMed](#)]
6. Welker, M.; von Döhren, H. Cyanobacterial peptides—Nature’s own combinatorial biosynthesis. *FEMS Microbiol. Rev.* **2006**, *30*, 530–563. [[CrossRef](#)]
7. Van Apeldoorn, M.E.; Van Egmond, H.P.; Speijers, G.J.A.; Bakker, G.J.I. Toxins of cyanobacteria. *Mol. Nutr. Food Res.* **2007**, *51*, 7–60. [[CrossRef](#)]
8. Bailiu-Rodriguez, D.; Peraino, N.J.; Premathilaka, S.H.; Birbeck, J.A.; Baliu-Rodriguez, T.; Westrick, J.A.; Isailovic, D. Identification of novel microcystins using high-resolution MS and MSn with python code. *Environ. Sci. Technol.* **2022**, *56*, 1652–1663. [[CrossRef](#)]
9. Cao, L.; Huang, F.; Massey, I.Y.; Wen, C.; Zheng, S.; Xu, S.; Yang, F. Effects of microcystin-LR on the microstructure and inflammation-related factors of jejunum in mice. *Toxins* **2019**, *11*, 482. [[CrossRef](#)]
10. Catherine, A.; Bernard, C.; Spoof, L.; Bruno, M. Microcystins and Nodularins. In *Handbook of Cyanobacterial Monitoring and Cyanotoxin Analysis*; Meriluoto, J., Lisa Spoof, L., Codd, G.A., Eds.; John Wiley & Sons, Ltd.: Chichester, UK, 2017; pp. 107–126.
11. Sivonen, K.; Jones, G. Cyanobacterial Toxins. In *Toxic Cyanobacteria in Water: A Guide to Their Public Health Consequences, Monitoring and Management*; Chorus, I., Bartram, J., Eds.; E and FN Spon: London, UK, 1999; pp. 41–111.
12. WHO. Cyanobacterial Toxins: Microcystins. In *Guidelines for Drinking Water Quality and Guidelines for Safe Recreational Water Environments*; World Health Organization: Geneva, Switzerland, 2020.
13. Lad, A.; Breidenbach, J.D.; Su, R.C.; Murray, J.; Kuang, R.; Mascarenhas, A.; Najjar, J.; Patel, S.; Hegde, P.; Youssef, M.; et al. As we drink and breathe: Adverse health effects of microcystins and other harmful algal bloom toxins in the liver, gut, lungs and beyond. *Toxins* **2022**, *12*, 418. [[CrossRef](#)]
14. Drobac, D.; Tokodi, N.; Simeunović, J.; Baltić, V.; Stanić, D.; Svirčev, Z. Human exposure to cyanotoxins and their effects on health. *Arhiv za Higijenu Rada i Toksikologiju* **2013**, *64*, 119–130. [[CrossRef](#)]
15. Jochimsen, E.M.; Carmichael, W.W.; An, J.S.; Cardo, D.M.; Cookson, S.T.; Holmes, C.E.; Antunes, M.B.; de Melo Filho, D.A.; Lyra, T.M.; Barreto, V.S.; et al. Liver failure and death after exposure to microcystins at a hemodialysis center in Brazil. *N. Eng. J. Med.* **1998**, *338*, 873–878. [[CrossRef](#)] [[PubMed](#)]
16. Ueno, Y.; Nagata, S.; Tsutsumi, T.; Hasegawa, A.; Watanabe, M.F.; Park, H.D.; Chen, G.C.; Yu, S.Z. Detection of microcystins, a blue-green algal hepatotoxin, in drinking water sampled in Haimen and Fusui, endemic areas of primary liver cancer in China, by highly sensitive immunoassay. *Carcinogenesis* **1996**, *17*, 1317–1321. [[CrossRef](#)]
17. Chen, J.; Xie, P.; Li, L.; Xu, J. First identification of the hepatotoxic microcystins in the serum of a chronically exposed human population together with indication of hepatocellular damage. *Toxicol. Sci.* **2009**, *108*, 81–89. [[CrossRef](#)] [[PubMed](#)]
18. Svirčev, Z.; Drobac, D.; Tokodi, N.; Lužanin, Z.; Munjas, A.M.; Nikolin, B.; Vuleta, D.; Meriluoto, J. Epidemiology of cancers in Serbia and possible connection with cyanobacterial blooms. *J. Environ. Sci. Health Part C* **2014**, *32*, 319–337. [[CrossRef](#)] [[PubMed](#)]
19. MacKintosh, C.; Beattie, K.A.; Klumpp, S.; Cohen, P.; Codd, G.A. Cyanobacterial microcystin-LR is a potent and specific inhibitor of protein phosphatases 1 and 2A from both mammals and higher plants. *FEBS Lett.* **1990**, *264*, 187–192. [[CrossRef](#)]
20. Dawson, R.M. The toxicology of microcystins. *Toxicon* **1998**, *36*, 953–962. [[CrossRef](#)]
21. Greer, B.; Meneely, J.P.; Elliott, C.T. Uptake and accumulation of Microcystin-LR based on exposure through drinking water: An animal model assessing the human health risk. *Sci. Rep.* **2018**, *8*, 4913. [[CrossRef](#)]
22. Fischer, W.J.; Altheimer, S.; Cattori, V.; Meier, P.J.; Dietrich, D.R.; Hagenbuch, B. Organic anion transporting polypeptides expressed in liver and brain mediate uptake of microcystin. *Toxicol. Appl. Pharmacol.* **2005**, *203*, 257–263. [[CrossRef](#)]
23. Lezcano, N.; Sedán, D.; Lucotti, I.; Giannuzzi, L.; Vittone, L.; Andrinolo, D.; Mundiña-Weilenmann, C. Subchronic microcystin-LR exposure increased hepatic apoptosis and induced compensatory mechanisms in mice. *J. Biochem. Mol. Toxicol.* **2012**, *26*, 131–138. [[CrossRef](#)]
24. Zhou, W.; Zhang, X.; Xie, P.; Liang, H.; Zhang, X. The suppression of hematopoiesis function in Balb/c mice induced by prolonged exposure of microcystin-LR. *Toxicol. Lett.* **2013**, *219*, 194–201. [[CrossRef](#)]
25. Campos, A.; Redouane, E.M.; Freitas, M.; Amaral, S.; Azevedo, T.; Loss, L.; Máthé, C.; Mohamed, Z.A.; Oudra, B.; Vasconcelos, V. Impacts of microcystins on morphological and physiological parameters of agricultural plants: A review. *Plants* **2021**, *10*, 639. [[CrossRef](#)] [[PubMed](#)]
26. Liang, C.; Wang, W. Response and recovery of rice (*Oryza sativa*) seedlings to irrigation with microcystin-contaminated water. *Environ. Ear. Sci.* **2015**, *73*, 4573–4580. [[CrossRef](#)]
27. Xiang, L.; Li, Y.W.; Liu, B.L.; Zhao, H.M.; Li, H.; Cai, Q.Y.; Mo, C.H.; Wong, M.H.; Li, Q.X. High ecological and human health risks from microcystins in vegetable fields in southern China. *Environ. Int.* **2019**, *133*, 105142. [[CrossRef](#)]
28. De Figueiredo, D.R.; Azeiteiro, U.M.; Esteves, S.M.; Gonçalves, F.J.M.; Pereira, M.J. Microcystin-producing blooms—A serious global public health issue. *Ecol. Environ. Saf.* **2004**, *59*, 151–163. [[CrossRef](#)] [[PubMed](#)]
29. Gobler, C.J.; Burkholder, J.M.; Davis, T.W.; Harke, M.J.; Johengen, T.; Stow, C.A.; Waal, D.B.V. The dual role of nitrogen supply in controlling the growth and toxicity of cyanobacterial blooms. *Harmful Algae* **2016**, *56*, 87–97. [[CrossRef](#)]

30. Mutoti, M.; Gumbo, J.; Jideani, A.I.O. Occurrence of cyanobacteria in water used for food production: A review. *Phys. Chem. Earth*. **2022**, *125*, 103101. [\[CrossRef\]](#)
31. Bláha, L.; Babica, P.; Maršálek, B. Toxins produced in cyanobacterial water blooms—Toxicity and risks. *Interdiscip. Toxicol.* **2009**, *2*, 36–41. [\[CrossRef\]](#) [\[PubMed\]](#)
32. Massey, I.Y.; Yang, F. A mini review on microcystins and bacterial degradation. *Toxins* **2020**, *12*, 268. [\[CrossRef\]](#)
33. Schmidt, J.R.; Wilhelm, S.W.; Boyer, G.L. The fate of microcystins in the environment and challenges for monitoring. *Toxins* **2014**, *6*, 3354–3387. [\[CrossRef\]](#)
34. Chorus, I.; Fastner, J.; Welker, M. Cyanobacteria and cyanotoxins in a changing environment: Concepts, controversies, challenges. *Water* **2021**, *13*, 2463. [\[CrossRef\]](#)
35. Peng, L.; Liu, Y.; Chen, W.; Liu, L.; Kent, M.; Song, L. Health risks associated with consumption of microcystin-contaminated fish and shellfish in three Chinese lakes: Significance for freshwater aquacultures. *Ecotoxicol. Environ. Saf.* **2010**, *73*, 1804–1811. [\[CrossRef\]](#) [\[PubMed\]](#)
36. Wu, Y.; Li, L.; Gan, N.; Zheng, L.; Ma, H.; Shan, K.; Liu, J.; Xiao, B.; Song, L. Seasonal dynamics of water bloom-forming *Microcystis* morphospecies and the associated extracellular microcystin concentrations in large, shallow, eutrophic Dianchi Lake. *J. Environ. Sci.* **2014**, *26*, 1921–1929. [\[CrossRef\]](#)
37. Chen, W.T. Analysis on eutrophication and cyanobacteria blooms characteristics in Dashahe Reservoir. *Guangdong Water Resour. Hydropower* **2015**, *6*, 33–35.
38. Harada, K.; Tsuji, K.; Watanabe, M.F.; Kondo, F. Stability of microcystins from cyanobacteria—III. Effect of pH and temperature. *Phycologia* **1996**, *35*, 83–88. [\[CrossRef\]](#)
39. Tsuji, K.; Naito, S.; Kondo, F.; Ishikawa, N.; Watanabe, M.F.; Suzuki, M.; Harada, K. Stability of microcystins from cyanobacteria: Effect of light on decomposition and isomerization. *Environ. Sci. Technol.* **1994**, *28*, 173–177. [\[CrossRef\]](#)
40. Rastogi, R.P.; Sinha, R.P.; Incharoensakdi, A. The cyanotoxin-microcystins: Current overview. *Rev. Environ. Sci. Biol. Technol.* **2014**, *13*, 215–249. [\[CrossRef\]](#)
41. Chen, W.; Song, L.; Gan, N.; Li, L. Sorption, degradation and mobility of microcystins in Chinese agriculture soils: Risk assessment for groundwater protection. *Environ Pollut.* **2006**, *144*, 752–758. [\[CrossRef\]](#)
42. Mohamed, Z.A.; Al Shehri, A.M. Microcystins in groundwater wells and their accumulation in vegetable plants irrigated with contaminated waters in Saudi Arabia. *J. Hazard Mater.* **2009**, *172*, 310–315. [\[CrossRef\]](#)
43. Yang, Z.; Kong, F.; Zhang, M. Groundwater contamination by microcystin from toxic cyanobacteria blooms in Lake Chaohu, China. *Environ. Monit. Assess.* **2016**, *188*, 280. [\[CrossRef\]](#)
44. Wen, H.F. Toxicological effects of microcystin-LR on earthworm (*Eisenia fetida*) in soil. *Biol. Fertil. Soils* **2017**, *53*, 849–860. [\[CrossRef\]](#)
45. Ding, Q.; Liu, K.; Song, Z.; Sun, R.; Zhang, J.; Yin, L.; Pu, Y. Effects of microcystin-LR on metabolic functions and structure succession of sediment bacterial community under anaerobic conditions. *Toxins* **2020**, *12*, 183. [\[CrossRef\]](#) [\[PubMed\]](#)
46. Corbel, S.; Mougin, C.; Nélieu, S.; Delarue, G.; Bouaïcha, N. Evaluation of the transfer and the accumulation of microcystins in tomato (*Solanum lycopersicum* cultivar Micro Tom) tissues using a cyanobacterial extract containing microcystins and the radiolabeled microcystin-LR (14 C-MC-LR). *Sci. Total Environ.* **2016**, *541*, 1052–1058. [\[CrossRef\]](#) [\[PubMed\]](#)
47. Cao, Q.; Steinman, A.D.; Wan, X.; Xie, L. Bioaccumulation of microcystin congeners in soil-plant system and human health risk assessment: A field study from Lake Taihu region of China. *Environ. Pollut.* **2018**, *240*, 44–50. [\[CrossRef\]](#) [\[PubMed\]](#)
48. Esterhuizen-Londt, M.; Pflugmacher, S. Vegetables cultivated with exposure to pure and naturally occurring β -N-methylaminol-alanine (BMAA) via irrigation. *Environ. Res.* **2019**, *169*, 357–361. [\[CrossRef\]](#) [\[PubMed\]](#)
49. Peuthert, A.; Chakrabarti, S.; Pflugmacher, S. Uptake of Microcystins-LR and -LF (cyanobacterial toxins) in seedlings of several important agricultural plant species and the correlation with cellular damage (lipid peroxidation). *Environ. Toxicol.* **2007**, *22*, 436–442. [\[CrossRef\]](#)
50. Crush, J.R.; Briggs, L.R.; Sprosen, J.M.; Nichols, S.N. Effect of irrigation with lake water containing microcystins on microcystin content and growth of ryegrass, clover, rape, and lettuce. *Environ. Toxicol.* **2008**, *23*, 246–252. [\[CrossRef\]](#)
51. Bouaïcha, N. Cyanobacterial toxins: Modes of actions, fate in aquatic and soil ecosystems, phytotoxicity and bioaccumulation in agricultural crops. *Chemosphere* **2014**, *96*, 1–15.
52. Singh, J.S.; Kumar, A.; Rai, A.N.; Singh, D.P. Cyanobacteria: A precious bio-resource in agriculture, ecosystem, and environmental sustainability. *Front. Microbiol.* **2016**, *7*, 529. [\[CrossRef\]](#)
53. Falconer, I.R. Tumor promotion and liver-injury caused by oral consumption of cyanobacteria. *Environ. Toxicol. Water Qual.* **1991**, *6*, 177–184. [\[CrossRef\]](#)
54. Wu, X.; Xiao, B.; Li, R.; Wang, C.; Huang, J.; Wang, Z. Mechanisms and factors affecting sorption of microcystins onto natural sediments. *Environ. Sci. Technol.* **2011**, *45*, 2641–2647. [\[CrossRef\]](#)
55. Yuhui, L.; Wang, Y.; Yin, L.; Pu, Y.; Wang, D. Using the nematode *Caenorhabditis elegans* as a model animal for assessing the toxicity induced by microcystin-LR. *J. Environ. Sci.* **2009**, *21*, 395–401.
56. Cao, Q.; Steinman, A.D.; Yao, L.; Xie, L. Effects of light, microorganisms, farming chemicals and water content on the degradation of microcystin-LR in agricultural soils. *Ecotoxicol. Environ. Saf.* **2018**, *156*, 141–147. [\[CrossRef\]](#) [\[PubMed\]](#)
57. Machado, J.; Campos, A.; Vasconcelos, V.; Freitas, M. Effects of microcystin-LR and cylindrospermopsin on plant-soil systems: A review of their relevance for agricultural plant quality and public health. *Environ. Res.* **2017**, *153*, 191–204. [\[CrossRef\]](#)

58. Saqrane, S.; Ghazali, I.E.; Oudra, B.; Bouarab, L.; Vasconcelos, V. Effects of cyanobacteria producing microcystins on seed germination and seedling growth of several agricultural plants. *J. Environ. Sci. Health Part B* **2008**, *43*, 443–451. [[CrossRef](#)]
59. Mikalsen, B.; Bojson, G.; Skulberg, O.M.; Fastner, J.; Davies, W.; Gabrielsen, T.M.; Rudi, K.; Jakobsen, K.S. Natural variation in the microcystin synthetase operon mcyABC and impact on microcystin production in *Microcystis* strains. *J. Bacteriol.* **2003**, *185*, 2774–2785. [[CrossRef](#)] [[PubMed](#)]
60. Felnagle, E.A.; Jackson, E.E.; Chan, Y.A.; Podevels, A.M.; Berti, A.D.; McMahon, M.D.; Thomas, M.G. Nonribosomal peptide synthetases involved in the production of medically relevant natural products. *Mol. Pharm.* **2008**, *5*, 191–211. [[CrossRef](#)]
61. Bouaïcha, N.; Miles, C.O.; Beach, D.G.; Labidi, Z.; Djabri, A.; Benayache, N.Y.; Nguyen-Quang, T. Structural diversity, characterization and toxicology of microcystins. *Toxins* **2019**, *11*, 714. [[CrossRef](#)]
62. Fontanillo, M.; Köhn, M. Microcystins: Synthesis and structure–activity relationship studies toward PP1 and PP2A. *Bioorg. Med. Chem.* **2018**, *26*, 1118–1126. [[CrossRef](#)]
63. Craig, M.; Luu, H.A.; McCready, T.L.; Williams, D.; Andersen, R.J.; Holmes, C.F. Molecular mechanisms underlying the interaction of motuporin and microcystins with type-1 and type-2A protein phosphatases. *Biochem. Cell Biol.* **1996**, *74*, 569–578. [[CrossRef](#)] [[PubMed](#)]
64. Ferrão-Filho, A.D.S.; Kozłowski-Suzuki, B. Cyanotoxins: Bioaccumulation and effects on aquatic animals. *Mar. Drugs* **2011**, *9*, 2729–2772. [[CrossRef](#)]
65. Carmichael, W.W.; Boyer, G.L. Health impacts from cyanobacteria harmful algae blooms: Implications for the North American Great Lakes. *Harmful Algae* **2016**, *54*, 194–212. [[CrossRef](#)]
66. Toivola, D.M.; Omary, M.B.; Ku, N.O.; Peltola, O.; Baribault, H.; Eriksson, J.E. Protein phosphatase inhibition in normal and keratin 8/18 assembly-incompetent mouse strains supports a functional role of keratin intermediate filaments in preserving hepatocyte integrity. *Hepatology* **1998**, *28*, 116–128. [[CrossRef](#)] [[PubMed](#)]
67. Frangež, R.; Žužek, M.C.; Mrkun, J.; Šuput, D.; Sedmak, B.; Kosek, M. Microcystin-LR affects cytoskeleton and morphology of rabbit primary whole embryo cultured cells in vitro. *Toxicol.* **2003**, *41*, 999–1005. [[CrossRef](#)]
68. Pappas, D.; Gkelis, S.; Panteris, E. The effects of microcystin-LR in *Oryza sativa* root cells: F-actin as a new target of cyanobacterial toxicity. *Plant Biol.* **2020**, *22*, 839–849. [[CrossRef](#)] [[PubMed](#)]
69. Abe, T.; Lawson, T.; Weyers, J.D.B.; Codd, G.A. Microcystin-LR inhibits photosynthesis of *Phaseolus vulgaris* primary leaves: Implications for current spray irrigation practice. *New Phytol.* **1996**, *133*, 651–658. [[CrossRef](#)]
70. Llana-Ruiz-Cabello, M.; Jos, A.; Cameán, A.; Oliveira, F.; Barreiro, A.; Machado, J.; Azevedo, J.; Pinto, E.; Almeida, A.; Campos, A.; et al. Analysis of the use of cylindrospermopsin and/or microcystin-contaminated water in the growth, mineral content, and contamination of *Spinacia oleracea* and *Lactuca sativa*. *Toxins* **2019**, *11*, 624. [[CrossRef](#)]
71. Freitas, M.; Azevedo, J.; Pinto, E.; Neves, J.; Campos, A.; Vasconcelos, V. Effects of microcystin-LR, cylindrospermopsin and a microcystin-LR/cylindrospermopsin mixture on growth, oxidative stress and mineral content in lettuce plants (*Lactuca sativa* L.). *Ecotoxicol. Environ. Saf.* **2015**, *116*, 59–67. [[CrossRef](#)]
72. Machado, J.; Azevedo, J.; Freitas, M.; Pinto, E.; Almeida, A.; Vasconcelos, V.; Campos, A. Analysis of the use of microcystin-contaminated water in the growth and nutritional quality of the root-vegetable, *Daucus carota*. *Environ. Sci. Pollut. Res.* **2017**, *24*, 752–764. [[CrossRef](#)]
73. Zhang, Y.; Whalen, J.K.; Sauvé, S. Phytotoxicity and bioconcentration of microcystins in agricultural plants: Meta-analysis and risk assessment. *Environ. Pollut.* **2020**, *272*, 115966. [[CrossRef](#)]
74. Malaïssi, L.; Vaccarini, C.A.; Hernández, M.P.; Ruscitti, M.; Arango, C.; Busquets, F.; Arambarri, A.M.; Giannuzzi, L.; Andrinolo, D.; Sedan, D. [D-Leu1] MC-LR and MC-LR: A small–large difference: Significantly different effects on *Phaseolus vulgaris* L. (Fabaceae) growth and phototropic response after single contact during imbibition with each of these microcystin variants. *Toxins* **2020**, *12*, 585. [[CrossRef](#)]
75. Sedan, D.; Malaïssi, L.; Vaccarini, C.A.; Ventosi, E.; Laguens, M.; Rosso, L.; Giannuzzi, L.; Andrinolo, D. [D-Leu1] MC-LR has lower PP1 inhibitory capability and greater toxic potency than MC-LR in animal and plant tissues. *Toxins* **2020**, *12*, 632. [[CrossRef](#)] [[PubMed](#)]
76. Cao, Q.; Rediske, R.R.; Yao, L.; Xie, L. Effect of microcystins on root growth, oxidative response, and exudation of rice (*Oryza sativa*). *Ecotoxicol. Environ. Saf.* **2018**, *149*, 143–149. [[CrossRef](#)] [[PubMed](#)]
77. Do Carmo Bittencourt-Oliveira, M.; Cordeiro-Araújo, M.K.; Chia, M.A.; de Toledo Arruda-Neto, J.D.; de Oliveira, Ê.T.; dos Santos, F. Lettuce irrigated with contaminated water: Photosynthetic effects, antioxidative response and bioaccumulation of microcystin congeners. *Ecotoxicol. Environ. Saf.* **2016**, *128*, 83–90. [[CrossRef](#)] [[PubMed](#)]
78. Corbel, S.; Bouaïcha, N.; Nélieu, S.; Mougin, C. Soil irrigation with water and toxic cyanobacterial microcystins accelerates tomato development. *Environ. Chem. Lett.* **2015**, *13*, 447–452. [[CrossRef](#)]
79. Pflugmacher, S. Reduction in germination rate and elevation of peroxidase activity in *Zea mays* seedlings due to exposure to different microcystin analogues and toxic cell free cyanobacterial crude extract. *J. Appl. Bot. Food Qual.* **2007**, *81*, 45–48.
80. Pflugmacher, S.; Hofmann, J.; Hübner, B. Effects on growth and physiological parameters in wheat (*Triticum aestivum* L.) grown in soil and irrigated with cyanobacterial toxin contaminated water. *Environ. Toxicol. Chem.* **2007**, *26*, 2710–2716. [[CrossRef](#)]
81. Pflugmacher, S.; Aulhorn, M.; Grimm, B. Influence of a cyanobacterial crude extract containing microcystin-LR on the physiology and antioxidative defence systems of different spinach variants. *New Phytol.* **2007**, *175*, 482–489. [[CrossRef](#)]

82. Lee, S.; Jiang, X.; Manubolu, M.; Riedl, K.; Ludsin, S.A.; Martin, J.F.; Lee, J. Fresh produce and their soils accumulate cyanotoxins from irrigation water: Implications for public health and food security. *Food Res. Int.* **2017**, *102*, 234–245. [[CrossRef](#)]
83. Cevallos-Casals, B.A.; Cisneros-Zevallos, L. Impact of germination on phenolic content and antioxidant activity of 13 edible seed species. *Food Chem.* **2010**, *119*, 1485–1490. [[CrossRef](#)]
84. Pereira, S.; Saker, M.L.; Vale, M.; Vasconcelos, V.M. Comparison of sensitivity of grasses (*Lolium perenne* L. and *Festuca rubra* L.) and lettuce (*Lactuca sativa* L.) exposed to water contaminated with microcystins. *Bull. Environ. Contam. Toxicol.* **2009**, *83*, 81–84. [[CrossRef](#)]
85. Gutiérrez-Praena, D.; Campos, A.; Azevedo, J.; Neves, J.; Freitas, M.; Guzmán-Guillén, R.; Cameán, A.M.; Renaut, J.; Vasconcelos, V. Exposure of *Lycopersicon esculentum* to microcystin-LR: Effects in the leaf proteome and toxin translocation from water to leaves and fruits. *Toxins* **2014**, *6*, 1837–1854. [[CrossRef](#)] [[PubMed](#)]
86. Romero-Oliva, C.S.; Contardo-Jara, V.; Pflugmacher, S. Time dependent uptake, bioaccumulation and biotransformation of cell free crude extract microcystins from Lake Amatitlán, Guatemala by *Ceratophyllum demersum*, *Egeria densa* and *Hydrilla verticillata*. *Toxicon* **2015**, *105*, 62–73. [[CrossRef](#)]
87. Verbitsky, V.B.; Kurbatova, S.A.; Berezina, N.A.; Korneva, L.G.; Meteleva, N.Y.; Makarova, O.S.; Sharov, A.N.; Ershov, I.Y.; Malysheva, O.A.; Russkikh, Y.V.; et al. Responses of aquatic organisms to cyanobacteria and Elodea in microcosms. *Dokl Biol. Sci.* **2019**, *488*, 136–140. [[CrossRef](#)] [[PubMed](#)]
88. Mitrovic, S.M.; Allis, O.; Furey, A.; James, K.J. Bioaccumulation and harmful effects of microcystin-LR in the aquatic plants *Lemna minor* and *Wolffia arrhiza* and the filamentous alga *Chladophora fracta*. *Ecotoxicol. Environ. Saf.* **2005**, *61*, 345–352. [[CrossRef](#)]
89. Zanchett, G.; Oliveira-Filho, E.C. Cyanobacteria and cyanotoxins: From impacts on aquatic ecosystems and human health to anticarcinogenic effects. *Toxins* **2013**, *5*, 1896–1917. [[CrossRef](#)]
90. Saqrane, S.; Ouahid, Y.; El Ghazali, I.; Oudra, B.; Bouarab, L.; del Campo, F.F. Physiological changes in *Triticum durum*, *Zea mays*, *Pisum sativum* and *Lens esculenta* cultivars, caused by irrigation with water contaminated with microcystins: A laboratory experimental approach. *Toxicon* **2009**, *53*, 786–796. [[CrossRef](#)] [[PubMed](#)]
91. Saqrane, S.; Ghazali, I.E.; Ouahid, Y.; Hassni, M.E.; Hadrami, I.E.; Bouarab, L.; del Campo, F.F.; Oudra, B.; Vasconcelos, V. Phytotoxic effects of cyanobacteria extract on the aquatic plant *Lemna gibba*: Microcystin accumulation, detoxication and oxidative stress induction. *Aquat. Toxicol.* **2007**, *83*, 284–294. [[CrossRef](#)]
92. Melaram, R. Environmental risk factors implicated in liver disease: A mini-review. *Front. Public Health* **2021**, *9*, 683719. [[CrossRef](#)]