


# Clinical Course from Diagnosis to Death in Patients with Well-Differentiated Thyroid Cancer

Hyunju Park, Jun Park, So Young Park, Tae Hyuk Kim, Sun Wook Kim, and Jae Hoon Chung


**Table S1.** Clinicopathological characteristics of DTC patients based on time interval.

Characteristics		1996-2002	2003-2009	2010-2018	P-value
<b>Number of Patients (Inoperable Patients)</b>		27 (2)	36 (3)	16 (1)	
Mean survival (month, SD)		74.37 (9.39)	56.25 (5.23)	63.19 (13.5)	
Sex ( <i>n</i> , %)	Female	19 (70.4)	26 (72.2)	7 (43.8)	0.113
Primary tumor size ( <i>n</i> , %)	<4cm	19 (70.4)	28 (77.8)	12 (75.0)	0.799
	4cm or more	8 (29.6)	8 (22.2)	4 (25.0)	
	1	4 (16.0)	4 (12.1)	1 (6.7)	
pT ( <i>n</i> , %)	2	5 (20.0)	3 (9.1)	1 (6.7)	0.419
	3	11 (44.0)	11 (33.3)	8 (53.3)	
	4	5 (20.0)	15 (45.5)	5 (33.3)	
pN ( <i>n</i> , %)	Not assessed	1 (4.0)	1 (3.0)	1 (6.7)	0.251
	N0	9 (36.0)	17 (51.5)	6 (40.0)	
	N1a	8 (32.0)	3 (9.1)	1 (6.7)	
	N1b	7 (28.0)	12 (36.4)	7 (46.7)	

SD, standard deviation; *n*, number of patients.



**Figure S1.** Overall survival based on time period.



**Figure S2.** Timelines from diagnosis to death in each group. In late-DM, first column represented time from diagnosis to distant metastasis, and second column represented time from distant metastasis to death (median, interquartile range).



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).