Supplementary Material: Extracellular Neuroglobin as a Stress-Induced Factor Activating Pre-Adaptation Mechanisms Against Oxidative Stress and Chemotherapy-Induced Cell Death in Breast Cancer

Marco Fiocchetti, Virginia Solar Fernandez, Marco Segatto, Stefano Leone, Paolo Cercola, Annalisa Massari, Francesco Cavaliere and Maria Marino





Figure S1. Graphic representation of experimental protocol for the generation of conditioned media (for details see Section 4).






Figure S2. Whole blots images from Figure 2.




Figure 3. Whole blots images from Figure 3B,C.





Figure S4. Whole blots images from Figure 4B,D,F.



95

94

111

111

87

15

9

19

DTX+E2

CM-Veh 4h +DTX

CM-E2 4h +DTX

CM-H2O2 4h +DTX




Figure 5D arbitrary units		
	PARP1 89 KDa	TUB
Veh	1	46
E2	2	52
CM-Veh 24h	1	28
CM-E2 24h	3	31
CM-H2O2 24h	2	40
DTX	17	68
DTX+E2	7	55
CM-Veh 24h +DTX	22	53
CM-E2 24h +DTX	28	71
CM-H2O2 24h +DTX	5	27

Figure 5C arbitrary units		
	Bcl-2 26 kDa	TUB
Veh	17	36
E2	46	46
CM-Veh 24h	11	36
CM-E2 24h	36	40
CM-H2O2 24h	64	59

Figure S5. Whole blots images from Figure 5C,D,G.