



Supplementary Materials: TNF α Enhances Tamoxifen Sensitivity through Dissociation of ER α -p53-NCOR1 Complexes in ER α -Positive Breast Cancer

Hyunhee Kim, Seung-Ho Park, Jangho Lee, Gi-Jun Sung, Ji-Hye Song, Sungmin Kwak, Ji-Hoon Jeong, Min-Jeong Kong, Jin-Taek Hwang, Hyo-Kyoung Choi and Kyung-Chul Choi




Figure S1. Analysis of the association between NCOR1 and P53 in ER+ breast cancer. The association between NCOR1 and P53 signaling in ER+ breast cancer. The transcriptome data of breast cancer cohort from The Cancer Genome Atlas (TCGA) ($n = 1,904$) was analyzed. (A, B) *Ncor1* mRNA levels in ER positive or ER negative breast cancer. ER status was measured by RNA-Seq (A) or IHC (B). (C, D) The correlation between *Ncor1* and *Tp53* mRNA expressions in ER positive (E) or ER negative breast cancer (E). Venn diagram for the biological processes enriched in ER+ and/or ER- breast tumors. (F) The biological processes enriched only in ER+ breast tumor.

Table S1. Antibodies used for WB and ChIP assays.

Antibodies	Vendor Cat no.	Local	Country	Titer	Usage
α H3K9ac	Merck 06-942	Darmstadt	Germany	(1:100)	
α H3K4me3	Merck 05-1339	Darmstadt	Germany	(1:100)	
α H3K9me3	Merck 05-1250	Darmstadt	Germany	(1:100)	W. B (ChIP)
α ER α	Santa Cruz Biotechnol- ogy SC-543	Dallas, Texas	USA	1:1000 (1:100)	
α p53	Santa Cruz Biotechnol- ogy SC-126	Dallas, Texas	USA	1:1000 (1:100)	
α p-p53	Abcam ab134190	Cambridge	UK	1:1000	
α NCoR	ATGEN	Seongnam, Gyeonggido	Korea	1:1000	
α p-NCoR	ATGEN	Seongnam, Gyeonggido	Korea	1:1000	
PARP-1	Cell signaling 9542S	Danvers, MA	USA	1:1000	
Bax	Cell signaling 5023S	Danvers, MA	USA	1:1000	W.B
Bcl-2	Santa Cruz Biotechnol- ogy SC-492	Dallas, Texas	USA	1:1000	
α β -Actin	Sigma-Aldrich A5441	Saint Louis, MO	USA	1:10000	

Abbreviation: α H3K9ac, anti-H3K9 acetylation; α H3K4me3, anti-H3K4 trimethylation; α H3K9me3, anti-H3K9 trimethylation; α ER α , anti-Estrogen Receptor α ; α p53, anti-p53; α p-p53, anti-phospho-p53; α NCoR, anti-Nuclear receptor co-repressor; α -PARP-1, anti-PARP-1; α Bax, anti-Bax α Bcl-2, anti-Bcl-2; α β -Actin, anti-beta Actin; W.B, Western blotting; ChIP, Chromatin immunoprecipitation.

Table S2. Primers used for qRT-PCR and ChIP assays.

Primer List			
NCoR1	Forward	5' ACACCGCAGTATTGTCCAAAT 3'	qRT-PCR for human
	Reverse	5' CACCTGGTTGTCTTGATGTTCT 3'	
GAPDH	Forward	5' TGATGACATCAAGAAGGTGGTGAAG 3'	qPCR for ChIP assay
	Reverse	5' TCCTTGGAGGCCATGTAGGCCAT 3'	
p21 pro- moter	Forward	5' GTAAATCCTTGCCTGCCAGAG 3'	qPCR for ChIP assay
	Reverse	5' GCGCTGGACACATTCCCCA 3'	

Abbreviation: NCoR1, Nuclear receptor co-repressor1; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; p21, cyclin-dependent kinase inhibitor 1; ChIP assay, chromatin immunoprecipitation assay.

Table S3. NCOR1 shRNA sequences for knockdown system.

shRNA sequences list	
<i>shNCOR1 #1</i>	5' CCGGGCCCACAGATGATGAAGAAATCTGAGATTCTT CATCATCTGTGGGCTTTTG 3'
<i>shNCOR1 #2</i>	5' CCGGGCCATCAAACACAATGTCAAACACTCGAGTTGACA TTGTGTTGATGGCTTTTG 3'

Abbreviation: NCoR1, Nuclear receptor co-repressor1.