

Table S1. Improved compliance and time to postoperative mobilization with the recommendation after the mobilization recommendation was introduced. Fisher's exact tests were used for *p*-values, except for mobilization (*Mann Whitney Test).

	All cases <i>n</i> =445	Before <i>n</i> =164	After <i>n</i> =281	Odds ratio (95% CI)	<i>p</i> -value
Compliance					
Yes	267 (60)	88 (54)	179 (64)	1.52 (1.00-2.28)	0.045
No	178 (40)	76 (46)	102 (36)		
Mobilized within 24 hours					
Yes	196 (44)	17 (10)	179 (64)	15.2 (8.5-28.4)	<0.001
No	249 (56)	147 (90)	102 (36)		
Mobilized within 48 hours					
Yes	342 (77)	88 (54)	254 (90)	8.1 (4.8-13.9)	<0.001
No	103 (23)	76 (46)	27 (10)		
Mobilized within 72 hours					
Yes	405 (91)	136 (83)	269 (96)	4.6 (2.2-10.3)	<0.001
No	40 (9)	28 (17)	12 (4)		
Time to mobilization					
Mean ± SD	2.2 ± 2.5	2.9 ± 2.0	1.7 ± 2.6		
Median (IQR)	2 (1-2)	2 (2-3)	1 (1-2)	<i>z</i> = 11.48	<0.001*

Table S2. Logistic regression analysis (controlling for alcohol) of compliance with guideline concordant care, postoperative complications, and length of stay. The reference level for odds ratios is indicated with *italics*.

Characteristic	All cases	Compliance=Yes	Compliance=No	OR	95% CI	p-value
<i>n (%)</i>	445 (100)	267 (60)	178 (40)			
Major complication						
Yes	70 (15.8)	29 (10.9)	39 (21.9)	0.44	0.25-0.76	0.004
No	374 (84.2)	238 (89.1)	123 (69.1)			
Pneumonia						
Yes	37 (8.3)	15 (5.6)	22 ()	0.35	0.16-0.75	0.007
No	408 (91.7)	252 (94.4)	156 ()			
Length of stay						
<i>POD 0-10</i>	204 (46)	131 (49.1)	109 (61.6)	0.58	0.38-0.59	0.013
After POD 10	240 (54)	136 (50.9)	68 (38.4)			

Table S3. Predictors of major complications using elastic net regularization followed by multivariable logistic regression. Mobilization (within 48 hours) substituted for compliance with guideline-recommended mobilization.

Characteristic (reference level)	Major complications			
	Odds ratio	95% CI	z	p-value
Age (centred at the mean)	1.01	0.98-1.04	0.56	0.558
BMI category (<i>healthy range</i>)				
Underweight	1.93	0.57-6.54	1.05	0.293
Overweight	0.44	0.19-1.02	-1.91	0.056
Obese	0.74	0.32-1.73	-0.69	0.493
Sex (<i>male</i>)				
Female	0.70	0.33-1.50	-0.91	0.362
Smoking Status (<i>never</i>)				
Former smoker	0.57	0.24-1.35	-1.28	0.201
Current smoker	0.23	0.08-0.65	-2.74	0.006
Alcohol status (<i>never</i>)				
Light/Moderate	1.86	0.75-4.61	-1.34	0.181
Heavy	1.32	0.40-4.32	0.46	0.645
Former drinker	0.52	0.12-2.27	-0.87	0.383
Comorbidity (<i>none</i>)				
One	1.41	0.57-3.49	0.74	0.460
Two or more	2.64	1.14-6.15	2.25	0.024
Primary Site (<i>oral cavity</i>)				
Pharynx & Larynx	1.42	0.49-4.07	0.65	0.516
Paranasal/Nasal	0.31	0.05-1.81	-1.30	0.192
Skin	0.17	0.01-1.90	-1.44	0.150
Other	0.32	0.05-1.92	-1.25	0.212
Cancer stage (<i>I-II</i>)				
III-IV	2.29	0.96-5.48	1.86	0.063
Flap type (<i>radial forearm</i>)				
Fibula	1.28	0.52-3.14	0.53	0.597
Anterolateral thigh	1.39	0.45-4.29	0.57	0.566
Other	2.80	1.12-7.01	2.20	0.028
Unit arrival (<i>POD 0-1</i>)				
After POD 1	2.62	1.17-5.87	2.33	0.020
Compliance (<i>no</i>)				
Yes	0.57	0.29-1.13	-1.62	0.106
Tracheostomy (<i>no</i>)				
Yes	0.38	1.13-1.07	-1.83	0.067

Table S4. Predictors of hospital length of stay using elastic net regularization followed by multivariable logistic regression. Mobilization (within 48 hours) substituted for compliance with guideline-recommended mobilization.

Characteristic (reference level)	Major complications			
	Odds ratio	95% CI	z	p-value
Age (centred at the mean)	1.03	1.01-1.05	2.43	0.015
BMI category (healthy range)				
Underweight	1.00	0.34-2.96	-0.01	0.994
Overweight	0.64	0.34-1.20	-1.39	0.165
Obese	0.48	0.23-0.98	-2.02	0.043
Alcohol status (never)				
Light/Moderate	0.58	0.29-1.17	-1.51	0.130
Heavy	0.80	0.35-1.78	-0.56	0.577
Former drinker	2.37	0.90-6.21	1.75	0.081
Comorbidity (none)				
One	1.00	0.51-1.96	0.00	0.998
Two or more	1.50	0.75-3.01	1.14	0.253
Primary Site (oral cavity)				
Pharynx & Larynx	4.69	1.65-13.35	2.90	0.004
Paranasal/Nasal	0.06	0.01-0.33	-3.26	0.001
Skin	0.12	0.02-0.54	-2.75	0.006
Other	0.73	0.19-2.76	-0.46	0.646
Flap count (one)				
Two	0.21	0.01-6.44	-0.89	0.375
Resection extent (soft tissue only)				
Bone	1.15	0.60-2.20	0.43	0.669
Soft tissue & bone	27.10	0.49-1493.71	1.61	0.107
Compliance (no)				
Yes	0.89	0.51-1.55	-0.41	0.680
Tracheostomy (no)				
Yes	3.03	1.21-7.60	2.36	0.018
Any complication (no)				
Yes	6.69	3.75-11.95	6.42	<0.001
Major Complication (no)				
Yes	2.29	0.91-5.76	1.76	0.079