

Supplementary materials: Validation of the T Descriptor (TNM-8) in T3N0 Non-Small-Cell Lung Cancer Patients; a Bi-centric Cohort Analysis with Arguments for Redefinition

Philip Baum, Samantha Taber, Stella Erdmann, Thomas Muley, Mark Kriegsmann, Petros Christopoulos, Michael Thomas, Hauke Winter, Joachim Pfannschmidt and Martin E. Eichhorn

Table S1. Differences between Patients with chemotherapy vs. no chemotherapy. Patients with adjuvant Radio-Chemotherapy/ Radiotherapy are not included in the analysis.

	No chemotherapy		chemotherapy		p value
	No.	%	No.	%	
All	109	40.7	159	59.3	
Gender					<i>0.940</i>
Male	66	60.6	97	61.0	
Female	43	39.5	62	39.0	
ECOG					<i><0.001</i>
0	75	68.8	147	92.5	
≥1	34	30.3	12	7.6	
Age					<i>0.155</i>
<60	8	7.3	24	15.1	
60-69	25	22.9	32	20.1	
≥70	76	69.7	103	64.8	
Surgery					<i>0.749</i>
Lobectomy	89	81.7	135	84.9	
Sleeve Resection	3	2.8	3	1.9	
Sublobar Resection	7	6.4	6	3.8	
Bilobectomy	3	2.8	7	4.4	
Pneumonectomy	7	6.4	8	5.0	
Resection side*					<i>0.310</i>
Right	68	62.4	91	57.2	
Left	40	36.7	68	42.8	
Residue					<i>0.190</i>
0	105	96.3	157	98.7	
1	4	3.7	2	1.3	
Histology					<i>0.586</i>
AdC	49	45.0	77	48.4	
SqCC	47	43.1	59	37.1	
Other	13	11.9	23	14.5	

*one Patient with missing side.