



# **Clinical Landscape of PARP Inhibitors in Ovarian Cancer: Molecular Mechanisms and Clues to Overcome Resistance**

Satoru Kyo<sup>1,\*</sup>, Kosuke Kanno<sup>1</sup>, Masahiro Takakura<sup>2</sup>, Hitomi Yamashita<sup>1</sup>, Masako Ishikawa<sup>1</sup>, Tomoka Ishibashi<sup>1</sup>, Seiya Sato<sup>1</sup> and Kentaro Nakayama<sup>1</sup>

- <sup>1</sup> Department of Obstetrics and Gynecology, Shimane University Faculty of Medicine, Izumo 693-8501, Japan; kanno39@med.shimane-u.ac.jp (K.K.); meme1103@med.shimane-u.ac.jp (H.Y.); m-ishi@med.shimane-u.ac.jp (M.I.); tomoka@med.shimane-u.ac.jp (T.I.); sato\_seiya9534@yahoo.co.jp (S.S.); kn88@med.shimane-u.ac.jp (K.N.)
- <sup>2</sup> Department of Obstetrics and Gynecology, Kanazawa Medical University, Ishikawa 920-0293, Japan; takakura@kanazawa-med.ac.jp
- \* Correspondence: satoruky@med.shimane-u.ac.jp; Fax: +81-(0)853-20-2264

**Simple Summary:** Recent development of maintenance therapy using PARP inhibitors in ovarian cancer has led to a significant improvement in survival rates. However, resistance to these inhibitors can occur in patients, causing disease progression or relapse. Consequently, novel treatment strategies are urgently needed to overcome this resistance. This review article focuses on the precise molecular mechanisms by which PARP inhibitors exert their antitumor effects, as well as how they elicit resistance, in order to gain insight into novel therapeutic approaches to overcome PARP inhibitor resistance in ovarian cancer.

Abstract: The survival of patients with advanced or recurrent ovarian cancer has improved tremendously in the past decade, mainly due to the establishment of maintenance therapy with poly (ADP-ribose) polymerase (PARP) inhibitors (PARPis) after conservative chemotherapies. Despite their superior efficacy, resistance to PARPis has been reported, and patients with resistance have a much worse prognosis. Therefore, the development of novel treatment strategies to overcome PARPi resistance is urgently needed. The present review article focuses on the molecular mechanisms of how PARPis exert cytotoxic effects on cancer cells through DNA repair processes, especially the genetic background and tumor microenvironment favored by PARPis. Furthermore, currently available information on PARPi resistance mechanisms is introduced and discussed to develop a novel therapeutic approach against them.

**Keywords:** ovarian cancer; PARP inhibitor; maintenance therapy; homologous recombination deficiency; replication fork; DNA damage response; cGAS/STING; cytosolic immunity

## 1. Introduction

The concept of maintenance therapy after platinum-based chemotherapy has rapidly developed in the first-line setting of advanced epithelial ovarian, tubal, and peritoneal cancers (collectively referred to as ovarian cancer) or after recurrence, in which poly (ADP-ribose) polymerase inhibitors (PARPis) hold important roles, providing prolonged progression-free survival (PFS) and overall survival (OS) compared to conservative chemotherapy alone [1–6]. Nevertheless, a considerable proportion of patients still show disease progression or recurrence after maintenance therapies, and the prognosis is much worse, although several approaches, including repeated PARPi treatment, have been attempted to treat such patients [7]. To predict or protect against disease progression or the recurrence of ovarian cancers after maintenance therapy, the key is to develop a profound understanding of the molecular mechanisms by which PARPis exert specific anti-tumor effects, distinct from those of conservative chemotherapy. The present review article focuses on those



Citation: Kyo, S.; Kanno, K.; Takakura, M.; Yamashita, H.; Ishikawa, M.; Ishibashi, T.; Sato, S.; Nakayama, K. Clinical Landscape of PARP Inhibitors in Ovarian Cancer: Molecular Mechanisms and Clues to Overcome Resistance. *Cancers* **2022**, *14*, 2504. https://doi.org/10.3390/ cancers14102504

Academic Editors: Christina M. Annunziata and Adam R. Karpf

Received: 26 April 2022 Accepted: 18 May 2022 Published: 19 May 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). mechanisms, mentioning the genetic background and tumor microenvironment (TME) of ovarian cancers before, during, and after PARPi treatment. The importance of combination treatment strategies based on an understanding of complex molecular rationales is also stressed. Finally, methods to overcome PARPi resistance are discussed based on the currently available information on the detailed molecular mechanisms that cause resistance.

### 2. Two Distinct Therapeutic Mechanisms of PARPis

## 2.1. Enzymatic Inhibition

As a family member of enzymes composed of various isoforms, including PARP1/2 [8], PARP plays am essential role in the base excision repair (BER) of single-strand DNA breaks (SSBs). The founding family member, PARP1, has been the most extensively studied. It binds to the sites of SSBs via the zinc-finger DNA binding domains and catalyzes poly-ADPribosylation (PARylation) to proteins that repair SSBs using their enzymatic activity [9,10]. Poly-ADP-ribose (PAR) has a negative charge and serves as a platform to recruit proteins that repair SSBs with a positive charge, resulting in the efficient repair of SSBs. It is a major principle that PARP1 promotes repair of SSBs as an enzyme.

The enzymatic activity of PARP1 is targeted by PARPis, inhibiting PARylation. Eventually, PARPis disturb repair of SSBs, leading to the accumulation of double-strand DNA breaks (DSBs) via subsequent DNA replications. Such DSBs are repaired mainly by specific repair pathways, named homologous recombination repair (HR), a mechanism of DSB repair using sister chromatids as a template [11], although there is another machinery of DSB repair, non-homologous end-joining (NHEJ), that is used when HR is decreased. When cancer cells have HR defects (homologous recombination defect, HRD), they suffer from accumulated DSBs leading to increased genetic and chromosomal instability, finally resulting in cell death. This is mechanistic rationale, by which PARPis exert cytotoxic activity, especially in the presence of HRD, is called synthetic lethality [12,13].

## 2.2. PARP Trapping

Furthermore, PARPis exert an alternative effect on PARP1. Of note, PARP1 induces PARylation not only to proteins that repair SSBs, but also to PARP1 itself. The PARylation of PARP1 itself induces the release from DNA lesions due to charge repulsion, because of the negative charge of DNA. Thus, PARP1 leaves DNA lesions after PARylation by itself. Subsequently, PARPis inhibit the PARylation of PARP1 and, therefore, prevent PARP1 from leaving DNA lesions, allowing PARP1 to be continuously trapped on the DNA (PARP-trapping) [14]. This PARP-DNA complex interferes with DNA replication by stalling the replication fork and thereby functions as a so-called "DNA poison", which confers collapse of the replication fork and cell death [13,15]. This is an alternative mechanism by which PARPis exert cytotoxic activity. Various repair mechanisms, including HR, are mobilized to eliminate such poisonous complexes. However, the defects of repair mechanisms, such as HRD, allow continuous PARP-DNA complexes, promoting defective DNA replication and cell death.

Thus, two distinct types of PARPi actions, (1) enzymatic inhibition for synthetic lethality and/or (2) PARP trapping for fork collapse and cell death, occur in the context of maintenance therapy. It is important to determine which type of mechanism is clinically more active for cytotoxic activity. In previous in vitro studies [14,16], cancer cells were treated with various concentrations of PARPis, and the statuses of PARylation and PARP trapping were quantified and compared with cytotoxic efficacy. Overall, PARP trapping required higher concentrations of PARPis than enzymatic inhibition in most cell types, and cytotoxic activity was more dependent on PARP trapping than on enzymatic inhibition for PARylation; most cell types exhibited resistance to PARPis in lower concentrations that were sufficient for complete inhibition of PARylation, but insufficient for PARP trapping [14,16]. Therefore, PARP trapping appears to play predominant roles in cytotoxicity, especially at higher concentrations.

## 3. HRD as a Predictive Marker for the Efficacy and Survival Benefit of PARPis

The HR repair mechanism (HRR) involves multiple proteins, and germline *BRCA1/2* mutations are the most well-known genetic component of HRD in ovarian cancer, accounting for 10–15% of cases, especially for high-grade serous carcinoma (HGSOC), in which they account for 20–30% of patients [17–19]. Other causes of HRD include somatic *BRCA1/2* mutations, accounting for 1/3–1/2 of the germline *BRCA1/2* mutations [20], promoter silencing of *BRCA1* (approximately 10% of HGSOC) [17], or mutations of other genes involved in HRR. In the presence of HRD, PARPis are expected to show greater cytotoxic effects via enhanced synthetic lethality and PARP trapping.

Recent clinical trials with PARPis showed that the benefit with the highest clinical relevance was observed in BRCA-mutated or HRD cohorts [1–4], and, therefore, BRCAmutation and HRD status have been used clinically as companions to predict the efficacy of PARPis. The myChoice<sup>®</sup> CDx is the first and only FDA-approved tumor test that detects BRCA1/2 variants and/or scores HRD status [21]. The scoring is determined by assessing genomic instability with the following critical biomarkers: loss of heterozygosity (LOH), telomeric allelic imbalance, and large-scale state (defined as chromosomal breakage that generates 10 Mb or larger DNA fragments), based on the concept that impaired HR causes accumulation of DSBs and induces genetic and chromosomal instability. The sum of the above three biomarkers was calculated to represent genomic instability status (GIS) or an HRD score and used as a surrogate measure for genetic instability, in which a threshold of  $\geq$  42 or *BRCA* mutation was considered HRD-positive. Using this system and other candidate biomarkers, including mutations in other HRR genes (including BR IP1, CDK12, RAD54L, RAD51B, ATM, FANCA, FANCD2, FANCL, RAD51C, RAD52, and XRCC) and BRCA1 promoter methylation, tumor samples from olaparib maintenance monotherapy (study 19 [22]) were analyzed [23]. Most patients with BRCA mutations, and almost all patients with BRCA1 methylation, were identified as myriad HRD-positive, while less than half of patients with mutations in other HRR genes were identified as HRDpositive [23]. Patients with myriad BRCA-mutated tumors exhibited the greatest survival benefit from olaparib (progression-free survival, or PFS, hazard ratio, or HR, of 0.17, with a 95% confidence interval, or CI, of 0.09–0.30), while those with HRD-positive status showed a comparable survival effect (HR 0.24, 95% CI 0.15–0.39), but patients with BRCA-wild type/HRD-positive tumors did not have as high a survival benefit (PFS HR 0.48, 95%) CI 0.18–1.27) [23]. Of note is that patients with mutations in other HRR genes identified by gene sequencing had an unexpectedly better survival benefit (PFS HR 0.21, 95% CI 0.04–0.86) [23]. Patients with wild-type BRCA/HRD-negative status (PFS HR 0.60, 95% CI 0.31–1.17) or those with wild-type BRCA/HRR genes (PFS HR 0.71, 95% CI 0.37–1.35) showed the least benefit from olaparib [23].

Thus, *BRCA* mutations and/or HRD-positive status are superior biomarkers of survival benefit from olaparib, on which *BRCA* mutations have the greatest impact. Mutations in HRR genes other than *BRCA* may also have treatment benefit from olaparib. However, there were contradictory results in a recent study [24] analyzing the patient population of the phase III PAOLA-1 study [25], in which mutations in HRR genes were not predictive of PFS benefit from olaparib in combination with bevacizumab, compared with bevacizumab alone. Further analyses with larger patient populations are needed to determine the clinical significance of detecting HRR gene mutations other than of *BRCA* as biomarkers predictive of treatment benefit of not only olaparib, but also other PARPis.

## 4. Dynamics of HRD Status in the Treatment of Ovarian Cancer

The status of HRD is vigorously altered during the treatment of ovarian cancer.

Takaya et al. [26] examined the clonality index and LOH scores using tumor samples of HGSOC obtained at primary debulking surgery (PDS) and interval debulking surgery (IDS) after neoadjuvant chemotherapy (NACT) or secondary debulking surgery (SDS) for recurrent tumors, and found that the clonality index and LOH decreased at IDS compared to before NACT, and then subsequently re-elevated at SDS. These findings indicate that intratumoral heterogeneity decreases after chemotherapy, probably because platinumsensitive clones disappeared due to the platinum-based chemotherapy, leaving selectively grown chemo-resistant clones (Figure 1).



**Figure 1.** Changes in HRD status during the treatment of ovarian cancer. Even though primary tumors are predominantly composed of HRD-positive cells, they are likely to be eradicated by platinum-based chemotherapy, and residual tumors may be occupied by HRD-negative cells, representing the HRP phenotype. However, after a platinum-free period, remnant HRD-positive cells that have a growth advantage can form recurrent tumors, exhibiting the HRD phenotype. These acronyms are defined as follows: HRD, homologous recombination deficiency; HRP, homologous recombination proficiency; IDS, interval debulking surgery; LOH, loss of heterozygosity; NACT, neoadjuvant chemotherapy; SDS, secondary debulking surgery.

Since the LOH score is a component of HRD status, these findings indicate that HRDpositive cells in the primary tumors were likely to be eradicated by chemotherapy, whereas HRD-negative cells were selected and occupied residual tumors. Subsequent re-elevation of the clonality index and LOH score suggests that recurrent tumors are composed of heterogeneous clones, including HRD-positive cells, reminiscent of their aggressive regrowth potential during platinum-free periods. Sokolenko et al. examined the LOH of the BRCA1 allele of HGSOC with germline BRCA1 mutation (gBRCA1 MT) in primary tumors in pre-NACT status, residual tumor at IDS, and recurrent tumor at SDS [27], and they found that all primary tumors had BRCA1 LOH, whereas retention of the wild-type BRCA1 allele was detected at IDS. Thereafter, most recurrent tumors exhibited BRCA1 LOH again, indicating that NACT results in the rapid selection of pre-existing wild-type BRCA1proficient cells, whereas tumor relapses re-acquired *BRCA1* LOH during therapy holidays. A similar observation was reported by Patel et al. [28], in which change in HRD status was measured and compared between paired primary and recurrent HGSOC samples, demonstrating that recurrent tumors took over the HRD status of the primary tumors, even after the successful treatment of primary lesions. Thus, maintenance therapy after the primary treatment of HRD-positive tumors needs to target HRD-positive cells, despite tumor samples at IDS exhibiting an HRD-negative phenotype. Therefore, HRD-negative status at IDS does not precisely predict HRD status in recurrent tumors and should not be considered in the choice of PARPis after IDS.

## **5.** The Fate of PARP Trapping: Fork Stabilization or Degradation That Determines the Efficacy of PARPis

As described earlier, PARP trapping on DNA by PARPis blocks DNA replication via stalled replication forks, triggering a series of replication stress responses [29]. Unless stalled forks are successfully repaired by such stress responses, they are extensively degraded by nuclease proteins and undergo irreversible fork collapse, leading to genomic instability and cell death [30,31].



A series of stress responses triggered by stalled replication forks are illustrated in Figure 2.

Figure 2. Integrity of fork stabilization on PARP trapping determines PARPi sensitivity. PARP trapping induces stalled replication forks, triggering a series of replication stress responses. After fork stalling, ssDNAs are generated in both lagging and leading strands, and are coated by RPA. The ssDNA-RPA complex then induces activation of the replication checkpoint via ATR and subsequent CHK1 and Wee1 activation, arresting the cell cycle to allow the time necessary to relieve the stalled fork and protect against fork collapse. RPA is then replaced by RAD51, which promotes fork reversal, a phenomenon to bypass and unwind the stalled fork, and the BRCA2-RAD51 axis plays pivotal roles in stabilizing fork reversal. When BRCA1/2 is defective, MRE11 and/or EXO1 nucleases destabilize stalled forks, resulting in accumulated DSBs or fork collapse, leading to PARPi sensitivity. Spartan (SPRTN) facilitates the repair of bulky PARP1-DNA complexes via proteolytic digestion to ensure fork progression, conferring PARPi resistance. Thus, the integrity of fork stabilization on PARP trapping plays major roles in determining sensitivity to PARPis, and nuclear factors that are involved in this integrity can be potential targets to overcome PARPi resistance. These acronyms are defined as follows: ATM, ataxia-telangiectasia mutated protein kinase; ATR, ataxia-telangiectasia and rad3 related protein kinase; CHK1, checkpoint kinase 1; DSB, double strand break; EXO1, exonuclease 1; HRD, homologous recombination deficiency; HRP, homologous recombination proficiency; MRE11, meiotic recombination 11; MUS81, methyl methanesulfonate and ultraviolet-sensitive gene; POLD3, 3rd subunit of DNA polymerase δ; RADX, RPA-related RAD51-antagonist on X-chromosome; RECQ1, ATP-dependent DNA helicase Q1; RPA, replication protein A; SMARCAL1, SWI/SNF-related, matrixassociated, actin-dependent regulator of chromatin subfamily A-like 1; Wee1, Wee1 tyrosine kinase.

A stalled replication fork is very unstable and is exposed to the risk of fork collapse. There are rescue processes for stalled replication forks [32]. After fork stalling, ssDNAs are generated by polymerase-helicase uncoupling in both lagging and leading strands. These ssDNA regions are coated by replication protein A (RPA) to prevent the formation of secondary structure [33]. Thereafter, the ssDNA-RPA complex activates the replication checkpoint via ataxia-telangiectasia-mutated protein kinase (ATR) and the subsequent CHK1 and Wee1 activation [32,34], which arrests the cell cycle to allow the time necessary to relieve the stalled fork and protect against fork collapse, and, therefore, composes a part of DNA damage responses (DDRs).

Subsequently, RPA was replaced by RAD51, which promotes fork reversal, a phenomenon to bypass and unwind the stalled fork [35]. This reversal is achieved by nascent lagging strands caused by blocked leading strands, which serve as an alternative template for leading strand DNA synthesis toward the opposite direction of fork progression, which has protective and stabilizing effects on stalled forks. It is important to consider why this reversal phenomenon functions to stabilize stalled forks. Fork progression across template DNA lesions, such as SSBs, induces RecQ1-dependent DSB generation as the fork progresses, which finally results in fork collapse [36]. The reversal of the fork protects it from colliding with the DNA lesions, thus avoiding DSB-induced fork collapse. Furthermore, fork reversal towards the opposite direction to replication impediments may allow additional time and room sufficient for the repair machineries to remove them [37].

Stalled replication forks are characterized as exposed DNA ends, which render them susceptible to a variety of nucleases including MRE11 and EXO1 [38]. These nuclease activities destabilize not only stalled forks, but also reversal forks, thereby threatening fork collapse. The BRCA2-RAD51 axis plays pivotal roles in preventing such deleterious fork degradation [39]. As a result, BRCA2 is likely to localize to the stalled forks and promote the formation of stable RAD51 nucleoprotein filaments, therefore inhibiting MRE11-medicated fork degradation [40], in which BRCA1 also contributes to fork stabilization. Whereas several RAD51 paralogs, including XRCC2 and XRCC3, are also required for preventing MRE11-medicated fork degradation [41], various factors including RADX counteract with RAD51 nucleoprotein filaments and, thereby, permit degradation of the fork by MRE11 or EXO1 nuclease, leading to genomic instability or cell death [42]. When BRCA1/2 is defective, MRE11 targets unprotected reversed forks and starts fork resection, leading to the formation of ssDNA flaps in the reversed fork [43], which MUS81 cleaves, resulting in the generation and accumulation of DSBs [38]. These nuclease-induced DSBs are repaired by POLD3-dependent DNA synthesis, contributing to restarting replication recovery from cleaved forks via the break-induced replication (BIR) pathway [38]. Many factors are involved in the repair of such DSBs as parts of DNA damage signaling, including ataxiatelangiectasia-mutated protein (ATM).

Recently, spartan (SPRTN), a DNA-dependent metalloprotease, has been reported to be directly involved in the repair of PARP1-DNA complexes [44]. In the body, SPRTN is recruited to and interacts with trapped PARP1 at replication forks, and it facilitates the repair of bulky PARP1-DNA complexes via proteolytic digestion, leading to replication bypass of PARP1-DNA complexes to ensure fork progression, conferring PARPi resistance.

Overall, integrity of fork stabilization mechanisms on PARP trapping plays major roles in determining sensitivity to PARPis. A number of nuclear factors involved in this integrity can, therefore, be potential targets to overcome PARPi resistance. Especially, since some DDRs usually function to counteract fork stabilization, the inhibitors against constituents of DDR (DDRis) may have a potential to promote fork stabilization. The deficiency or inhibition of ATM are potential targets or candidates to improve the efficacy of PARPi [45]. Preclinical studies demonstrated that the ATM inhibitor AZD0156 enhanced the effects of olaparib across a panel of breast, gastric, and lung cancer cell lines, as well as potentiating the efficacy of olaparib in patient-derived xenografts (PDXs) of breast cancers [46]. Phase I studies of AZD0156 are currently ongoing (NCT02588105). The combined inhibition of PARP and ATR was shown to overcome PARPi resistance in PDXs of ovarian cancer [47]. The phase II trial (CAPRI trial) to use the ATR inhibitor ceralasertib (AZD6738) in combination with olaparib for recurrent ovarian cancer (NCT03462342) is ongoing [48], but has reported preliminary results of an overall response rate (ORR) of

approximately 50% for the HRD patients with acquired PARPi resistance. The CHK1 inhibitor prexasertib has been combined with olaparib for HGSOC in a phase II study, and an ORR of 29% was reported in *BRCA*-wild type recurrent HGSOC or high-grade endometrioid ovarian carcinoma (HGEOC) patients [49]. The Wee1 inhibitor adavosertib was combined with olaparib for patients with PARPi-resistant ovarian cancer in the phase II EFFORT study, showing an overall RR of 29% in combination vs. 23% for adavosertib alone, and a median PFS of 6.8 months in combination vs. 5.5 months for adavosertib alone [50]. Adavosertib was combined with olaparib for refractory or recurrent ovarian cancer patients with *TP53* and/or *KRAS* mutations in a phase-II OLAPCO trial (NCT02576444) [51], based on the proposed concept that the combination of Wee1 inhibitors and *TP53* loss will theoretically give coordinated efficacy, since Wee1 involves G2/M transition and p53 involves G1/S transition. Encouraging results, with an ORR of 43%, were obtained. The OLAPCO (NCT02576444) [52] and ATARI (NCT04065269) [53] trials are also ongoing for the combination of olaparib with ceralasertib.

### 6. PARP Inhibition Triggers Tumor Immunogenicity

### 6.1. Defective DNA Damage Responses Enhance Tumor Antigenicity

Maintenance of genome integrity through DNA replication is essential for cell growth and survival. To faithfully repair DNA lesions that threaten genome integrity, multiple DDRs are induced [54], some of which are composed of pathways for DNA repair, including BER, mismatch repair (MMR), HRR, and NHEJ. Defects of DDRs drive not only genomic instability, but also genomic mutability due to the inability to properly repair DNAs (Figure 3).

However, all DDR defects do not equally result in genomic mutability. MMR deficiency involves defects of mismatch repair proteins, including MLH1, PMS2, MSH2, and MSH6, that function to repair mismatched base pairs, insertions, and deletions at DNA duplication [55]. Eventually, MMR deficiency is likely to cause increased point mutations rather than large defects or gains of bases, generating the microsatellite instability (MSI) phenotype. It, therefore, provides readily available reservoirs of neoepitopes caused by elevated tumor mutational burden (TMB), as well as elevated tumor neoantigen burden (TNB), both of which lead to increased immunogenicity with enhanced recognition by T cells [56], triggering robust antitumor immunity. Therefore, superior responses to anti-PD1/anti-PD-L1 therapy can be expected in patients with MMR deficiency [57]. A correlation between increased TMB or TNB and superior effects of immune-checkpoint inhibitors (ICIs) has been proven in a variety of tumor types, leading to the tumor-agnostic approval of pembrolizumab for the treatment of advanced solid tumors with TMB [58]. Defects of other types of DDRs can also generate increased TMB. Defects in HR or BER are known to correlate with increased TMB or TNB, accompanying higher levels of tumor-infiltrating lymphocytes [59–62]. However, there are some conflicting data that HRD status does not result in increased TMB or TNB [63]. In ovarian cancers, genomic analyses showed less than 10 genetic mutations per megabase [64], and TMB or TNB is not likely to occur, whereas other genetic alterations with LOH, gene amplification, and telomeric allelic imbalance, evaluated as HRD scores, are more frequent. In fact, single-agent ICIs have been shown to exhibit less efficacy in ovarian cancer than in other tumor types [65–67].

Even though TMB or TNB due to defective DDRs is required for the initial priming of the antitumor immune responses, the scenario is not so simple, and they do not always lead to subsequent responses with T cell accumulation in some tumor types [68]. Some studies showed that the defects in HR can predict responses to ICI, but in a TMB-independent manner [69]. Therefore, understanding of not only tumor antigenicity, but also tumor adjuvanticity, is a key to exploiting the possible linkage of defective DDRs to tumor immunogenicity.



**Figure 3.** Dysfunction of DDR or PARPis induce the cGAS-STING pathway to activate cytosolic immunity. While some DDR dysfunction leads to increased tumor mutation burden (TMB) or tumor neoantigen burden (TNB), which enhances tumor antigenicity, other DDR dysfunction, such as via PARPis, causes stalled replication forks, triggering ATR/CHK1 activation leading to upregulated PD-L1, and replication stress responses inducing fork degradation via several nucleases, including MRE11, EXO1, and MUS81, result in the generation of cytosolic nuclear acids (cNAs). Fork collapse by the above nucleases induces genetic instability and generates micronuclei, which rupture and become cNAs. These cNAs are recognized by pattern recognition receptors (PRRs) as part of the innate immune response, triggering cGAS-STING signaling to increase cytosolic immunity. These acronyms are defined as follows: ALR, AIM2-like receptors; ATR, ataxia-telangiectasia-mutated protein kinase; cGAS, cyclic GMP-AMP synthase; CHK1, checkpoint kinase 1; DDR, DNA damage response; DSB, double strand break; FEN1, flap endonuclease 1; HR, homologous recombination; MMR, mismatch repair deficiency; PD-L1, programmed cell death ligand-1; RLR, RIG1-like receptors; STING, stimulator of interferon genes; XRCC1, X-ray repair cross-complementing 1.

### 6.2. PARPis Activate the cGAS-STING Pathway and Mediate Cytosolic Immunity

The interface between DDR and immunogenicity has recently attracted special attention in the field of immune-oncology, in which the cyclic GMP-AMP synthase-stimulator of interferon genes (cGAS-STING) pathway plays essential roles [70]. This pathway originally functions as a component of cellular host defense responding to pathogenic invasion, such as viral infection. Foreign cytosolic DNAs from invaded pathogens can be detected by the cytosolic DNA sensor, cGAS, which subsequently activates the STING, leading to activation of type I interferon (IFN) signaling, essential for innate immune responses [71]. Recently, the cGAS-STING pathway has been found to be activated by endogenous defects of DDRs [70,72–74], including defects of BER, HRR, or NHEJ (Figure 3). First, cGAS can detect tumor-derived DNAs, which are accumulated by defective DDRs. It is important to understand why tumor-derived DNAs accumulate in defective DDRs. For example, defects of HRR (HRD) leave DSBs unrepaired or insufficiently repaired, resulting in stalled replication forks, triggering replication stress responses, in which several nucleases are activated. Subsequently, MUS81 functions to resolve stalled replication forks by DNA end resection, and it can mediate the cleavage of DNA structures at stalled replication forks [75]. The resolution of Holliday junctions is promoted by EXO1 as a result of its exonuclease activity [76]. Thus, these nucleases generate fragmented DNAs as by-products of their nuclease activity, which are transported to the cytosol from the nucleus through passive spilling or active transport to generate cytosolic nucleic acids (cNAs). Generated cNAs are perceived by cGAS via various pattern recognition receptors (PRRs), with which innate immune systems recognize pathogenic microorganisms as non-self. This recognition activates synthase activity, leading to further activation of STING, inducing type I interferons and diverse cell-autonomous immune responses, leading to cytosolic immunity. Activation of cytosolic immunity via the cGAS-STING pathway functions to promote T cell infiltration and turn immunologically "cold" tumors into "hot" tumors.

Other than replication stress responses, there is another factor that triggers cGAS-STING signaling. Mitotic defects caused by inadequately repaired DNAs [77] or telomere dysfunction [78] are known to induce chromosomal instability and generate micronuclei via abnormal chromosome segregation, in which the fragile nuclear envelope of micronuclei is likely to rupture, leading to the release of chromatin and nuclear acid components into the cytosol, triggering cGAS-STING signaling (Figure 3). Thus, both DDR defects and mitotic defects result in the cGAS-STING pathway inducing cytosolic immunity. Consistent with this concept, recent studies have shown a novel aspect of the treatment strategy, that DDRtargeting agents [73,77,79,80] or radiotherapy [76,81] can activate cytosolic immunity via the cGAS-STING pathway. In addition, low levels of the cGAS-STING pathway are linked to a poor prognosis in various tumor types, and established tumor cell lines do not usually exhibit this pathway, indicating that the cGAS-STING pathway plays a role in preventing tumor progression [82]. As one of the representative DDR-targeting agents, PARPis were confirmed to elicit tumor-antitumor immunity via the cGAS-STING pathway [83-88]. The landmark study was published by Ding et al., demonstrating that olaparib triggers local and systemic antitumor immunity in a BRCA-deficient context via the activation of the cGAS-STING pathway using a genetically engineered mouse model of HGSOC [85]. They showed that olaparib treatment increased the number of intratumoral CD4<sup>+</sup> and  $CD8^+$  T cells, as well as enhanced production of IFN $\gamma$  and TNF $\alpha$ , together with increased contents of dsDNAs and micronuclei in the HRD context, but not in the homologous recombination proficiency (HRP) context, and the resultant activation of STING signaling was responsible for PARPi-induced antitumor immune responses, proven by STINGknockout mice [85]. These findings suggest the essential concept that PARP inhibition leads to STING-dependent antitumor immunity via generated dsDNAs and micronuclei in tumors with HRD.

In the clinical samples, PARPi-induced antitumor immunity via STING-mediated interferon responses was confirmed. In the RIO clinical trial (EudraCT 2014-003319-12), for triple-negative breast cancers treated with rucaparib, gene expression analysis was performed to compare the molecular profiling of tumor samples before and after the treatment with rucaparib [89]. Subsequently, RNA sequencing identified that type I and type II interferon signaling pathways were upregulated in the biopsy samples from patients treated using rucaparib, together with STING1 expression, which was observed only in HRD tumors.

### 6.3. Upregulation of PD-L1 by PARP Inhibition

Another aspect of PARPi-induced antitumor immunity is the modulation of immune checkpoint expression. Using breast cancer cell lines and mouse xenograft models, Jiao et al. demonstrated that PARPis upregulated PD-L1 expression, associated with GSK3β inactivation [90], and that PD-L1 upregulation by PARPis attenuated the therapeutic efficacy of PARPis via tumor-associated immunosuppression, as well as determining that simultaneous inhibition of PARP and PD-L1 increased sensitivity to PARPi therapy. An

in vitro experiment using ovarian cancer cell lines also confirmed that PD-L1 expression was upregulated after PARPi treatment through the CHK1 pathway [91] (Figure 3). Similar upregulation of PD-L1 by PARPis was more recently observed in pancreatic cancer cells in vitro and in vivo, in which JAK2/STAT3 signaling was involved [92].

## 6.4. Clinical Application to Combine PARPi with ICIs

The PARPi-induced activation of the cGAS-STING pathway and the upregulation of PD-L1 propose molecular rationales for a combination strategy with ICIs. The phase II study (NCT02484404) with olaparib plus the PD-L1 inhibitor (durvalumab), was performed for platinum-resistant recurrent ovarian cancer, and a response rate (RR) of 15% and stable disease (SD) of 38% yielded a disease control rate (DCR) of 53% [93]. A single-arm phase I-II trial of niraparib plus pembrolizumab in patients with platinum-resistant recurrent ovarian cancer reported complete response (CR) in 5% of patients, partial response (PR) in 13%, and SD in 47%, and the ORRs were consistent, irrespectively of platinum-sensitivity, previous bevacizumab treatment, and BRCA or HR status [94]. Of note is that responses in patients with wild-type *BRCA* or HRP were higher than expected with either agent as monotherapy. The phase II trial of a combination of dostarlimab, bevacizumab, and niraparib in patients with platinum-resistant recurrent ovarian cancer showed PR of 18% and SD in 59%, yielding a DCR of 77% [95]. For platinum-sensitive recurrent ovarian cancers, greater clinical benefit was observed. The phase II study of olaparib and durvalumab (MEDIOLA) for patients with gBRCA-mutated platinum-sensitive relapsed ovarian cancer reported ORR of 72% and a median PFS of 11.1 months [96]. The phase II study of olaparib, durvalumab, and bevacizumab (MEDIOLA) for patients with non-gBRCA-mutated platinum-sensitive recurrent ovarian cancer showed ORRs of 31% and 77% with an olaparib + durvalumab doublet and an olaparib + durvalumab + bevacizumab triplet, respectively, with a median PFS of 5.5 months and 14.7 months, respectively [97]. Many phase II-III studies are currently ongoing to compare the efficacy of PARPi monotherapy and in combination with ICIs, especially in the context of maintenance therapy after 1st-line chemotherapy, and the results of these studies will greatly affect and change the concept of maintenance therapy for ovarian cancer.

## 7. PARPis Can Cooperate with Antiangiogenic Agents

Hypoxia is a particular microenvironment of growing tumors, occurring due to the imbalance between tumor growth and angiogenesis. Decreased levels of HR factors, such as BRCA and RAD51, were observed in cancer cells during hypoxic exposure [98], mainly caused by epigenetic silencing of the *BRCA* promoter or the transcriptional repression of the *RAD51* promoter via an E2F4/p130 complex, a transcription factor induced by hypoxia that occupies the promoter to inhibit access of other transcriptional activators [99], compromising the HR pathway (Figure 4). Therefore, hypoxic conditions are likely to impair the function of HR, providing a TME favored by PARPis. On the contrary, hypoxic conditions provide another aspect for the TME of cancer cells exposed to PARPis. Hypoxia triggers the activation of VEGF expression via hypoxia-inducible factor 1 (HIF1) [100]. A recent study found that VEGF interacts with neuropilins (NRPs), a family of VEFG receptors [101], and VEGF/NRP signaling promotes HR via activation of the Hippo pathway transducers YAP and TAZ, which are critical downstream targets of VEGF signaling and which activate the *RAD51* promoter via the transcription factor TEAD (Figure 4).



**Figure 4.** In this figure, PARPis and anti-VEGF/VEGFRi cooperate to induce HRD. The hypoxiainducible factor (HIF1) increases VEGF expression, which interacts with neuropilins (NRPs), and VEGF/NRP signaling promotes homologous recombination (HR) through the Hippo pathway transducers YAP/TAZ, which activate the *RAD51* promoter via the transcription factor TEAD. In contrast, hypoxia inhibits *RAD51* via transcriptional repressor E2F4/p130 or inhibits *BRCA1* by epigenetic silencing, leading to impaired HR. Thus, hypoxia exhibits contradictory actions, inhibition, or activation of HR. In this context, anti-VEGF agents or VEGFR inhibitors (VEGFRis) suppress VEGF/NRP signaling, causing impaired HR, whereas PARPis inhibit BRCA1 and RAD51 expression via increased formation of the transcriptional repressor E2F4/p130, resulting in enhanced HRD. Thus, PARPis and anti-VEGF/VEGFRi can cooperate to elicit the HRD phenotype to achieve additive or synergistic effects.

Thus, the VEGF-NRP-YAP/TAZ axis plays a pivotal role in the hypoxia-induced activation of HR. Taken together, there are contradictory aspects of HR in hypoxia, in that HR potential is inhibited via decreased BRCA or RAD51 activity, whereas it is activated via the VEGF-NRP-YAP/TAZ axis. Therefore, targeting the VEGF-NRP-YAP/TAZ axis by anti-VEGF/VEGFR inhibitors effectively induces HRD in cancer cells in theory, a TME favored by PARPis. Various anti-VEGF/VEGFR inhibitors are thus challenged to enhance or cooperate in the effects of PARPis in clinical settings. It is of note that PARP inhibition is known to down-regulate BRCA1 and RAD51 expressions via increased occupancy of the promoters by the transcriptional repressor E2F4/p130 [102] (Figure 4). Thus, PARPis may create their favored TME by themselves, and can, therefore, exert enhanced efficacy.

Overall, these findings create the rationale for trying the combination of PARPis with antiangiogenics. Such combination therapy was initially studied in a phase II trial, in which olaparib plus cediranib, a VEGFR inhibitor, was compared to olaparib alone for platinum-sensitive recurrent ovarian cancer [103]. With the addition of cediranib, PFS was significantly improved, especially for *BRCA*-wild-type patients (median 16.5 months vs. 5.7 months, HR 0.32, p = 0.008), but not for those with *gBRCA* mutations (median 19.4 months vs. 16.5 months, p = 0.16). Subsequently, OS for *BRCA*-wild-type patients was reported to be significantly improved by the addition of cediranib (median 37.8 months vs. 23.0 months, p = 0.047) [104]. The important finding is that cediranib exerted additive survival effects in the HRP context, but not HRD, consistent with the above rationale. The AVANOVA2 study also investigated the benefit of adding bevacizumab, a VEGF antibody, to niraparib in platinum-sensitive recurrent ovarian cancer, and it confirmed the additive

effect (median PFS 11.9 months vs. 5.5 months, HR 0.35, p < 0.0001), irrespective of HRD status, for patients with HRP (HR 0.36) or HRD (0.40) [105]. The encouraging result of this study is that the additive survival effect was more significant in HRP patients than in HRD patients, considering the rationale of this study. The phase III PAOLA trial compared the efficacy of olaparib plus bevacizumab with bevacizumab alone in stage III-IV high-grade ovarian cancer as maintenance after first-line therapy [25], and it demonstrated the significant improvement of PFS by the addition of olaparib (median 22.1 months vs. 16.6 months, HR 0.59, p < 0.001), especially for the HRD cohorts (median 37.2 vs. 17.7 months, HR 0.33; 95% CI, 0.25–0.45) and for the HRD without BRCA mutations (median 28.1 months vs. 16.6 months, HR 0.43; 95% CI, 0.28–0.66). However, the results of the HRP cohorts, for which the combination strategy was expected to exert a greater additive effect, were disappointing (PFS median 16.9 months vs. 16.0 months, HR 1.00; 95% CI, 0.75–1.35). Furthermore, this study lacked an olaparib-alone arm to identify whether the combination (addition of bevacizumab) improves the outcomes compared to olaparib alone. Another phase III study (NRG-GY004) compared olaparib plus cediranib and olaparib alone to standard platinum-based therapy in recurrent platinum-sensitive ovarian cancer [106]. However, olaparib/cediranib did not improve PFS versus chemotherapy in the overall population (HR 0.86, 95% CI, 0.66–1.10). In gBRCA-mutant patients, olaparib/cediranib showed better PFS than chemotherapy (HR 0.55, 95% CI, 0.32–0.94), and also olaparib (HR 0.63, 95% CI, 0.37–1.07). In gBRCA wild-type patients, olaparib/cediranib or olaparib did not a show superior effect on PFS (HR 0.97, 95% CI, 0.73–1.30 and HR 1.41, 95% CI, 1.07–1.86, respectively). Several trials are ongoing to evaluate the efficacy of combinations of PARPis and antiangiogenic agents.

## 8. Molecular Factors Involved in PARPi Resistance

A variety of molecular factors involved in PARPi resistance have been reported or proposed using in vitro models, mouse models, and clinical samples. Some of these molecular factors affect HR genes or HRR, while others affect the pharmacokinetics of PARPi. The representative mechanisms and factors are described in the following sections.

### 8.1. Reversion Mutation

The representative PARPi resistance mechanisms include the restoration of HR-mediated DNA repair impairment via the reversion mutation of HR genes. Since PARPis do not target oncogenic driver genes, but rather the function of HR genes, the restoration of function of HR genes is likely to reduce the efficacy of PARPis. The secondary mutations of the *BRCA* gene after treatment with platinum or PARPi have the potential to restore the native open reading frame of the *BRCA* gene [107–109], and are therefore named "reversion mutations" (Figures 5 and 6A).

The BRCA reversions account for 20–30% of the cases of disease progression after treatment with platinum or PARPis [110]. Reversion mutations include true reversions (to wildtype sequence), as well as second-site reversions, which are usually intragenic deletions, mostly being flanked by short regions (1–6 bp) of DNA sequence microhomology [107,109] (Figure 5). This microhomology is thought to be the "scar" of DNA-repair processes that use the microhomology region to repair DSB, named microhomology-mediated end joining (MMEJ) [111]. Thus, MMEJ, which probably occurs as a result of impaired HR, may be the predominant cause of the reversion, indicating that HRD may be the primary upstream event driving reversion mutation. Although reversion events occur not only in tumors with BRCA mutation, but also in other HR genes, including the RAD51 gene, most reversion mutations occur in BRCA1/2 [112]. Even where patients had a common pathogenic mutation, the reversion mutations that emerged in each patient were unique [112] (Figure 5). Therefore, no strong propensity was observed for any particular reversion mutation to result from a particular pathogenic mutation. In *BRCA2*, reversion mutations occur in a position-dependent manner, and the N-terminal regions have a "hotspot" of reversion mutations while the C-terminal regions lack them [112]. This indicates that pathogenic

mutations in the C-terminal regions are unlikely to induce second-site mutation and are at lower risk of developing resistance via reversion. Splice-site pathogenic and missense mutations in *BRCA1/2* are not likely to cause reversion mutations, whereas truncating mutations cause them more frequently [112]. It is of note that patients with *BRCA* mutations that involve structural variants, including the homozygous deletion of the entire locus, that are essentially resistant to reversion mutations, are likely to show long-term responses to PARPis [113].



**Figure 5.** Representative types of reversion mutation. Architecture of the homologous recombination (HR) gene is shown. The HR gene with a pathogenic mutation (MT) and a resultant premature stop codon due to out of frame sequences are illustrated in the upper panel. The normal function of the gene is lost as a result of the MT and resultant premature stop codon. As shown in the lower panel, true reversion MT completely restores the pathogenic MT and open reading frame (ORF), while second site deletion, either encompassing or not encompassing the pathogenic MT, mostly being flanked by short regions of DNA sequence microhomology, generates an alternative ORF with a naive stop codon, leading to a restoration of gene function.

### 8.2. BRCA Hypomorphic Proteins

Some variants of BRCA1 have been reported with similar but weaker biological function of the wild-type BRCA1, named hypomorphic variants, which lack some of the entire domain of the protein (Figure 6B). The existence of such hypomorphic variants has been prevalently identified in a PARP-resistant, patient-derived, xenograft (PDX) model [114], suggesting that such hypomorphs play some roles in PARPi-resistance, probably via lesser but functional effects of the protein. Hypomorphic variants are presumed to be generated by gene rearrangements via various mechanisms, such as alternative mRNA splicing or alternative initiation of translation, that can bypass deleterious *gBRCA1* mutation and thereby restore HR function [115–118]. Whether these hypomorphs can exert biological effects equivalent to wild-type BRCA1, and if they can be related to the levels of PARPiresistance, should be tested by pre-clinical models, as well as in clinical settings.



**Figure 6.** Various factors that involve PARPi resistance. PARPi resistance is conferred by proper repair of single strand breaks (SSBs) via PARP-dependent base excision repair (BER) or double strand breaks (DSBs) via homologous recombination repair (HRR), whereas DSBs cannot be repaired in the presence of the 53BP-Shieldin complex, which sequesters the PALB2-BRCA2-RAD51 complex, alternative repair molecules for HRR, conferring the HRD phenotype, leading to replication stress-induced fork collapse and cell death. Of various factors involved in PARPi resistance, seven representative causes are shown by numbers in red boxes that function to support the BER or HRR pathway. The acronyms are defined as follows: HRD, homologous recombination deficiency; HRP, homologous recombination proficiency; PALB2, partner and localizer of BRCA2; PARG, poly (ADP-ribose) glycohydrolase; RAD51, RAD51 recombinase; SLFN11, Schlafen family member 11; 53BP1, p53-binding protein 1.

### 8.3. Epigenetic Reversion

Promoter silencing of the HR genes, including BRCA1 and RAD51C, has been detected in ovarian cancers associated with HRD phenotypes, and this is one of the major causes of gene inactivation [119] (Figure 6A). De-methylation of these promoters can re-express these genes as part of a process named "epigenetic reversion", which can potently convert the HRD to the HRP phenotype, thereby exhibiting PARPi resistance. Whole genome analyses of paired biopsy samples in primary and acquired platinum-resistant HGSOC have shown that the loss of *BRCA1* promoter methylation is associated with platinum-resistance [120]. In a PDX model of ovarian cancer lacking HR gene mutations but harboring RAD51C promoter methylation, treatment with niraparib induced a significant increase in mRAD51C mRNA, associated with loss of the RAD51C promoter methylation, indicating that RAD51C methylation can be reversed by the treatment pressure of PARPi, which restored RAD51C expression, causing PARPi resistance [121]. In patients with HGSOC enrolled in the ARIEL2 Part 1 PARPi study [122], zygosity of BRCA1 methylation was extensively analyzed in archival tumor and pre-treated biopsy samples with a quantitative method [123], and patients with homozygous BRCA1 methylation showed improved clinical outcomes with rucaparib treatment, compared to those with any methylated BRCA1 ever, indicating that homologous BRCA1 methylation is more likely to respond to PARPi, and that heterologous methylation is likely to be associated with resistance. Consistent with this, the PDX model with RAD51C methylation showed the loss of RAD51C methylation after the treatment with rucaparib or niraparib, in which a single unmethylated copy of *RAD51C* was able

to confer PARPi resistance [124]. Thus, methylation zygosity is a potential biomarker to predict PARPi efficacy and, therefore, should be assessed when considering PARPi therapy.

## 8.4. Re-Gaining HR Proficiency under BRCA Mutation

The above three mechanisms of HR-related PARPi resistance are conferred by changes in HR gene status, especially *BRCA1*, that regain HR proficiency. An additional mechanism is proposed to inactivate the *TP53BP1* gene encoding p53-binding protein 1 (53BP1) without changing *BRCA1* mutation status. The 53BP1 acts as the central component of the protein complex known as the 53BP1-Shieldin complex [125] (Figure 6C). In the *BRCA1* mutation setting, a *BRCA1*-independent HR mechanism acts in which PALB2-BRCA2-RAD51 plays an alternative role. The 53BP1-Shieldin complex interferes with this HR mechanism, sequestering the PALB2-BRCA2-RAD51 complex from DNA break sites, and it also plays a role in preventing one of the HR processes, namely nucleolytic DNA end processing, which is an initial process of HR [126–128]. Thus, the 53BP-Shieldin complex functions to disturb HR function, inducing HRD. Eventually, mutations in components of the 53BP1-Shieldin complex result in restored HR in the *BRCA1* mutation setting, leading to PARPi resistance, confirmed by a PDX model [114] or a clinical sample on PARPi progression [129].

## 8.5. PARP1 Mutation

Trapped PARP acts as a DNA poison that causes stalled and collapsed replication forks, leading to cell death. Therefore, the existence of PARP1 is indispensable for PARPi efficacy. Consistent with this, a mouse study using genetically engineered embryonic stem cells demonstrated that *PARP1* loss is the critical determinant of PARPi resistance [130] (Figure 6D). A recent mouse study also showed that *PARP* mutations that caused a loss of the function to bind DNA, even in the presence of PARPi, conferred PARPi resistance [131]. Of note is that a *PARP1* mutation (R591C; c.1771C > T) was detected in an ovarian cancer patient with olaparib resistance, with such mutation predicted to disturb PARP trapping [131]. However, the clinical relevance of *PARP1* mutations has not yet been fully clarified in a larger number of patients, and it will, therefore, be important to know the prevalence and potency to affect PARPi efficacy in ovarian cancer patients.

## 8.6. PARG Loss

It is understood that PARP1/2 are enzymes that catalyze PARylation of their target proteins. As mentioned above, they also catalyze PARylation of themselves (called auto-PARylation), facilitating their dissociation from DNA via electrostatic repulsion between PARylation products and DNA, both of which possess negative charges. This is a mechanism of PARPs to detach from DNA after completing SSB repair. In contrast to this, poly (ADPribose) glycohydrolase (PARG) exerts counteracting activity, degrading nuclear PAR [132]. Therefore, the loss of PARG function leads to enhanced auto-PARylation, allowing PARP to detach from DNA, thereby suppressing PARP-trapping by PARPis (Figure 6E). Genetic screening with multi-omics analyses of matched PARP-sensitive and PARP-resistant mouse breast tumors with BRCA2 mutation identified PARG loss as a major mechanism of PARPi resistance [133]. An invitro study using human cancer cells with BRCA1/2 deficiency demonstrated that chemically-mediated or shRNA-mediated PARG inhibition resulted in increased PAR and decreased PARP1 trapping, as well as increased resistance to PARPi [133]. The immunohistochemical analysis of HGSOC samples demonstrated that PARG protein was broadly expressed in most biopsied samples, but PARG-negative areas could be detected in a significant proportion of samples, which were tightly associated with increased levels of PAR [133]. Thus, PARG is an important mediator of the PARPi response, and the loss of PARG may be a biomarker potentially linked to PARPi resistance.

### 8.7. Schlafen11 Loss

Schlafen family member 11 (SLFN11) is a putative DNA/RNA helicase that is recruited to the stressed replication fork that halts the fork proceeding or promotes degradation,

independently of the canonical DDR and, therefore, suppresses cell proliferation [134]. Expression of SLFN11 was known to be a predictor of sensitivity to multiple DNA-damaging agents, including platinum agents and PARPis [135–137]. To identify the candidate markers of olaparib response, the correlation of gene expression profiling and olaparib responses was examined in various HGSOC cell lines, and SLFN11 expression was confirmed to be the most strongly associated with olaparib responses [138] (Figure 6F). Recently, a retrospective analysis of 110 HGSOC patients was performed from the phase II, randomized, placebo-controlled, olaparib maintenance trial (NCT00753545) [22], in which SLFN11 expression was evaluated [139]. The expression of SLFN11 significantly impacted PFS, in which SLFN11-high patients had a longer PFS of 12.4 months in the olaparib arm, compared to 3.1 months in the placebo arm. In contrast, the SLFN11-low group had a shorter PFS of 6.3 months in the olaparib arm and 5.1 months in the placebo arm. A similar trend also was observed in OS. Thus, high levels of SLFN11 were associated with improved clinical outcomes to olaparib in HGSOC, indicating that SLFN11 can be a candidate predictor of PARPi sensitivity.

## 8.8. ABCB1 Overexpression

One of the crucial drug resistance mechanisms includes the active removal of cytotoxic agents from cancer cells through the overexpression of the ATP-binding cassette (ABC) drug efflux transporter P-glycoprotein ABCB1, known as multidrug resistance 1 (MDR1); various types of anti-cancer agents are substrates of ABC drug efflux [140,141]. Overexpression of ABCB1 has been known to be involved in PARPi resistance in an early study using a genetically engineered mouse model for *BRCA1*-associated breast cancer [142] (Figure 6G). Using clinical samples of the trial that evaluated the efficacy of the olaparib-cediranib combination (NCT02681237) in HGSOC after progression on a PARPi, acquired genomic alterations at PARPi progression were analyzed, and ABCB1 upregulation was observed in approximately 15% of the patients, although other genetic alterations were detected concurrently [143]. Thus, overexpression of ABCB1 may be at least one cause of possible mechanisms of resistance to PARPis, especially in patients heavily treated with chemotherapies prior to PARPi treatment.

## 9. Conclusions

The present review article summarized the genetic background and molecular dynamics of HRD status in treatment with PARPis. Clinicians should pay particular attention to the timing of measuring HRD status during treatment as part of efforts to optimize subsequent therapies. Molecular mechanisms of how PARPis exert cytotoxic effects on stalled replication forks, of which stabilization or collapse is modified by various nuclear factors, that ultimately determine the fate of cells, that is, the sensitivity to PARPi, were described. Some of these factors are strong candidates that should be targeted in strategies to overcome PARPi resistance. Despite this accumulated knowledge, data with clinical samples, especially from post-PARPi tumors, are seriously lacking. Therefore, future clinical trials should be carefully planned to collect samples on enrolment that will be subjected to multi-omics analyses to uncover the factors related to drug responses, hopefully with new sources, such as liquid samples or single cells. These will help develop strategies to overcome PARPi resistance for ovarian cancer patients, especially in later treatment stages.

**Author Contributions:** S.K. wrote the original draft. H.Y., M.I., T.I., K.K., M.T., S.S. and K.N. helped with the review and editing. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

**Acknowledgments:** The authors are very grateful to all the members of the molecular pathology group in the Department of Obstetrics and Gynecology, Shimane University Faculty of Medicine.

**Conflicts of Interest:** S.K. received honoraria for educational seminars from AstraZeneca, Takeda Pharmaceutical Company, and Chugai Pharmaceutical Co., Ltd.

## References

- Coleman, R.L.; Fleming, G.F.; Brady, M.F.; Swisher, E.M.; Steffensen, K.D.; Friedlander, M.; Okamoto, A.; Moore, K.N.; Efrat Ben-Baruch, N.; Werner, T.L.; et al. Veliparib with First-Line Chemotherapy and as Maintenance Therapy in Ovarian Cancer. *N. Engl. J. Med.* **2019**, *381*, 2403–2415. [CrossRef] [PubMed]
- Coleman, R.L.; Oza, A.M.; Lorusso, D.; Aghajanian, C.; Oaknin, A.; Dean, A.; Colombo, N.; Weberpals, J.I.; Clamp, A.; Scambia, G.; et al. Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): A randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* 2017, 390, 1949–1961. [CrossRef]
- González-Martín, A.; Pothuri, B.; Vergote, I.; DePont Christensen, R.; Graybill, W.; Mirza, M.R.; McCormick, C.; Lorusso, D.; Hoskins, P.; Freyer, G.; et al. Niraparib in Patients with Newly Diagnosed Advanced Ovarian Cancer. *N. Engl. J. Med.* 2019, 381, 2391–2402. [CrossRef] [PubMed]
- Mirza, M.R.; Monk, B.J.; Herrstedt, J.; Oza, A.M.; Mahner, S.; Redondo, A.; Fabbro, M.; Ledermann, J.A.; Lorusso, D.; Vergote, I.; et al. Niraparib Maintenance Therapy in Platinum-Sensitive, Recurrent Ovarian Cancer. N. Engl. J. Med. 2016, 375, 2154–2164. [CrossRef] [PubMed]
- Moore, K.; Colombo, N.; Scambia, G.; Kim, B.G.; Oaknin, A.; Friedlander, M.; Lisyanskaya, A.; Floquet, A.; Leary, A.; Sonke, G.S.; et al. Maintenance Olaparib in Patients with Newly Diagnosed Advanced Ovarian Cancer. *N. Engl. J. Med.* 2018, 379, 2495–2505. [CrossRef]
- Pujade-Lauraine, E.; Ledermann, J.A.; Selle, F.; Gebski, V.; Penson, R.T.; Oza, A.M.; Korach, J.; Huzarski, T.; Poveda, A.; Pignata, S.; et al. Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): A double-blind, randomised, placebo-controlled, phase 3 trial. *Lancet Oncol.* 2017, 18, 1274–1284. [CrossRef]
- 7. Pujade-Lauraine, E.; Selle, F.; Scambia, G.; Asselain, B.; Marmé, F.; Lindemann, K.; Colombo, N.; Madry, R.; Glasspool, R.M.; Dubot, C.; et al. LBA33 Maintenance olaparib rechallenge in patients (pts) with ovarian carcinoma (OC) previously treated with a PARP inhibitor (PARPi): Phase IIIb OReO/ENGOT Ov-38 trial. *Ann. Oncol.* 2021, 32, S1308–S1309. [CrossRef]
- 8. Amé, J.C.; Spenlehauer, C.; de Murcia, G. The PARP superfamily. *Bioessays* 2004, 26, 882–893. [CrossRef]
- 9. Gibson, B.A.; Kraus, W.L. New insights into the molecular and cellular functions of poly(ADP-ribose) and PARPs. *Nat. Rev. Mol. Cell Biol.* **2012**, *13*, 411–424. [CrossRef]
- 10. Ray Chaudhuri, A.; Nussenzweig, A. The multifaceted roles of PARP1 in DNA repair and chromatin remodelling. *Nat. Rev. Mol. Cell Biol.* 2017, *18*, 610–621. [CrossRef]
- Roy, R.; Chun, J.; Powell, S.N. BRCA1 and BRCA2: Different roles in a common pathway of genome protection. *Nat. Rev. Cancer* 2011, 12, 68–78. [CrossRef] [PubMed]
- 12. Farmer, H.; McCabe, N.; Lord, C.J.; Tutt, A.N.; Johnson, D.A.; Richardson, T.B.; Santarosa, M.; Dillon, K.J.; Hickson, I.; Knights, C.; et al. Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature* **2005**, 434, 917–921. [CrossRef]
- 13. Lord, C.J.; Ashworth, A. PARP inhibitors: Synthetic lethality in the clinic. *Science* 2017, 355, 1152–1158. [CrossRef] [PubMed]
- 14. Murai, J.; Huang, S.Y.; Das, B.B.; Renaud, A.; Zhang, Y.; Doroshow, J.H.; Ji, J.; Takeda, S.; Pommier, Y. Trapping of PARP1 and PARP2 by Clinical PARP Inhibitors. *Cancer Res.* **2012**, *72*, 5588–5599. [CrossRef] [PubMed]
- 15. Slade, D. PARP and PARG inhibitors in cancer treatment. Genes Dev. 2020, 34, 360–394. [CrossRef]
- Zandarashvili, L.; Langelier, M.F.; Velagapudi, U.K.; Hancock, M.A.; Steffen, J.D.; Billur, R.; Hannan, Z.M.; Wicks, A.J.; Krastev, D.B.; Pettitt, S.J.; et al. Structural basis for allosteric PARP-1 retention on DNA breaks. *Science* 2020, *368*, eaax6367. [CrossRef] [PubMed]
- 17. The Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian carcinoma. *Nature* **2011**, 474, 609–615. [CrossRef]
- Alsop, K.; Fereday, S.; Meldrum, C.; deFazio, A.; Emmanuel, C.; George, J.; Dobrovic, A.; Birrer, M.J.; Webb, P.M.; Stewart, C.; et al. BRCA mutation frequency and patterns of treatment response in BRCA mutation-positive women with ovarian cancer: A report from the Australian Ovarian Cancer Study Group. J. Clin. Oncol. 2012, 30, 2654–2663. [CrossRef]
- Enomoto, T.; Aoki, D.; Hattori, K.; Jinushi, M.; Kigawa, J.; Takeshima, N.; Tsuda, H.; Watanabe, Y.; Yoshihara, K.; Sugiyama, T. The first Japanese nationwide multicenter study of BRCA mutation testing in ovarian cancer: CHARacterizing the cross-sectionaL approach to Ovarian cancer geneTic TEsting of BRCA (CHARLOTTE). Int. J. Gynecol. Cancer 2019, 29, 1043–1049. [CrossRef]
- Frugtniet, B.; Morgan, S.; Murray, A.; Palmer-Smith, S.; White, R.; Jones, R.; Hanna, L.; Fuller, C.; Hudson, E.; Mullard, A.; et al. The detection of germline and somatic BRCA1/2 genetic variants through parallel testing of patients with high-grade serous ovarian cancer: A national retrospective audit. *BJOG Int. J. Obstet. Gynaecol.* 2022, *129*, 433–442. [CrossRef]
- Timms, K.M.; Abkevich, V.; Hughes, E.; Neff, C.; Reid, J.; Morris, B.; Kalva, S.; Potter, J.; Tran, T.V.; Chen, J.; et al. Association of BRCA1/2 defects with genomic scores predictive of DNA damage repair deficiency among breast cancer subtypes. *Breast Cancer Res.* 2014, *16*, 475. [CrossRef] [PubMed]
- 22. Ledermann, J.; Harter, P.; Gourley, C.; Friedlander, M.; Vergote, I.; Rustin, G.; Scott, C.L.; Meier, W.; Shapira-Frommer, R.; Safra, T.; et al. Olaparib maintenance therapy in patients with platinum-sensitive relapsed serous ovarian cancer: A preplanned retrospective analysis of outcomes by BRCA status in a randomised phase 2 trial. *Lancet Oncol.* **2014**, *15*, 852–861. [CrossRef]

- Hodgson, D.R.; Dougherty, B.A.; Lai, Z.; Fielding, A.; Grinsted, L.; Spencer, S.; O'Connor, M.J.; Ho, T.W.; Robertson, J.D.; Lanchbury, J.S.; et al. Candidate biomarkers of PARP inhibitor sensitivity in ovarian cancer beyond the BRCA genes. *Br. J. Cancer* 2018, 119, 1401–1409. [CrossRef] [PubMed]
- Pujade-Lauraine, E.; Brown, J.; Barnicle, A.; Rowe, P.; Lao-Sirieix, P.; Criscione, S.; du Bois, A.; Lorusso, D.; Romero, I.; Petru, E.; et al. Homologous recombination repair mutation gene panels (excluding BRCA) are not predictive of maintenance olaparib plus bevacizumab efficacy in the first-line PAOLA-1/ENGOT-ov25 trial. *Gynecol. Oncol.* 2021, 162, S26–S27. [CrossRef]
- Ray-Coquard, I.; Pautier, P.; Pignata, S.; Pérol, D.; González-Martín, A.; Berger, R.; Fujiwara, K.; Vergote, I.; Colombo, N.; Mäenpää, J.; et al. Olaparib plus Bevacizumab as First-Line Maintenance in Ovarian Cancer. *N. Engl. J. Med.* 2019, 381, 2416–2428. [CrossRef]
- Takaya, H.; Nakai, H.; Sakai, K.; Nishio, K.; Murakami, K.; Mandai, M.; Matsumura, N. Intratumor heterogeneity and homologous recombination deficiency of high-grade serous ovarian cancer are associated with prognosis and molecular subtype and change in treatment course. *Gynecol. Oncol.* 2020, 156, 415–422. [CrossRef]
- Sokolenko, A.P.; Bizin, I.V.; Preobrazhenskaya, E.V.; Gorodnova, T.V.; Ivantsov, A.O.; Iyevleva, A.G.; Savonevich, E.L.; Kotiv, K.B.; Kuligina, E.S.; Imyanitov, E.N. Molecular profiles of BRCA1-associated ovarian cancer treated by platinum-based therapy: Analysis of primary, residual and relapsed tumors. *Int. J. Cancer* 2020, *146*, 1879–1888. [CrossRef]
- Patel, J.N.; Braicu, I.; Timms, K.M.; Solimeno, C.; Tshiaba, P.; Reid, J.; Lanchbury, J.S.; Darb-Esfahani, S.; Ganapathi, M.K.; Sehouli, J.; et al. Characterisation of homologous recombination deficiency in paired primary and recurrent high-grade serous ovarian cancer. *Br. J. Cancer* 2018, *119*, 1060–1066. [CrossRef]
- 29. Cortez, D. Preventing replication fork collapse to maintain genome integrity. DNA Repair 2015, 32, 149–157. [CrossRef]
- 30. Lopes, M.; Cotta-Ramusino, C.; Pellicioli, A.; Liberi, G.; Plevani, P.; Muzi-Falconi, M.; Newlon, C.S.; Foiani, M. The DNA replication checkpoint response stabilizes stalled replication forks. *Nature* **2001**, *412*, 557–561. [CrossRef]
- 31. Sogo, J.M.; Lopes, M.; Foiani, M. Fork reversal and ssDNA accumulation at stalled replication forks owing to checkpoint defects. *Science* 2002, 297, 599–602. [CrossRef] [PubMed]
- 32. Liao, H.; Ji, F.; Helleday, T.; Ying, S. Mechanisms for stalled replication fork stabilization: New targets for synthetic lethality strategies in cancer treatments. *EMBO Rep.* 2018, 19, e46263. [CrossRef] [PubMed]
- Fan, J.; Pavletich, N.P. Structure and conformational change of a replication protein A heterotrimer bound to ssDNA. *Genes Dev.* 2012, 26, 2337–2347. [CrossRef] [PubMed]
- Cimprich, K.A.; Cortez, D. ATR: An essential regulator of genome integrity. *Nat. Rev. Mol. Cell Biol.* 2008, 9, 616–627. [CrossRef]
  [PubMed]
- 35. Sugiyama, T.; Kowalczykowski, S.C. Rad52 protein associates with replication protein A (RPA)-single-stranded DNA to accelerate Rad51-mediated displacement of RPA and presynaptic complex formation. *J. Biol. Chem.* **2002**, 277, 31663–31672. [CrossRef]
- Berti, M.; Ray Chaudhuri, A.; Thangavel, S.; Gomathinayagam, S.; Kenig, S.; Vujanovic, M.; Odreman, F.; Glatter, T.; Graziano, S.; Mendoza-Maldonado, R.; et al. Human RECQ1 promotes restart of replication forks reversed by DNA topoisomerase I inhibition. *Nat. Struct. Mol. Biol.* 2013, 20, 347–354. [CrossRef]
- 37. Couch, F.B.; Cortez, D. Fork reversal, too much of a good thing. Cell Cycle 2014, 13, 1049–1050. [CrossRef]
- Lemaçon, D.; Jackson, J.; Quinet, A.; Brickner, J.R.; Li, S.; Yazinski, S.; You, Z.; Ira, G.; Zou, L.; Mosammaparast, N.; et al. MRE11 and EXO1 nucleases degrade reversed forks and elicit MUS81-dependent fork rescue in BRCA2-deficient cells. *Nat. Commun.* 2017, *8*, 860. [CrossRef]
- Jensen, R.B.; Carreira, A.; Kowalczykowski, S.C. Purified human BRCA2 stimulates RAD51-mediated recombination. *Nature* 2010, 467, 678–683. [CrossRef]
- Schlacher, K.; Wu, H.; Jasin, M. A distinct replication fork protection pathway connects Fanconi anemia tumor suppressors to RAD51-BRCA1/2. *Cancer Cell* 2012, 22, 106–116. [CrossRef]
- 41. Somyajit, K.; Saxena, S.; Babu, S.; Mishra, A.; Nagaraju, G. Mammalian RAD51 paralogs protect nascent DNA at stalled forks and mediate replication restart. *Nucleic Acids Res.* **2015**, *43*, 9835–9855. [CrossRef] [PubMed]
- Dungrawala, H.; Bhat, K.P.; Le Meur, R.; Chazin, W.J.; Ding, X.; Sharan, S.K.; Wessel, S.R.; Sathe, A.A.; Zhao, R.; Cortez, D. RADX Promotes Genome Stability and Modulates Chemosensitivity by Regulating RAD51 at Replication Forks. *Mol. Cell* 2017, 67, 374–386.e5. [CrossRef] [PubMed]
- Schlacher, K.; Christ, N.; Siaud, N.; Egashira, A.; Wu, H.; Jasin, M. Double-strand break repair-independent role for BRCA2 in blocking stalled replication fork degradation by MRE11. *Cell* 2011, 145, 529–542. [CrossRef] [PubMed]
- 44. Saha, L.K.; Murai, Y.; Saha, S.; Jo, U.; Tsuda, M.; Takeda, S.; Pommier, Y. Replication-dependent cytotoxicity and Spartan-mediated repair of trapped PARP1-DNA complexes. *Nucleic Acids Res.* **2021**, *49*, 10493–10506. [CrossRef] [PubMed]
- 45. Gout, J.; Perkhofer, L.; Morawe, M.; Arnold, F.; Ihle, M.; Biber, S.; Lange, S.; Roger, E.; Kraus, J.M.; Stifter, K.; et al. Synergistic targeting and resistance to PARP inhibition in DNA damage repair-deficient pancreatic cancer. *Gut* 2021, *70*, 743–760. [CrossRef]
- Riches, L.C.; Trinidad, A.G.; Hughes, G.; Jones, G.N.; Hughes, A.M.; Thomason, A.G.; Gavine, P.; Cui, A.; Ling, S.; Stott, J.; et al. Pharmacology of the ATM Inhibitor AZD0156: Potentiation of Irradiation and Olaparib Responses Preclinically. *Mol. Cancer* 2020, 19, 13–25. [CrossRef]
- Kim, H.; Xu, H.; George, E.; Hallberg, D.; Kumar, S.; Jagannathan, V.; Medvedev, S.; Kinose, Y.; Devins, K.; Verma, P.; et al. Combining PARP with ATR inhibition overcomes PARP inhibitor and platinum resistance in ovarian cancer models. *Nat. Commun.* 2020, 11, 3726. [CrossRef]

- 48. Shah, P.D.; Wethington, S.L.; Pagan, C.; Latif, N.; Tanyi, J.; Martin, L.P.; Morgan, M.; Burger, R.A.; Haggerty, A.; Zarrin, H.; et al. Combination ATR and PARP Inhibitor (CAPRI): A phase 2 study of ceralasertib plus olaparib in patients with recurrent, platinum-resistant epithelial ovarian cancer. *Gynecol. Oncol.* **2021**, *163*, 246–253. [CrossRef]
- Lee, J.M.; Nair, J.; Zimmer, A.; Lipkowitz, S.; Annunziata, C.M.; Merino, M.J.; Swisher, E.M.; Harrell, M.I.; Trepel, J.B.; Lee, M.J.; et al. Prexasertib, a cell cycle checkpoint kinase 1 and 2 inhibitor, in BRCA wild-type recurrent high-grade serous ovarian cancer: A first-in-class proof-of-concept phase 2 study. *Lancet Oncol.* 2018, 19, 207–215. [CrossRef]
- Westin, S.N.; Coleman, R.L.; Fellman, B.M.; Yuan, Y.; Sood, A.K.; Soliman, P.T.; Wright, A.A.; Horowitz, N.S.; Campos, S.M.; Konstantinopoulos, P.A.; et al. EFFORT: EFFicacy Of adavosertib in parp ResisTance: A randomized two-arm non-comparative phase II study of adavosertib with or without olaparib in women with PARP-resistant ovarian cancer. J. Clin. Oncol. 2021, 39, 5505. [CrossRef]
- 51. Leijen, S.; van Geel, R.M.; Sonke, G.S.; de Jong, D.; Rosenberg, E.H.; Marchetti, S.; Pluim, D.; van Werkhoven, E.; Rose, S.; Lee, M.A.; et al. Phase II Study of WEE1 Inhibitor AZD1775 Plus Carboplatin in Patients With TP53-Mutated Ovarian Cancer Refractory or Resistant to First-Line Therapy Within 3 Months. J. Clin. Oncol. 2016, 34, 4354–4361. [CrossRef]
- 52. Mahdi, H.; Hafez, N.; Doroshow, D.; Sohal, D.; Keedy, V.; Do, K.T.; LoRusso, P.; Jürgensmeier, J.; Avedissian, M.; Sklar, J.; et al. Ceralasertib-Mediated ATR Inhibition Combined With Olaparib in Advanced Cancers Harboring DNA Damage Response and Repair Alterations (Olaparib Combinations). *JCO Precis. Oncol.* **2021**, *5*, 1432–1442. [CrossRef] [PubMed]
- Banerjee, S.; Stewart, J.; Porta, N.; Toms, C.; Leary, A.; Lheureux, S.; Khalique, S.; Tai, J.; Attygalle, A.; Vroobel, K.; et al. ATARI trial: ATR inhibitor in combination with olaparib in gynecological cancers with ARID1A loss or no loss (ENGOT/GYN1/NCRI). *Int. J. Gynecol. Cancer* 2021, 31, 1471–1475. [CrossRef] [PubMed]
- 54. Lord, C.J.; Ashworth, A. The DNA damage response and cancer therapy. Nature 2012, 481, 287–294. [CrossRef] [PubMed]
- 55. Yamamoto, H.; Imai, K. Microsatellite instability: An update. Arch. Toxicol. 2015, 89, 899–921. [CrossRef]
- 56. Chan, T.A.; Yarchoan, M.; Jaffee, E.; Swanton, C.; Quezada, S.A.; Stenzinger, A.; Peters, S. Development of tumor mutation burden as an immunotherapy biomarker: Utility for the oncology clinic. *Ann. Oncol.* **2019**, *30*, 44–56. [CrossRef]
- 57. Le, D.T.; Uram, J.N.; Wang, H.; Bartlett, B.R.; Kemberling, H.; Eyring, A.D.; Skora, A.D.; Luber, B.S.; Azad, N.S.; Laheru, D.; et al. PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N. Engl. J. Med.* **2015**, *372*, 2509–2520. [CrossRef]
- Le, D.T.; Durham, J.N.; Smith, K.N.; Wang, H.; Bartlett, B.R.; Aulakh, L.K.; Lu, S.; Kemberling, H.; Wilt, C.; Luber, B.S.; et al. Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science* 2017, 357, 409–413. [CrossRef]
- McAlpine, J.N.; Porter, H.; Köbel, M.; Nelson, B.H.; Prentice, L.M.; Kalloger, S.E.; Senz, J.; Milne, K.; Ding, J.; Shah, S.P.; et al. BRCA1 and BRCA2 mutations correlate with TP53 abnormalities and presence of immune cell infiltrates in ovarian high-grade serous carcinoma. *Mod. Pathol.* 2012, 25, 740–750. [CrossRef]
- Nolan, E.; Savas, P.; Policheni, A.N.; Darcy, P.K.; Vaillant, F.; Mintoff, C.P.; Dushyanthen, S.; Mansour, M.; Pang, J.B.; Fox, S.B.; et al. Combined immune checkpoint blockade as a therapeutic strategy for BRCA1-mutated breast cancer. *Sci. Transl. Med.* 2017, 9, eaal4922. [CrossRef]
- 61. Viel, A.; Bruselles, A.; Meccia, E.; Fornasarig, M.; Quaia, M.; Canzonieri, V.; Policicchio, E.; Urso, E.D.; Agostini, M.; Genuardi, M.; et al. A Specific Mutational Signature Associated with DNA 8-Oxoguanine Persistence in MUTYH-defective Colorectal Cancer. *EBioMedicine* **2017**, *20*, 39–49. [CrossRef] [PubMed]
- Kraya, A.A.; Maxwell, K.N.; Wubbenhorst, B.; Wenz, B.M.; Pluta, J.; Rech, A.J.; Dorfman, L.M.; Lunceford, N.; Barrett, A.; Mitra, N.; et al. Genomic Signatures Predict the Immunogenicity of BRCA-Deficient Breast Cancer. *Clin. Cancer Res.* 2019, 25, 4363–4374. [CrossRef] [PubMed]
- 63. Solinas, C.; Marcoux, D.; Garaud, S.; Vitória, J.R.; Van den Eynden, G.; de Wind, A.; De Silva, P.; Boisson, A.; Craciun, L.; Larsimont, D.; et al. BRCA gene mutations do not shape the extent and organization of tumor infiltrating lymphocytes in triple negative breast cancer. *Cancer Lett.* **2019**, *450*, 88–97. [CrossRef]
- 64. Alexandrov, L.B.; Nik-Zainal, S.; Wedge, D.C.; Aparicio, S.A.; Behjati, S.; Biankin, A.V.; Bignell, G.R.; Bolli, N.; Borg, A.; Børresen-Dale, A.L.; et al. Signatures of mutational processes in human cancer. *Nature* **2013**, *500*, 415–421. [CrossRef]
- 65. Matulonis, U.A.; Shapira-Frommer, R.; Santin, A.D.; Lisyanskaya, A.S.; Pignata, S.; Vergote, I.; Raspagliesi, F.; Sonke, G.S.; Birrer, M.; Provencher, D.M.; et al. Antitumor activity and safety of pembrolizumab in patients with advanced recurrent ovarian cancer: Results from the phase II KEYNOTE-100 study. *Ann. Oncol.* **2019**, *30*, 1080–1087. [CrossRef] [PubMed]
- 66. Hamanishi, J.; Takeshima, N.; Katsumata, N.; Ushijima, K.; Kimura, T.; Takeuchi, S.; Matsumoto, K.; Ito, K.; Mandai, M.; Nakai, H.; et al. Nivolumab Versus Gemcitabine or Pegylated Liposomal Doxorubicin for Patients With Platinum-Resistant Ovarian Cancer: Open-Label, Randomized Trial in Japan (NINJA). J. Clin. Oncol. 2021, 39, 3671–3681. [CrossRef] [PubMed]
- 67. Pujade-Lauraine, E.; Fujiwara, K.; Ledermann, J.A.; Oza, A.M.; Kristeleit, R.; Ray-Coquard, I.L.; Richardson, G.E.; Sessa, C.; Yonemori, K.; Banerjee, S.; et al. Avelumab alone or in combination with chemotherapy versus chemotherapy alone in platinum-resistant or platinum-refractory ovarian cancer (JAVELIN Ovarian 200): An open-label, three-arm, randomised, phase 3 study. *Lancet Oncol.* **2021**, *22*, 1034–1046. [CrossRef]
- Spranger, S.; Luke, J.J.; Bao, R.; Zha, Y.; Hernandez, K.M.; Li, Y.; Gajewski, A.P.; Andrade, J.; Gajewski, T.F. Density of immunogenic antigens does not explain the presence or absence of the T-cell-inflamed tumor microenvironment in melanoma. *Proc. Natl. Acad. Sci. USA* 2016, 113, e7759–e7768. [CrossRef]

- Hsiehchen, D.; Hsieh, A.; Samstein, R.M.; Lu, T.; Beg, M.S.; Gerber, D.E.; Wang, T.; Morris, L.G.T.; Zhu, H. DNA Repair Gene Mutations as Predictors of Immune Checkpoint Inhibitor Response beyond Tumor Mutation Burden. *Cell Rep. Med.* 2020, 1, 100034. [CrossRef]
- 70. Chen, Q.; Sun, L.; Chen, Z.J. Regulation and function of the cGAS-STING pathway of cytosolic DNA sensing. *Nat. Immunol.* **2016**, 17, 1142–1149. [CrossRef]
- Ishikawa, H.; Barber, G.N. STING is an endoplasmic reticulum adaptor that facilitates innate immune signalling. *Nature* 2008, 455, 674–678. [CrossRef] [PubMed]
- 72. Barber, G.N. STING: Infection, inflammation and cancer. Nat. Rev. Immunol. 2015, 15, 760–770. [CrossRef] [PubMed]
- Parkes, E.E.; Walker, S.M.; Taggart, L.E.; McCabe, N.; Knight, L.A.; Wilkinson, R.; McCloskey, K.D.; Buckley, N.E.; Savage, K.I.; Salto-Tellez, M.; et al. Activation of STING-Dependent Innate Immune Signaling By S-Phase-Specific DNA Damage in Breast Cancer. J. Natl. Cancer Inst. 2017, 109, djw199. [CrossRef] [PubMed]
- 74. Chabanon, R.M.; Rouanne, M.; Lord, C.J.; Soria, J.C.; Pasero, P.; Postel-Vinay, S. Targeting the DNA damage response in immuno-oncology: Developments and opportunities. *Nat. Rev. Cancer* 2021, 21, 701–717. [CrossRef]
- Ho, S.S.; Zhang, W.Y.; Tan, N.Y.; Khatoo, M.; Suter, M.A.; Tripathi, S.; Cheung, F.S.; Lim, W.K.; Tan, P.H.; Ngeow, J.; et al. The DNA Structure-Specific Endonuclease MUS81 Mediates DNA Sensor STING-Dependent Host Rejection of Prostate Cancer Cells. *Immunity* 2016, 44, 1177–1189. [CrossRef]
- Erdal, E.; Haider, S.; Rehwinkel, J.; Harris, A.L.; McHugh, P.J. A prosurvival DNA damage-induced cytoplasmic interferon response is mediated by end resection factors and is limited by Trex1. *Genes Dev.* 2017, *31*, 353–369. [CrossRef]
- 77. Harding, S.M.; Benci, J.L.; Irianto, J.; Discher, D.E.; Minn, A.J.; Greenberg, R.A. Mitotic progression following DNA damage enables pattern recognition within micronuclei. *Nature* 2017, 548, 466–470. [CrossRef]
- Chen, Y.A.; Shen, Y.L.; Hsia, H.Y.; Tiang, Y.P.; Sung, T.L.; Chen, L.Y. Extrachromosomal telomere repeat DNA is linked to ALT development via cGAS-STING DNA sensing pathway. *Nat. Struct. Mol. Biol.* 2017, 24, 1124–1131. [CrossRef]
- Kitai, Y.; Kawasaki, T.; Sueyoshi, T.; Kobiyama, K.; Ishii, K.J.; Zou, J.; Akira, S.; Matsuda, T.; Kawai, T. DNA-Containing Exosomes Derived from Cancer Cells Treated with Topotecan Activate a STING-Dependent Pathway and Reinforce Antitumor Immunity. J. Immunol. 2017, 198, 1649–1659. [CrossRef]
- Härtlova, A.; Erttmann, S.F.; Raffi, F.A.; Schmalz, A.M.; Resch, U.; Anugula, S.; Lienenklaus, S.; Nilsson, L.M.; Kröger, A.; Nilsson, J.A.; et al. DNA damage primes the type I interferon system via the cytosolic DNA sensor STING to promote antimicrobial innate immunity. *Immunity* 2015, 42, 332–343. [CrossRef]
- Mackenzie, K.J.; Carroll, P.; Martin, C.A.; Murina, O.; Fluteau, A.; Simpson, D.J.; Olova, N.; Sutcliffe, H.; Rainger, J.K.; Leitch, A.; et al. cGAS surveillance of micronuclei links genome instability to innate immunity. *Nature* 2017, 548, 461–465. [CrossRef] [PubMed]
- Yum, S.; Li, M.H.; Frankel, A.E.; Chen, Z.J.J. Roles of the cGAS-STING Pathway in Cancer Immunosurveillance and Immunotherapy. Annu. Rev. Cancer Biol. 2019, 3, 323–344. [CrossRef]
- Chabanon, R.M.; Muirhead, G.; Krastev, D.B.; Adam, J.; Morel, D.; Garrido, M.; Lamb, A.; Hénon, C.; Dorvault, N.; Rouanne, M.; et al. PARP inhibition enhances tumor cell-intrinsic immunity in ERCC1-deficient non-small cell lung cancer. *J. Clin. Investig.* 2019, 129, 1211–1228. [CrossRef] [PubMed]
- Pantelidou, C.; Sonzogni, O.; De Oliveria Taveira, M.; Mehta, A.K.; Kothari, A.; Wang, D.; Visal, T.; Li, M.K.; Pinto, J.; Castrillon, J.A.; et al. PARP Inhibitor Efficacy Depends on CD8(+) T-cell Recruitment via Intratumoral STING Pathway Activation in BRCA-Deficient Models of Triple-Negative Breast Cancer. *Cancer Discov.* 2019, *9*, 722–737. [CrossRef]
- 85. Ding, L.; Kim, H.J.; Wang, Q.; Kearns, M.; Jiang, T.; Ohlson, C.E.; Li, B.B.; Xie, S.; Liu, J.F.; Stover, E.H.; et al. PARP Inhibition Elicits STING-Dependent Antitumor Immunity in Brca1-Deficient Ovarian Cancer. *Cell Rep.* **2018**, *25*, 2972–2980.e5. [CrossRef]
- Reisländer, T.; Lombardi, E.P.; Groelly, F.J.; Miar, A.; Porru, M.; Di Vito, S.; Wright, B.; Lockstone, H.; Biroccio, A.; Harris, A.; et al. BRCA2 abrogation triggers innate immune responses potentiated by treatment with PARP inhibitors. *Nat. Commun.* 2019, 10, 3143. [CrossRef]
- 87. Wang, Z.; Sun, K.; Xiao, Y.; Feng, B.; Mikule, K.; Ma, X.; Feng, N.; Vellano, C.P.; Federico, L.; Marszalek, J.R.; et al. Niraparib activates interferon signaling and potentiates anti-PD-1 antibody efficacy in tumor models. *Sci. Rep.* **2019**, *9*, 1853. [CrossRef]
- Kim, C.; Wang, X.D.; Yu, Y. PARP1 inhibitors trigger innate immunity via PARP1 trapping-induced DNA damage response. *eLife* 2020, 9, e60637. [CrossRef]
- 89. Chopra, N.; Tovey, H.; Pearson, A.; Cutts, R.; Toms, C.; Proszek, P.; Hubank, M.; Dowsett, M.; Dodson, A.; Daley, F.; et al. Homologous recombination DNA repair deficiency and PARP inhibition activity in primary triple negative breast cancer. *Nat. Commun.* **2020**, *11*, 2662. [CrossRef]
- Jiao, S.; Xia, W.; Yamaguchi, H.; Wei, Y.; Chen, M.K.; Hsu, J.M.; Hsu, J.L.; Yu, W.H.; Du, Y.; Lee, H.H.; et al. PARP Inhibitor Upregulates PD-L1 Expression and Enhances Cancer-Associated Immunosuppression. *Clin. Cancer Res.* 2017, 23, 3711–3720. [CrossRef]
- Xue, C.; Xu, Y.; Ye, W.; Xie, Q.; Gao, H.; Xu, B.; Zhang, D.; Jiang, J. Expression of PD-L1 in ovarian cancer and its synergistic antitumor effect with PARP inhibitor. *Gynecol. Oncol.* 2020, 157, 222–233. [CrossRef] [PubMed]
- Wang, Y.; Zheng, K.; Xiong, H.; Huang, Y.; Chen, X.; Zhou, Y.; Qin, W.; Su, J.; Chen, R.; Qiu, H.; et al. PARP Inhibitor Upregulates PD-L1 Expression and Provides a New Combination Therapy in Pancreatic Cancer. *Front. Immunol.* 2021, 12, 762989. [CrossRef] [PubMed]

- Lee, J.M.; Annunziata, C.M.; Houston, N.; Kohn, E.C.; Lipkowitz, S.; Minasian, L.; Nichols, E.; Trepel, J.; Trewhitt, K.; Zia, F.; et al. A phase II study of durvalumab, a PD-L1 inhibitor and olaparib in recurrent ovarian cancer (OvCa). *Ann. Oncol.* 2018, 29, viii334. [CrossRef]
- Konstantinopoulos, P.A.; Waggoner, S.; Vidal, G.A.; Mita, M.; Moroney, J.W.; Holloway, R.; Van Le, L.; Sachdev, J.C.; Chapman-Davis, E.; Colon-Otero, G.; et al. Single-Arm Phases 1 and 2 Trial of Niraparib in Combination With Pembrolizumab in Patients With Recurrent Platinum-Resistant Ovarian Carcinoma. *JAMA Oncol.* 2019, *5*, 1141–1149. [CrossRef] [PubMed]
- 95. Liu, J.; Gaillard, S.; Hendrickson, A.W.; Moroney, J.; Yeku, O.; Diver, E.; Gunderson, C.; Arend, R.; Ratner, E.; Samnotra, V.; et al. An open-label phase II study of dostarlimab (TSR-042), bevacizumab (bev), and niraparib combination in patients (pts) with platinum-resistant ovarian cancer (PROC): Cohort A of the OPAL trial. *Gynecol. Oncol.* **2021**, *162*, S17–S18. [CrossRef]
- 96. Drew, Y.; Kaufman, B.; Banerjee, S.; Lortholary, A.; Hong, S.H.; Park, Y.H.; Zimmermann, S.; Roxburgh, P.; Ferguson, M.; Alvarez, R.H.; et al. Phase II study of olaparib + durvalumab (MEDIOLA): Updated results in germline BRCA-mutated platinum-sensitive relapsed (PSR) ovarian cancer (OC). *Ann. Oncol.* **2019**, *30*, v485–v486. [CrossRef]
- 97. Drew, Y.; Penson, R.T.; O'Malley, D.M.; Kim, J.W.; Zimmermann, S.; Roxburgh, P.; Sohn, J.; Stemmer, S.M.; Bastian, S.; Ferguson, M.; et al. 814MO Phase II study of olaparib (O) plus durvalumab (D) and bevacizumab (B) (MEDIOLA): Initial results in patients (pts) with non-germline BRCA-mutated (non-gBRCAm) platinum sensitive relapsed (PSR) ovarian cancer (OC). Ann. Oncol. 2020, 31, S615–S616. [CrossRef]
- Bindra, R.S.; Gibson, S.L.; Meng, A.; Westermark, U.; Jasin, M.; Pierce, A.J.; Bristow, R.G.; Classon, M.K.; Glazer, P.M. Hypoxiainduced down-regulation of BRCA1 expression by E2Fs. *Cancer Res.* 2005, 65, 11597–11604. [CrossRef]
- 99. Bindra, R.S.; Glazer, P.M. Repression of RAD51 gene expression by E2F4/p130 complexes in hypoxia. *Oncogene* 2007, 26, 2048–2057. [CrossRef]
- 100. Forsythe, J.A.; Jiang, B.H.; Iyer, N.V.; Agani, F.; Leung, S.W.; Koos, R.D.; Semenza, G.L. Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1. *Mol. Cell. Biol.* **1996**, *16*, 4604–4613. [CrossRef]
- Elaimy, A.L.; Amante, J.J.; Zhu, L.J.; Wang, M.; Walmsley, C.S.; FitzGerald, T.J.; Goel, H.L.; Mercurio, A.M. The VEGF receptor neuropilin 2 promotes homologous recombination by stimulating YAP/TAZ-mediated Rad51 expression. *Proc. Natl. Acad. Sci.* USA 2019, 116, 14174–14180. [CrossRef] [PubMed]
- 102. Hegan, D.C.; Lu, Y.; Stachelek, G.C.; Crosby, M.E.; Bindra, R.S.; Glazer, P.M. Inhibition of poly(ADP-ribose) polymerase down-regulates BRCA1 and RAD51 in a pathway mediated by E2F4 and p130. *Proc. Natl. Acad. Sci. USA* 2010, 107, 2201–2206. [CrossRef] [PubMed]
- 103. Liu, J.F.; Barry, W.T.; Birrer, M.; Lee, J.M.; Buckanovich, R.J.; Fleming, G.F.; Rimel, B.; Buss, M.K.; Nattam, S.; Hurteau, J.; et al. Combination cediranib and olaparib versus olaparib alone for women with recurrent platinum-sensitive ovarian cancer: A randomised phase 2 study. *Lancet Oncol.* 2014, 15, 1207–1214. [CrossRef]
- 104. Liu, J.F.; Barry, W.T.; Birrer, M.; Lee, J.M.; Buckanovich, R.J.; Fleming, G.F.; Rimel, B.J.; Buss, M.K.; Nattam, S.R.; Hurteau, J.; et al. Overall survival and updated progression-free survival outcomes in a randomized phase II study of combination cediranib and olaparib versus olaparib in relapsed platinum-sensitive ovarian cancer. Ann. Oncol. 2019, 30, 551–557. [CrossRef]
- 105. Mirza, M.R.; Avall Lundqvist, E.; Birrer, M.J.; dePont Christensen, R.; Nyvang, G.B.; Malander, S.; Anttila, M.; Werner, T.L.; Lund, B.; Lindahl, G.; et al. Niraparib plus bevacizumab versus niraparib alone for platinum-sensitive recurrent ovarian cancer (NSGO-AVANOVA2/ENGOT-ov24): A randomised, phase 2, superiority trial. *Lancet Oncol.* 2019, 20, 1409–1419. [CrossRef]
- 106. Liu, J.F.; Brady, M.F.; Matulonis, U.A.; Miller, A.; Kohn, E.C.; Swisher, E.M.; Cella, D.; Tew, W.P.; Cloven, N.G.; Muller, C.Y.; et al. Olaparib With or Without Cediranib Versus Platinum-Based Chemotherapy in Recurrent Platinum-Sensitive Ovarian Cancer (NRG-GY004): A Randomized, Open-Label, Phase III Trial. J. Clin. Oncol. 2022, 40, JCO2102011. [CrossRef]
- 107. Sakai, W.; Swisher, E.M.; Karlan, B.Y.; Agarwal, M.K.; Higgins, J.; Friedman, C.; Villegas, E.; Jacquemont, C.; Farrugia, D.J.; Couch, F.J.; et al. Secondary mutations as a mechanism of cisplatin resistance in BRCA2-mutated cancers. *Nature* 2008, 451, 1116–1120. [CrossRef]
- 108. Swisher, E.M.; Sakai, W.; Karlan, B.Y.; Wurz, K.; Urban, N.; Taniguchi, T. Secondary BRCA1 mutations in BRCA1-mutated ovarian carcinomas with platinum resistance. *Cancer Res.* 2008, *68*, 2581–2586. [CrossRef]
- Edwards, S.L.; Brough, R.; Lord, C.J.; Natrajan, R.; Vatcheva, R.; Levine, D.A.; Boyd, J.; Reis-Filho, J.S.; Ashworth, A. Resistance to therapy caused by intragenic deletion in BRCA2. *Nature* 2008, 451, 1111–1115. [CrossRef]
- 110. Tobalina, L.; Armenia, J.; Irving, E.; O'Connor, M.J.; Forment, J.V. A meta-analysis of reversion mutations in BRCA genes identifies signatures of DNA end-joining repair mechanisms driving therapy resistance. *Ann. Oncol.* **2021**, *32*, 103–112. [CrossRef]
- Sinha, S.; Villarreal, D.; Shim, E.Y.; Lee, S.E. Risky business: Microhomology-mediated end joining. *Mutat. Res. Mol. Mech. Mutagen.* 2016, 788, 17–24. [CrossRef] [PubMed]
- 112. Pettitt, S.J.; Frankum, J.R.; Punta, M.; Lise, S.; Alexander, J.; Chen, Y.; Yap, T.A.; Haider, S.; Tutt, A.N.J.; Lord, C.J. Clinical BRCA1/2 Reversion Analysis Identifies Hotspot Mutations and Predicted Neoantigens Associated with Therapy Resistance. *Cancer Discov.* 2020, 10, 1475–1488. [CrossRef] [PubMed]
- 113. Swisher, E.M.; Kristeleit, R.S.; Oza, A.M.; Tinker, A.V.; Ray-Coquard, I.; Oaknin, A.; Coleman, R.L.; Burris, H.A.; Aghajanian, C.; O'Malley, D.M.; et al. Characterization of patients with long-term responses to rucaparib treatment in recurrent ovarian cancer. *Gynecol. Oncol.* 2021, 163, 490–497. [CrossRef] [PubMed]

- 114. Cruz, C.; Castroviejo-Bermejo, M.; Gutiérrez-Enríquez, S.; Llop-Guevara, A.; Ibrahim, Y.H.; Gris-Oliver, A.; Bonache, S.; Morancho, B.; Bruna, A.; Rueda, O.M.; et al. RAD51 foci as a functional biomarker of homologous recombination repair and PARP inhibitor resistance in germline BRCA-mutated breast cancer. *Ann. Oncol.* 2018, 29, 1203–1210. [CrossRef]
- 115. Wang, Y.; Bernhardy, A.J.; Cruz, C.; Krais, J.J.; Nacson, J.; Nicolas, E.; Peri, S.; van der Gulden, H.; van der Heijden, I.; O'Brien, S.W.; et al. The BRCA1-Δ11q Alternative Splice Isoform Bypasses Germline Mutations and Promotes Therapeutic Resistance to PARP Inhibition and Cisplatin. *Cancer Res.* 2016, *76*, 2778–2790. [CrossRef]
- 116. Wang, Y.; Bernhardy, A.J.; Nacson, J.; Krais, J.J.; Tan, Y.F.; Nicolas, E.; Radke, M.R.; Handorf, E.; Llop-Guevara, A.; Balmaña, J.; et al. BRCA1 intronic Alu elements drive gene rearrangements and PARP inhibitor resistance. *Nat. Commun.* **2019**, *10*, 5661. [CrossRef]
- 117. Wang, Y.; Krais, J.J.; Bernhardy, A.J.; Nicolas, E.; Cai, K.Q.; Harrell, M.I.; Kim, H.H.; George, E.; Swisher, E.M.; Simpkins, F.; et al. RING domain-deficient BRCA1 promotes PARP inhibitor and platinum resistance. *J. Clin. Investig.* **2016**, *126*, 3145–3157. [CrossRef]
- 118. Drost, R.; Dhillon, K.K.; van der Gulden, H.; van der Heijden, I.; Brandsma, I.; Cruz, C.; Chondronasiou, D.; Castroviejo-Bermejo, M.; Boon, U.; Schut, E.; et al. BRCA1185delAG tumors may acquire therapy resistance through expression of RING-less BRCA1. *J. Clin. Investig.* 2016, 126, 2903–2918. [CrossRef]
- 119. Knijnenburg, T.A.; Wang, L.; Zimmermann, M.T.; Chambwe, N.; Gao, G.F.; Cherniack, A.D.; Fan, H.; Shen, H.; Way, G.P.; Greene, C.S.; et al. Genomic and Molecular Landscape of DNA Damage Repair Deficiency across The Cancer Genome Atlas. *Cell Rep.* 2018, 23, 239–254.e6. [CrossRef]
- 120. Patch, A.M.; Christie, E.L.; Etemadmoghadam, D.; Garsed, D.W.; George, J.; Fereday, S.; Nones, K.; Cowin, P.; Alsop, K.; Bailey, P.J.; et al. Whole-genome characterization of chemoresistant ovarian cancer. *Nature* **2015**, *521*, 489–494. [CrossRef]
- 121. Hurley, R.M.; McGehee, C.D.; Nesic, K.; Correia, C.; Weiskittel, T.M.; Kelly, R.L.; Venkatachalam, A.; Hou, X.; Pathoulas, N.M.; Meng, X.W.; et al. Characterization of a RAD51C-silenced high-grade serous ovarian cancer model during development of PARP inhibitor resistance. NAR Cancer 2021, 3, zcab028. [CrossRef] [PubMed]
- 122. Swisher, E.M.; Lin, K.K.; Oza, A.M.; Scott, C.L.; Giordano, H.; Sun, J.; Konecny, G.E.; Coleman, R.L.; Tinker, A.V.; O'Malley, D.M.; et al. Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): An international, multicentre, open-label, phase 2 trial. *Lancet Oncol.* 2017, 18, 75–87. [CrossRef]
- 123. Kondrashova, O.; Topp, M.; Nesic, K.; Lieschke, E.; Ho, G.Y.; Harrell, M.I.; Zapparoli, G.V.; Hadley, A.; Holian, R.; Boehm, E.; et al. Methylation of all BRCA1 copies predicts response to the PARP inhibitor rucaparib in ovarian carcinoma. *Nat. Commun.* 2018, *9*, 3970. [CrossRef] [PubMed]
- 124. Nesic, K.; Kondrashova, O.; Hurley, R.M.; McGehee, C.D.; Vandenberg, C.J.; Ho, G.Y.; Lieschke, E.; Dall, G.; Bound, N.; Shield-Artin, K.; et al. Acquired RAD51C Promoter Methylation Loss Causes PARP Inhibitor Resistance in High-Grade Serous Ovarian Carcinoma. *Cancer Res.* 2021, *81*, 4709–4722. [CrossRef] [PubMed]
- 125. Mirman, Z.; de Lange, T. 53BP1: A DSB escort. Genes Dev. 2020, 34, 7–23. [CrossRef] [PubMed]
- 126. Zong, D.; Adam, S.; Wang, Y.; Sasanuma, H.; Callén, E.; Murga, M.; Day, A.; Kruhlak, M.J.; Wong, N.; Munro, M.; et al. BRCA1 Haploinsufficiency Is Masked by RNF168-Mediated Chromatin Ubiquitylation. *Mol. Cell* **2019**, *73*, 1267–1281.e7. [CrossRef]
- 127. Callen, E.; Zong, D.; Wu, W.; Wong, N.; Stanlie, A.; Ishikawa, M.; Pavani, R.; Dumitrache, L.C.; Byrum, A.K.; Mendez-Dorantes, C.; et al. 53BP1 Enforces Distinct Pre- and Post-resection Blocks on Homologous Recombination. *Mol. Cell* 2020, 77, 26–38.e7. [CrossRef]
- 128. Belotserkovskaya, R.; Raga Gil, E.; Lawrence, N.; Butler, R.; Clifford, G.; Wilson, M.D.; Jackson, S.P. PALB2 chromatin recruitment restores homologous recombination in BRCA1-deficient cells depleted of 53BP1. *Nat. Commun.* 2020, *11*, 819. [CrossRef]
- 129. Waks, A.G.; Cohen, O.; Kochupurakkal, B.; Kim, D.; Dunn, C.E.; Buendia Buendia, J.; Wander, S.; Helvie, K.; Lloyd, M.R.; Marini, L.; et al. Reversion and non-reversion mechanisms of resistance to PARP inhibitor or platinum chemotherapy in BRCA1/2-mutant metastatic breast cancer. *Ann. Oncol.* 2020, *31*, 590–598. [CrossRef]
- 130. Pettitt, S.J.; Rehman, F.L.; Bajrami, I.; Brough, R.; Wallberg, F.; Kozarewa, I.; Fenwick, K.; Assiotis, I.; Chen, L.; Campbell, J.; et al. A genetic screen using the PiggyBac transposon in haploid cells identifies Parp1 as a mediator of olaparib toxicity. *PLoS ONE* 2013, *8*, e61520. [CrossRef]
- 131. Pettitt, S.J.; Krastev, D.B.; Brandsma, I.; Dréan, A.; Song, F.; Aleksandrov, R.; Harrell, M.I.; Menon, M.; Brough, R.; Campbell, J.; et al. Genome-wide and high-density CRISPR-Cas9 screens identify point mutations in PARP1 causing PARP inhibitor resistance. *Nat. Commun.* 2018, *9*, 1849. [CrossRef] [PubMed]
- 132. Pascal, J.M.; Ellenberger, T. The rise and fall of poly(ADP-ribose): An enzymatic perspective. *DNA Repair* **2015**, *32*, 10–16. [CrossRef] [PubMed]
- 133. Gogola, E.; Duarte, A.A.; de Ruiter, J.R.; Wiegant, W.W.; Schmid, J.A.; de Bruijn, R.; James, D.I.; Guerrero Llobet, S.; Vis, D.J.; Annunziato, S.; et al. Selective Loss of PARG Restores PARylation and Counteracts PARP Inhibitor-Mediated Synthetic Lethality. *Cancer Cell* 2018, 33, 1078–1093.e12. [CrossRef] [PubMed]
- 134. Murai, J.; Tang, S.W.; Leo, E.; Baechler, S.A.; Redon, C.E.; Zhang, H.; Al Abo, M.; Rajapakse, V.N.; Nakamura, E.; Jenkins, L.M.M.; et al. SLFN11 Blocks Stressed Replication Forks Independently of ATR. *Mol. Cell* **2018**, *69*, 371–384.e6. [CrossRef]
- Zoppoli, G.; Regairaz, M.; Leo, E.; Reinhold, W.C.; Varma, S.; Ballestrero, A.; Doroshow, J.H.; Pommier, Y. Putative DNA/RNA helicase Schlafen-11 (SLFN11) sensitizes cancer cells to DNA-damaging agents. *Proc. Natl. Acad. Sci. USA* 2012, 109, 15030–15035. [CrossRef]
- Murai, J.; Feng, Y.; Yu, G.K.; Ru, Y.; Tang, S.W.; Shen, Y.; Pommier, Y. Resistance to PARP inhibitors by SLFN11 inactivation can be overcome by ATR inhibition. *Oncotarget* 2016, 7, 76534–76550. [CrossRef]

- 137. Murai, J.; Thomas, A.; Miettinen, M.; Pommier, Y. Schlafen 11 (SLFN11), a restriction factor for replicative stress induced by DNA-targeting anti-cancer therapies. *Pharmacol. Ther.* **2019**, 201, 94–102. [CrossRef]
- 138. Amuzu, S.; Carmona, E.; Mes-Masson, A.M.; Greenwood, C.M.T.; Tonin, P.N.; Ragoussis, J. Candidate Markers of Olaparib Response from Genomic Data Analyses of Human Cancer Cell Lines. *Cancers* **2021**, *13*, 1296. [CrossRef]
- Willis, S.E.; Winkler, C.; Roudier, M.P.; Baird, T.; Marco-Casanova, P.; Jones, E.V.; Rowe, P.; Rodriguez-Canales, J.; Angell, H.K.; Ng, F.S.L.; et al. Retrospective analysis of Schlafen11 (SLFN11) to predict the outcomes to therapies affecting the DNA damage response. *Br. J. Cancer* 2021, 125, 1666–1676. [CrossRef]
- 140. Raymond, M.; Rose, E.; Housman, D.E.; Gros, P. Physical mapping, amplification, and overexpression of the mouse mdr gene family in multidrug-resistant cells. *Mol. Cell. Biol.* **1990**, *10*, 1642–1651. [CrossRef]
- 141. Wu, C.P.; Calcagno, A.M.; Ambudkar, S.V. Reversal of ABC drug transporter-mediated multidrug resistance in cancer cells: Evaluation of current strategies. *Curr. Mol. Pharm.* **2008**, *1*, 93–105. [CrossRef] [PubMed]
- 142. Rottenberg, S.; Jaspers, J.E.; Kersbergen, A.; van der Burg, E.; Nygren, A.O.; Zander, S.A.; Derksen, P.W.; de Bruin, M.; Zevenhoven, J.; Lau, A.; et al. High sensitivity of BRCA1-deficient mammary tumors to the PARP inhibitor AZD2281 alone and in combination with platinum drugs. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 17079–17084. [CrossRef] [PubMed]
- 143. Lheureux, S.; Oaknin, A.; Garg, S.; Bruce, J.P.; Madariaga, A.; Dhani, N.C.; Bowering, V.; White, J.; Accardi, S.; Tan, Q.; et al. EVOLVE: A Multicenter Open-Label Single-Arm Clinical and Translational Phase II Trial of Cediranib Plus Olaparib for Ovarian Cancer after PARP Inhibition Progression. *Clin. Cancer Res.* 2020, 26, 4206–4215. [CrossRef] [PubMed]