

Table S1: Distribution of patients in the extended cohort ($n=17$) in the different CTCAE groups for renal toxicity and distribution of the patients according to KDIGO.

		Baseline	%	After one cycle	%	After two cycles	%	After three cycles	%	After four cycles	%	After five cycles	%
CTCAE Creatinine	CTC 0°	14	82.4	15	88.2	14	82.4	14	82.4	13	76.5	12	70.6
	CTC I°	2	11.8	2	11.8	3	17.6	3	17.6	3	17.6	4	23.5
	CTC II°	1	5.9	0	0.0	0	0.0	0	0.0	1	5.9	1	5.9
	CTC III°/IV°	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
CTCAE eGFR	CTC 0°	4	23.5	4	23.5	5	29.4	5	29.4	4	23.5	5	29.4
	CTC I°	10	58.8	11	64.7	9	52.9	8	47.1	10	58.8	9	52.9
	CTC II°	3	17.6	2	11.8	3	17.6	4	23.5	3	17.6	3	17.6
	CTC III°/IV°	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
KDIGO	G1	4	23.5	4	23.5	5	29.4	5	29.4	4	23.5	5	29.4
	G2	10	58.8	11	64.7	9	52.9	8	47.1	10	58.8	9	52.9
	G3a	2	11.8	2	11.8	2	11.8	3	17.6	1	5.9	0	0.0
	G3b	1	5.9	0	0.0	1	5.9	1	5.9	2	11.8	3	17.6
	G4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

Table S2: Distribution of patients in the extended cohort ($n=17$) in the different CTCAE groups for hematotoxicity.

[illegible]