

Edmonton Symptom Assessment Scale (ESAS)			Control			Switch		
	Items	PSG *	T0 Median (IQR)	T1 Median (IQR)	P value	T0 Median (IQR)	T1 Median (IQR)	P value
Pain	3	3 (1,5)	3 (2,6)	1	4 (2.3,6.8)	3 (2,5)	<0.05	
Tiredness	3	6 (3,8)	4 (3.3,8)	0.70	7 (5.5,8)	6 (3,8)	<0.05	
Drowsiness	2	4 (2,7)	4 (2,7)	0.55	6 (4,8)	4 (2,7)	0.18	
Nausea	1	1 (1,1)	1 (1,1)	0.21	2 (1,4)	1 (1,3)	0.28	
Appetite	3	1 (1,5)	1.5 (1,4.8)	0.15	4 (2.3,8)	5 (1,7)	0.68	
Shortness of Breath	2	1 (1,6.3)	1 (1,4.75)	0.34	4 (1,5)	2 (1,7)	0.97	
Depression	2	1 (1,5)	1.5 (1,5)	0.55	4 (2,6.8)	2 (1,5)	0.23	
Anxiety	2	3 (1,5)	1.5 (1,4.5)	0.12	2 (1,5.8)	2 (1,5)	0.78	
Wellbeing	2	5 (3,7)	5 (2.8,7)	0.76	5 (4,5.5)	5 (3,6)	0.70	

*PSG = Personalised Symptom Goal

Supplementary Table S1: Comparison of Adverse Event scores Across Opioid Switch and Control Groups at T0 and T1