

Supplementary Material

Biotechnological Approach for the Production of Enantiomeric Hydroxylactones Derived from Benzaldehyde and Evaluation of Their Cytotoxic Activity

Marcelina Mazur ^{1,*}, Anna Kudrynska ¹, Aleksandra Pawlak ², Beatriz Hernandez-Suarez ², Bożena Obmińska-Mrukowicz ² and Witold Gładkowski ¹

¹ Department of Chemistry, Wrocław University of Environmental and Life Sciences, Norwida 25, 50-375 Wrocław, Poland; aaahhaaa@ukr.net (A.K.); witold.gladkowski@upwr.edu.pl (W.G.)

² Department of Pharmacology and Toxicology, Wrocław University of Environmental and Life Sciences, Norwida 31, 50-375 Wrocław, Poland; aleksandra.pawlak@upwr.edu.pl (A.P.); beatriz.hernandez-suarez@upwr.edu.pl (B.H.-S.); bozena.obminska-mrukowicz@upwr.edu.pl (B.O.-M.)

* Correspondence: marcelina.mazur@upwr.edu.pl (M.M.); Tel.: +48-713205197






Figure S1. IR spectrum of lactone **2**.

Figure S2. ^{13}C -NMR spectrum of lactone 2.




Figure S4. HMBC spectrum of lactone 2.




Figure S5. The comparison of GC-MS analysis of iodolactone 1 and hydroxylactone 2.