

Correction

Correction: Hayat et al. A Superficial Intramolecular Alignment of Carbon Nitride through Conjugated Monomer for Optimized Photocatalytic CO₂ Reduction. *Catalysts* 2021, 11, 935

Asif Hayat ^{1,†} , Muhammad Sohail ^{2,†}, T.A. Taha ^{3,4} , Asma M. Alenad ⁵ , Ikram Uddin ⁶, Ashiq Hayat ⁷, Tariq Ali ⁸ , Rahim Shah ⁸, Ahmad Irfan ⁹, Wasim Ullah Khan ¹⁰, Arkom Palamanit ¹¹, Yas Al-Hadeethi ^{12,13} , Jawad Ali Shah Syed ¹⁴, Mohammed A. Amin ^{15,*}, Javid Khan ^{16,*} and Sunil Kumar Baburao Mane ^{17,*}

- ¹ State Key Laboratory of Photocatalysis on Energy and Environment, College of Chemistry, Fuzhou University, Fuzhou 350116, China
- ² Institute for Advanced Study, Shenzhen University, Shenzhen 518060, China
- ³ Physics Department, College of Science, Jouf University, Sakaka 75471, Saudi Arabia
- ⁴ Physics and Engineering Mathematics Department, Faculty of Electronic Engineering, Menoufia University, Menouf 32952, Egypt
- ⁵ Chemistry Department, College of Science, Jouf University, Sakaka 75471, Saudi Arabia
- ⁶ CAS Key Laboratory of Science and Technology on Applied Catalysis, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China
- ⁷ Department of Physics, Quaid-e-Azam University, Islamabad 45620, Pakistan
- ⁸ Soochow Institute for Energy and Materials Innovations, College of Energy, Soochow University, Suzhou 215006, China
- ⁹ Department of Chemistry, College of Science, King Khalid University, Abha 61413, Saudi Arabia
- ¹⁰ State Key Laboratory of Optoelectronic Materials and Technologies, School of Materials Science and Engineering, Sun Yat-sen University, Guangzhou 510275, China
- ¹¹ Energy Technology Program, Department of Specialized Engineering, Faculty of Engineering, Prince of Songkla University, 15 Karnjanavanich Rd., Songkhla 90110, Thailand
- ¹² Department of Physics, Faculty of Science, King AbdulAziz University, Jeddah 21589, Saudi Arabia
- ¹³ Lithography in Devices Fabrication Research Group, Deanship of Scientific Research, King AbdulAziz University, Jeddah 21589, Saudi Arabia
- ¹⁴ Department of Material Science and Engineering, College of Engineering and Applied Sciences, Nanjing University, Nanjing 210023, China
- ¹⁵ Department of Chemistry, College of Science, Taif University, Taif 21944, Saudi Arabia
- ¹⁶ School of Chemistry, Sun Yat-Sen University, Guangzhou 510275, China
- ¹⁷ Department of Chemistry, Khaja Bhandanawaz University, Gulbarga 585101, India
- * Correspondence: mohamed@tu.edu.sa (M.A.A.); khanj3@mail.sysu.edu.cn (J.K.); mane@mail.sysu.edu.cn (S.K.B.M.); Tel.: +96-6545707507 (M.A.A.); +86-20-84111823 (J.K.); +91-8095027725 (S.K.B.M.)
- † These authors are equally contributed.



Citation: Hayat, A.; Sohail, M.; Taha, T.A.; Alenad, A.M.; Uddin, I.; Hayat, A.; Ali, T.; Shah, R.; Irfan, A.; Khan, W.U.; et al. Correction: Hayat et al. A Superficial Intramolecular Alignment of Carbon Nitride through Conjugated Monomer for Optimized Photocatalytic CO₂ Reduction. *Catalysts* 2021, 11, 935. *Catalysts* **2023**, *13*, 309. <https://doi.org/10.3390/catal13020309>

Received: 2 January 2023

Revised: 6 January 2023

Accepted: 10 January 2023

Published: 31 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

In Figure 2a [1], the SEM measurements show that the CN nanosheets are closely packed. However, the morphology did not sustain after adding the TDP monomer (change from nanosheets to irregular, rough surface nanoparticles). At the same time, Figure 2b shows magnified irregular nanoparticles with an average diameter of 70–100 nm for CN-TDPx. Asymmetrical, thick, and compressed nanosheets can be seen in Figure 2c for CN, while suppressed surface morphology was observed for CN-TDP_{7.0}, with transparent, smooth, tinny, and much-reduced size, as well as silk-like nanostructures for CN-TDP_{7.0} (Figure 2d).




Figure 2. (a) SEM image for CN and (b) for CN-TDP_{7.0}. (c) TEM image for CN and (d) for CN-TDP_{7.0} samples.

Reference

1. Hayat, A.; Sohail, M.; Taha, T.A.; Alenad, A.M.; Uddin, I.; Hayat, A.; Ali, T.; Shah, R.; Irfan, A.; Khan, W.U.; et al. A superficial intramolecular alignment of carbon nitride through conjugated monomer for optimized photocatalytic CO₂ reduction. *Catalysts* **2021**, *11*, 935. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.