
Supporting Information for

Design of High-Performance Polybenzoxazines with Tunable Extended Networks Based on Resveratrol and Allyl Functional Benzoxazine

Yunliang Xing ¹, Xianru He ^{1,*}, Rui Yang ², Kan Zhang ^{2,*}, Shengfu Yang ³

¹ School of New Energy and Materials, Southwest Petroleum University, Chengdu 610500, China; Phoebe@coryes.com (Y.X.)

² Research School of Polymeric Materials, School of Materials Science and Engineering, Jiangsu University, Zhenjiang 212013, China; yr183346@163.com (R.Y.); zhangkan@ujs.edu.cn (K.Z.)

³ Department of Chemistry, University of Leicester, Leicester LE1 7RH, United Kingdom; sfy1@le.ac.uk (S.Y.)

* Correspondence: xrhe@swpu.edu.cn (X.H.); zhangkan@ujs.edu.cn (K. Z.); Tel.: +86-028-83037406 (X.H.); +86-0571-85016616 (K.Z.)

Received: 22 October 2020; Accepted: 24 November 2020; Published: date




Figure S1. ^1H - ^1H NOESY 2D NMR spectrum of RES-al.




Figure S2. ^1H - ^{13}C HMQC 2D NMR spectrum of RES-al.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).