

Supporting Information

Assessment of physicochemical and *in vivo* biological properties of polymeric nanocapsules based on chitosan and poly(*N*-vinyl pyrrolidone-*alt*-itaconic anhydride)

Kheira Zanoune Dellali¹, Mohammed Dellali¹, Delia Mihaela Rata^{2*}, Anca Niculina Cadinoiu², Leonard Ionut Atanase^{2*}, Marcel Popa^{2,3*}, Mihaela-Claudia Spataru⁴, Carmen Solcan⁴

¹Faculty of Technology, University Hassiba Benbouali, BP 151 02000, Chlef, Algeria

²“Apollonia” University of Iasi, Pacurari Street, No. 11, 700511, Iasi, Romania

³Academy of Romanian Scientists, Splaiul Independentei Street, No 54, 050094, Bucharest, Romania

⁴“Ion Ionescu de la Brad” University of Life Sciences, Mihail Sadoveanu Alley, No 8, 700489, Iasi, Romania




Figure S1: Evolution of mice bodyweight through the experiment (the first measurement was made in the first day of the experiment, the second one at 10 day and the last one at the final of the experiment)

Table S1. *p*-values of Student's t-test for simple 5-FU compared with 5-FU loaded NCs

<i>p</i>-value	CN-1-5FU	CN-2-5FU	CN-3-5FU	CN-4-5FU	CN-5-5FU	CN-6-5FU	CN-7-5FU
5-FU	0.0217	0.0350	0.0308	0.0304	0.0394	0.0454	0.0238

The statistical significance of drug release kinetics was analyzed by the Student's t-test. The values are expressed as mean \pm SE of three parallel measurements, $p < 0.05$ being considered significant.