

Activity of a Recombinant Chitinase of the *Atta sexdens* Ant on Different Forms of the Chitin, and Its Fungicidal Effect Against *Lasiodiplodia theobromae*

Katia Celina Santos Correa^a, William Marcondes Facchinatto^b, Filipe Biagioni Habitzreuter^c, Gabriel Henrique Ribeiro^d, Lucas Gomes Rodrigues^a, Kelli Cristina Micocci^a, Sérgio Paulo Campana-Filho^c, Luiz Alberto Colnago^d, Dulce Helena Ferreira Souza^a

^a Department of Chemistry, Federal University of São Carlos (UFSCar), Rd. Washington Luis, km 235, CEP – 13565-905, São Carlos, SP, Brasil.

^b Aveiro Institute of Materials (CICECO), Department of Chemistry, University of Aveiro, R. Santiago, CP – 3810-193, Aveiro, Portugal.

^c São Carlos Institute of Chemistry, University of São Paulo (IQSC-USP), Av. Trabalhador São-carlense 400, CEP – 13560-590, São Carlos, SP, Brasil.

^d Brazilian Agricultural Research Corporation (Embrapa Instrumentation), R. XV de Novembro 1452, CEP – 13560-970, São Carlos, SP, Brasil.

*Corresponding author:

Dulce Helena F. Souza

E-mail address: dulce@ufscar.br

Phone number: +55 16 3351 8074

Department of Chemistry, Federal University of São Carlos
Rd. Washington Luis, km 235, CEP – 13565-905
São Carlos - SP, Brasil

Summary

Captions	Page
Figure S1. GlcNAc chemical structure.	S2
Figure S2. COZY ^1H - ^1H NMR spectrum for GlcNAc.	S4
Figure S3. Enzymatic activity in the presence of colloidal α -chitin substrate at 28 °C.	S5

Caption	Page
Table S1. Assignments of the ^1H NMR spectra in D_2O for the N-acetyl-d-glucosamine compound	S3




Figure 1S. GlcNAc chemical structure.

Table 1S. Assignments of the ^1H NMR spectra in D_2O for the N-acetyl-d-glucosamine compound. Chemical shifts (ppm), multiplicity, and coupling constants (Hz)

No.	^1H
1'	3.88 (m)
2'	5.19 d (2.8 Hz)
3'	3.40 – 3.95 (m)
4'	3.40 – 3.59 (m)
5'	3.40 – 3.95 (m)
6'	3.40 – 3.95 (m)
NH	8.09 d (8.0 Hz)
7'	-
8'	2.03 s




Figure S2. COZY ^1H - ^1H NMR spectrum for GlcNAc. Signals of the N-acetyl-d-glucosamina hydrolyzed product are highlighted.




Figure S3. Enzymatic activity in the presence of colloidal α -chitin substrate at 28 °C.