

Supplementary Materials: Epirubicin-Complexed Polypeptide Micelle Effectively and Safely Treats Hepatocellular Carcinoma

Qinglin Zhang, Jianxun Ding, Chenwei Lv, Weiguo Xu, Xun Sun and Xiangwei Meng




Figure S1. D_h changes of mPEG-*b*-PGA/EPI in PBS and BSA/PBS of pH 7.4 at 25 °C *versus* time. Data were presented as mean \pm standard deviation ($n = 3$).




Figure S2. *In vitro* cytotoxicity of mPEG-*b*-PGA toward HepG2 cells after incubation for 72 h.




Figure S3. Hemolysis of free EPI (A), mPEG-*b*-PGA/EPI (B), and mPEG-*b*-PGA (C).




Figure S4. Diet amounts (e.g., food and water intake amount) of Kunming mice after intravenous injection of free EPI or mPEG-*b*-PGA/EPI at EPI equivalent dose of 5.0, 10.0, 15.0, or 20.0 mg·(kg BW)⁻¹, or mPEG-*b*-PGA at dose of 250.0, 500.0, 750.0, or 1000.0 mg·(kg BW)⁻¹.



© 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons by Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).