

Supporting Information

Copolyesters Based on 2,5-Furandicarboxylic acid (FDCA): Effect of 2,2,4,4-Tetramethyl-1,3-Cyclobutanediol Units on Their Properties

Jinggang Wang ^{1,2}, Xiaoqing Liu ^{1*}, Jin Zhu ¹ and Yanhua Jiang ¹

¹ Ningbo Institute of Materials Technology and Engineering, Chinese Academy of Sciences,
Ningbo, Zhejiang 315201, (P. R. China)

² University of Chinese Academy of Sciences, Beijing 100049 (P. R. China)

*Correspondence: liuxq@nimte.ac.cn (Xiaoqing Liu); Tel.: 86-574-86685925




Figure S1 ¹H-NMR spectra of dimethyl furan-2,5-dicarboxylate




Figure S2 ¹³C-NMR spectra of dimethyl furan-2,5-dicarboxylate

Table S1 The ^1H -NMR signal assignments for PEF and PETF 10/18

Sample	f_1	f_2	a	b_{cis}	b_{trans}	c_{cis}	c_{trans}
	CH in furan ring	CH in furan ring	CH ₂ in EG	CH in cis-CBDO	CH in trans-CBDO	CH ₃ in cis-CBDO	CH ₃ in trans-CBDO
PEF (ppm)	7.20	-	4.63	-	-	-	-
PETF10/18	7.20-7.23	7.24-7.26	4.63	4.53	4.70	1.07,1.24	1.14

Table S2 The ^1H -NMR signal assignments for PPF and PPTF 10/18

Sample	f_1	f_2	d	e	b_{cis}	b_{trans}	c_{cis}	c_{trans}
	CH in furan ring	CH in furan ring	CH ₂ in PPD	CH ₂ in PPD	CH in cis-CBDO	CH in trans-CBDO	CH ₃ in cis-CBDO	CH ₃ in trans-CBDO
PPF	7.20	-	4.43	2.15	-	-	-	-
PPTF10-18	7.20-7.23	7.24-7.26	4.43	2.15	4.53	4.70	1.07,1.24	1.14

Table S3 The ^1H -NMR signal assignment for PBF and PBTF 10/18

Sample	f_1	f_2	g	k	b_{cis}	b_{trans}	c_{cis}	c_{trans}
	CH in furan ring	CH in furan ring	CH ₂ in BDO	CH ₂ in BDO	CH in cis-CBDO	CH in trans-CBDO	CH ₃ in cis-CBDO	CH ₃ in trans-CBDO
PBF	7.20	-	4.61	2.08	-	-	-	-
PBTF10-18	7.20-7.23	7.24-7.26	4.61	2.08	4.53	4.70	1.07,1.24	1.14




Figure S3 Chemical structures of EFE, EFT, TFT and the peak assignment in ^{13}C -NMR spectra




Figure S4 TGA curves for PEF, PPF, PBF, PETF-18, PPTF-18 and PBTF-18