



Article Fungal Diseases in Two North-West Spain Vineyards: Relationship with Meteorological Conditions and Predictive Aerobiological Model

Jose A. Cortiñas Rodríguez ¹, Estefanía González-Fernández ¹, María Fernández-González ^{1,2,*}, Rosa A. Vázquez-Ruiz ³ and María Jesús Aira ⁴

- ¹ Department of Plant Biology and Soil Sciences, Sciences Faculty of Ourense, University of Vigo, As Lagoas s/n. 32004 Ourense, Spain; jcortinas@uvigo.gal (J.A.C.R.); estefaniaglezf@gmail.com (E.G.-F.)
- ² Earth Sciences Institute (ICT), Pole of the Faculty of Sciences University of Porto, 4169-007 Porto, Portugal
- ³ Department of Botany, Higher Polytechnic School, University of Santiago of Compostela, 27002 Lugo, Spain; rosana.vazquez@usc.es
- ⁴ Department of Botany, Biology Faculty, University of Santiago of Compostela, 15782 Santiago of Compostela, Spain; mariajesus.aira@usc.es
- * Correspondence: mfgonzalez@uvigo.es

Received: 19 December 2019; Accepted: 28 January 2020; Published: 3 February 2020

Abstract: Grey mould, powdery mildew, and downy mildew are the most frequent fungal diseases among vineyards worldwide. In the present study, we analysed the influence of the fungi causing these diseases (*Botrytis, Erysiphe*, and *Plasmopara*, respectively) on two viticulture areas from Northwestern (NW) Spain during three growth seasons (2016, 2017, and 2018). The obtained results showed the predominant concentration of the *Botrytis* airborne spores, mainly from the beginning of the Inflorescence emerge phenological stage (S-5) until the end of the Flowering phenological stage (S-6). *Erysiphe* and *Plasmopara* airborne spore peak concentrations were more localised around Flowering (S-6) and Development of fruits (S-7) phenological stages. We applied a Spearman's correlation test and a Principal Component Analysis to determine the influence of the variables with the highest correlation coefficient, we developed multiple regression models to forecast the phytopathogenic fungal spore concentrations. The *Botrytis* model regression equation explained between 59.4–70.9% of spore concentration variability. The *Erysiphe* equation explained between 57.6–61% and the *Plasmopara* explained between 39.9–55.8%. In general, we found better prediction results for mean daily concentrations than sporadic spore peaks.

Keywords: Botrytis; Erysiphe; Plasmopara; vineyards; incidence; multiple linear regression

1. Introduction

Among the cryptogamic diseases, the fungal grey mould, powdery mildew, and downy mildew epidemics have the highest incidence in European vineyards [1–4]. Grey mould caused by *Botrytis spp.* Pers.: Fr. (teleomorph: *Botryotinia fuckeliana* (de Bary) Whetzel) affects more than 200 plants, mainly dicot plants, and it can cause serious economic damage, especially in grapevines [5–7]. In addition to an important crop yield loss, this disease can also reduce the wine quality by providing an unstable colour, oxidative damages, premature aging, unpleasant flavours, and clarification difficulties [8]. The *Botrytis* infections, due to the fungal laccase enzyme oxidation, can compromise both the grapes and wine quality [9]. This fungus overwinters as sclerotia and mycelium in buds or trunk cracks, infecting plants again in the following spring mainly through wounds caused by hailstones, mechanical injuries or insects [10–12].

Powdery mildew caused by *Erysiphe necator* Schwein.: (anamorph: *Oidium tuckeri* Berk) came from the United States to Europe through England in 1845, spreading to the entire Mediterranean region and other geographical areas [13]. The American native grapevine has a high resistance level against this pathogen, while the European species *Vitis vinifera* is susceptible as it can present severe disease symptoms in different plant parts [14]. This fungus notably decreases the grape yield and quality, altering the must and wine organoleptic characteristics as it reduces the soluble solids and increases the total acidity [15]. In red wines, infected fruits at the beginning of ripening reach a lower phenolic compounds content, which also has an impact on their sensory properties [16].

Finally, *Plasmopara viticola* (Berk. & Curtis) Berl. & De Toni, responsible for downy mildew, is an obligate oomycete that overwinters as oospores in soil leaves and vegetal debris. These propagules mature as a function of temperature and precipitation, causing subsequent secondary infections. *Plasmopara* is native to North America and was first detected in Spain in 1878. Since then, it has been considered as one of the worst grapevine diseases that occur during favourable weather conditions [17]. The fungus bears fruit only on the host plant surface, as such, it can be only diagnosed by observing sporulation or fruiting on the plant [18]. It causes direct inflorescence and bunch losses or indirect decreases of photosynthetic activity on affected leaves. In severe attacks, the plants suffer partial desiccation resulting in the premature falling of leaves, and the withering of branches [19].

The incidence of these three vine pathogens was widely studied, as well as considering the analysis of cultural control practices. Different measures, such as drainage improvement, green pruning near to flowering, defoliation near to veraison for the improvement of ventilation for bunches, and the use of a training system to achieve appropriate air circulation, can reduce the pathology severity caused by any of these fungi [20–22]. The plant vigour can be controlled by avoiding excessive growth and vegetative development is also beneficial to reduce the incidence of phytopathogens [23]. Regarding chemical control, contact products, such as copper in the form of wettable powder and applied in a foliar spray or sulphur as a powder for dusting, should be used as infection prevention systems, although their use should be avoided during flowering as it can affect fruit set [24]. However, when diseases are evident it is necessary to apply penetrating and systemic treatments with curative effects, regulating their frequency according to the pathogen's intensity [25].

The main goals of this research work are to analyse the incidence of these three fungal grapevine diseases (grey mould, powdery mildew, and downy mildew) in North-western (NW) Spain vineyards and to develop prediction models to detect the airborne spore presence that fungi produce. Besides cost reduction in grape production, this kind of study also impacts environmental protection and quality since it allows a phytosanitary treatment application when a real infection is detected [26,27].

2. Materials and Methods

2.1. Location and Climatic Characteristics of the Study Area

The present study was conducted in two wine-making regions, Ribeiro and Ribeira Sacra, in the Cenlle (117 m a. s.l. 42°18′55.7″ N; 8°6′2.54″ W) and O Mato (332 m a. s.l. 42°30′32.3″ N; 7°30′3.02″ W) vineyards, respective ly (Figure 1) during 2016, 2017 and 2018.



Figure 1. Location of the Ribeiro (Cenlle) and Ribeira Sacra (O Mato) Designation Origin areas in the Northwestern Spain.

The prevailing cultivar in the Cenlle vineyard is Godello, and in O Mato vineyard it is Mencía, both included in the preferential category according to the Designation of Origin Ribeiro (DOG N° 149, 2009) and Ribeira Sacra (DOG N° 194, 2009) regulations. The Godello cultivar has a high vigour and an erect bearing (tending to grow more vertically than horizontally) with an early sprout and maturation. Considering the pathogen susceptibility of the plant according to the three categories established in the classification of the Godello cultivar is classified as having low sensitivity to *Botrytis*, medium sensitivity to *Plasmopara*, and sensitivity to *Erysiphe*. The Mencía cultivar has a medium vigour with early sprouting and semi-late maturation. This cultivar is sensitive to the three considered fungal pathogens [28].

For climatic characterization, we considered the meteorological data of the maximum, minimum and mean temperature, relative humidity, daylight (sunshine) hours, rainfall and wind-speed variables (Table 1), provided by two stations located very close to the studied vineyards, the Leiro and Monforte stations belonging to the weather service Meteogalicia https://www.meteogalicia.gal (accessed on 21/06/2019).

		Ribeiro			Ribeira Sacra		
	Year	2016	2017	2018	2016	2017	2018
	Max T (°C)	22.8	23.9	23.7	20.6	21.9	21.4
Ammunal Awaraga	Min T (°C)	7.3	6.4	8.2	6.7	6.1	7.3
Meteorological Data	Mean T (°C)	13.8	13.9	15.2	12.8	13.1	13.7
	Mean RH (%)	78.4	75.8	75.9	80.9	76.9	78.6
	Sunshine (hours)	5.3	5.9	4.9	8.1	8.4	8.4
	Wind-Speed (Km/h)	2.1	1.8	2.2	2.1	1.8	2.5
Annual Total	Rainfall (L/m ²)	1211.8	814.5	775.2	940.4	616.2	660.5
Maximum	Daily rainfall (l/m ²)	49.0	100.8	39.4	42.8	77.5	25.7
	Date	10 Jan	10 Dec	28 Feb	10 Jan	10 Dec	11 Mar

Table 1. Meteorological parameters, maximum temperature, minimum temperature, average temperature, average relative humidity, sunshine, wind speed and rainfall, in the study area (2016–2018).

Maximum temperature (Max T), minimum temperature (Min T), average temperature (Mean T), average relative humidity (Mean RH).

2.2. Fieldwork and Laboratory Analysis

We studied the phenological stages for both grapevine cultivars following the [29] scale adopted by the BBCH (*Biologische Bundesanstalt, Bundessortenamt and Chemical industry*) as a standardized scale. Phenological observations were applied on 10 vines of each cultivar and this information was correlated with atmospheric spore (or sporangia for *Plasmopara*) concentrations.

The study was carried out during the active grapevine cycle, from the 1st of March to the grape harvest date in the month of September. We used a volumetric method by means of a Hirst type Lanzoni VPPS-2000[®] (Lanzoni s.r.l., Bologna, Italy) pollen-spore trap [30] for the sampling of airborne reproductive structures. The volumetric traps were placed at a height of two meters, according to the vine's foliar arrangement on each studied vineyard. We identified and quantified the three considered pathogens spores/sporangia (*Botrytis, Erysiphe* and *Plasmopara*) in the collected samples. We followed the Spanish Aerobiology Network (REA) proposed protocol [31] for spore count and sample preparation. Slide counts were performed on two longitudinal transects along the slides and the airborne fungal spore concentrations were expressed as a daily average of fungal spores/m³ of air recommended terms were used for the aerobiological terminology [32].

Table 2 shows the phytochemical treatments applied in the plots of the study. In the Ribeiro D.O: (Cenlle), a mean of three treatments/year against *Botrytis* was applied from stage 6 (flowering), six treatments/year against Mildew + Oidium mainly from stage 5 (inflorescence emerges), and two specific treatments/year against Mildew in the 7th and 8th stages (development of fruits and ripening of berries). In the Ribeira Sacra D.O. (O Mato), no treatments against *Botrytis* were applied, four treatments/year (none in 2017) against Mildew + Oidium, mainly in stage 5 (Inflorescence emerge) and four specific treatments/year against Mildew (none in 2017). For all cases, the fungicides used until Flowering (E6) are of the systemic type in years that had a high incidence of fungi, and from this moment of contact (mainly sulphur-based powder products until Fruit set, and copper in wet dust until Softening of berries).

					-		
Against		S 0	S1	S 5	S6	S 7	S 8
	Cenlle 2016				1	2	
	Cenlle 2017				1	1	
Detutio	Cenlle 2018				2	2	1
Botrytis	O Mato 2016						
	O Mato 2017						
	O Mato 2018						
	Cenlle 2016			2	2	2	
	Cenlle 2017			2	1	3	1
	Cenlle 2018		1	1	1	1	2
Mildew + Oldium	O Mato 2016			1	2	1	
	O Mato 2017						
	O Mato 2018			2	1	2	
	Cenlle 2016						1
	Cenlle 2017						2
N (*1.1	Cenlle 2018					2	2
Mildew	O Mato 2016		2				2
	O Mato 2017						
	O Mato 2018	1	1				2

Table 2. Number of phytosanitary treatments applied in the studied plots.

Phenological stages: Bud development (S0), Leaf development(S1), Inflorescence emerge (S5), Flowering (S6), Development of fruits (S7) and Ripening of berries (S8).

2.3. Statistical Analysis

In order to determine the influence of the main meteorological variables on airborne spore concentrations, we applied Spearman's correlation test, as the considered variables didn't show a normal distribution. We considered the correlations for the same day and the previous 1–7 days as spore production can be directly affected by meteorological conditions or indirectly through its effect on the colonised substrates [26]. Moreover, we applied a Principal Component Analysis (PCA) to complement the analysis of these variables, considering the meteorological influence of all variables as a whole. For this analysis we considered for both vineyards the maximum, mean, and minimum temperatures; relative humidity; rainfall; and wind-speed meteorological parameters, independent of the vineyard location.

With the obtained results, we developed models based on Lineal Regressions using the meteorological variables with the highest correlation coefficients as estimators of the fungal spore concentrations. For model validation, we carried out an internal validation and we studied the residual scores as they show the differences between observed and predicted data for each analysed year.

3. Results

3.1. Total Spore Concentrations and Spatial Distribution

The active grapevine period slightly varied according to the vineyard location and the considered year. The sprouting phenological phase (S-0) began between mid-March to the beginning of April, while the harvest date occurred between the end of August to mid-September. Within these dates, the collection of aerobiological samples was carried out and the fungal spores were counted.

The *Botrytis* Seasonal Spore Integral (SSIn) was markedly higher than for the other considered pathogens, with a maximum of 49620 spores in the O Mato vineyard in 2018 and SSIn values of 27656 and 17240 spores for the 2016 and 2017 years, respectively. The *Erysiphe* SSIn noted a record in 2016 for both studied vineyards of 17269 spores in Cenlle and 12946 spores in O Mato. *Plasmopara* was the pathogen with the lowest atmospheric presence. The highest SSIn value was recorded in 2018 at the O Mato vineyard with 5656 spores, varying between 1363 for 2017 and 3618 for 2016 SSIn, and from 679 for 2017 to 3605 for 2018 SSIn values in Cenlle (Table 3).

Cenlle	Botrytis	Erysiphe	Plasmopara
Study Period	19	9 Sep 2016 to 18 Sep 2016	5
SSIn	16806	17269	1910
Daily Maximum	637	399	104
Maximum Date	22 Jun	16 May	8 Jun
Study Period	16	Mar 2017 to 30 Aug 201	7
SSIn	15378	3344	679
Daily Maximum	696	423	54
Maximum Date	28 May 2017	26 May 2017	8 May 2017
Study Period	29	Mar 2018 to 10 Sep 2018	3
SSIn	24214	3116	3605
Daily maximum	1210	316	460
Maximum date	7 Jul 2018	4 Jun 2018	7 Jul 2018
O Mato			
Study period	24	Mar 2016 to 30 Sep 2016	6
SSIn	27656	12946	3618
Daily Maximum	1435	597	502

Table 3. SSIn data, daily maximum spore concentration and maximum date for *Botrytis*, *Erysiphe*, and *Plasmopara* in Ribeiro (Cenlle) and Ribeira Sacra (O Mato) DOs during the study period (spores/m³ of air).

Maximum Date	27 May 2016	3 Jun 2016	7 Jul 2016			
Study Period	04 Mar 2017 to 08 Sep 2017					
SSIn	17240	1686	1363			
Daily Maximum	826	108	45			
Maximum Date	4 Jun 2017	19 Jun 2017	3 Jun 2017			
Study Period	07	Apr 2018 to 07 Sep 2018	3			
SSIn	49620	5632	5656			
Daily Maximum	1547	197	345			
Maximum date	9 Jun 2018	29 May 2018	22 Jul 2018			

SSIn - Seasonal Spore Integral.

Taking into account the maximum daily values we also found a clear *Botrytis* spore's dominance with a maximum of 1547 spores/m³ in O Mato on 9 June 2018. The highest daily peaks of *Erysiphe* and *Plasmopara* were recorded during the 2016 season in the O Mato vineyard, with 597 *Erysiphe* spores/m³ on 3 June 2016 and 502 *Plasmopara* spores/m³ on 7 July 2016.

Considering the pathogen relation with the different grapevine phenological phases, we found a considerable variability depending on the pathogen, season, and vineyard location. The *Botrytis* and *Erysiphe* airborne spore presence were almost constantly along the grapevine growth cycle, while the *Plasmopara* sporangia did not show a continuous daily record. For the considered 2016–2018 period, we found 29 days/season average of *Plasmopara* 0-record in Cenlle, and 20 days/season in O Mato.

The highest *Botrytis* incidence was detected from the inflorescence emergence (S-5) phenological stage until the end of flowering (S-6). At the Cenlle plot, the maximum atmospheric *Botrytis* spore peak was recorded in mid-June in 2016, the end of May in 2017, and several spore peaks were detected between the beginnings of June and the end of July in 2018, with a maximum peak on 7 July. At the



Agronomy 2020, 10, 219

Mato plot, the *Botrytis* spore peaks occurred within the May–July period, with the maximum peaks on 27 May 2016, on 4 June 2017, and 9 June 2018. We also found early infections during the leaf development (S-1) phenological stage at the Cenlle vineyard in 2017, and at the O Mato vineyard in 2016 and 2017 (Figure 2).

Figure 2. Phenology, spore concentrations, and phytosanitary treatments applied in the three study years for Ribeiro (Cenlle) and Ribeira Sacra (O Mato) vineyards.

Most of the *Erysiphe* and *Plasmopara* infections coincided with the end of flowering (S-6) and development of fruits (S-7) phenological stages. At the Cenlle vineyard, there were several *Erysiphe* and *Plasmopara* spore peaks detected in the May–July period in 2016, while the maximum *Erysiphe* spore peak was on 26 May 2017 and 4 June 2018, and the maximum *Plasmopara* sporangia peaks were registered at the beginning of May in 2017, and during July in 2018. At the O Mato vineyard, the highest *Erysiphe* spore peaks were also detected in the May–July period in 2016 with a maximum spore peak on 3 June. In 2017 and 2018, the highest daily spore peaks were lower than in 2016, registered on 19 June and 29 May, respectively. The *Plasmopara* highest sporangia peaks occurred at the beginning of July in 2016 and the end of July in 2018, while in 2017 the peak was detected at the beginning of June, with a markedly lower value (Figure 2).

3.2. Analysis of Meteorological Parameters

Taking into account the annual average maximum temperature value, it was higher in Ribeiro (Cenlle) with a 2 °C difference over Ribeira Sacra (O Mato) for the same considered years. The same occurred with minimum and mean temperatures but with the lowest difference in this case, around 1 °C. For both vineyards, Cenlle and O Mato, 2017 was warmer than the other two considered years with a maximum temperature of 23.9 °C and 21.9 °C respectively. Mean and minimum temperatures were higher in 2018, with 15.2 °C and 13.7 °C mean temperatures, and 8.2 °C and 7.3 °C minimum temperatures, respectively. The average relative humidity was higher in Ribeira Sacra than in Ribeiro, reaching its maximum in 2016 with 80.9%. The same happened with the sunshine hours, with the maximum values for Ribeira Sacra found in 2017 and 2018. The rainiest year was 2016 for both Ribeiro (with 1211.8 mm) and Ribeira Sacra (940.4 mm) areas. Nevertheless, the rainiest day was 10 December 2017 for both vineyards (Table 1).

3.3. Statistical Results

The statistical analysis between the spore daily concentrations and the daily values of the main meteorological variables along the grapevine reproductive cycle was conducted, while also considering the daily values of the meteorological variables during the previous 7 days in regards to the presence of spores in the atmosphere of the vineyard. The statistical analysis between the spore concentrations and the main meteorological variables showed that rainfall and relative humidity had a statistically significant influence in most cases. Nevertheless, the influence sign was not so clear for temperature, as we found the same number of positive and negative correlations for maximum and mean temperature. The parameters with the highest influence on each pathogen varied depending on the vineyard location and the study year.

Within the significant *Botrytis* correlations, we found the highest Spearman's r positive coefficients for the *Botrytis* spores obtained four days before (*Botrytis-4*) vs. rainfall, and *Botrytis-4* vs. relative humidity correlations for Cenlle in 2017. In the same Cenlle vineyard in 2018, we found the *Botrytis-1* vs. minimum temperature, and *Botrytis* vs. mean temperature correlations. We also found a high Spearman's r positive coefficient for the *Botrytis-2* vs. relative humidity correlation at the O Mato vineyard in 2016. For the *Erysiphe* airborne spores we found that the highest Spearman's r significant coefficient was negative, corresponding to *Erysiphe-2* vs. minimum temperature, and *Erysiphe-2* vs. mean temperature correlations at Cenlle vineyard in 2016. At the O Mato vineyard, we found also high correlations, positive in this case, between *Erysiphe* vs. minimum temperature, and *Erysiphe* vs. mean temperature in 2017. The strongest correlations found for *Plasmopara* airborne sporangia corresponded to *Plasmopara* vs. minimum temperature, *Plasmopara* vs. mean temperature

positive correlations at the Cenlle vineyard in 2017, and *Plasmopara-6* vs. rainfall, *Plasmopara-2* vs. relative humidity positive correlations at the O Mato vineyard in 2016. However, we also found considerable negative correlations for *Plasmopara-7* vs. maximum temperature and *Plasmopara-7* vs. mean temperature at the O Mato vineyard in 2016 (Table 4)

		2016			2017			2018	
Cenlle	Botrytis	Erysiphe	Plasmopara	Botrytis	Erysiphe	Plasmopara	Botrytis	Erysiphe	Plasmopara
Deinfell	-0.167 *	-0.365 *	0.355 *	0.401 **	0.196 *	0.172 *	-0.190 *	NC	-0.255 **
Kainiali	Botrytis	Erysiphe-7	Plasmopara-2	Botrytis-4	Erysiphe-7	Plasmopara-4	Botrytis-1	IN.5.	Plasmopara
DII	0.382 *	-0.321 *	0.342 *	0.507 **	0.248 **	NC	0.263 **	0.203 **	NC
КП	Botrytis-2	Erysiphe-2	Plamopara-2	Botrytis-4	Erysiphe-5	IN.5.	Botrytis-5	Erysiphe–7	N.5.
Т	0.288 **	0.440 **	-0.335 *	-0.228 **	-0.180 *	-0.300**	0.387 **	NC	0.387 **
1 max	Botrytis	Erysiphe-2	Plasmopara-2	Botrytis-4	Erysiphe-7	Plasmopara-6	Botrytis	IN.5.	Plasmopara
Tania	0.323 **	-0.674 *	0.346 *	NC	NC	-0.411 **	0.634 **	0.197 *	0.489 **
1 min	Botrytis	Erysiphe-2	Plasmopara-2	IN.5.	IN.5.	Plasmopara-7	Botrytis-1	Erysiphe	Plasmopara
Tmesen	0.307 **	-0.666 **	NC	-0.188 *	-0.154 *	-0.362 **	0.556 **	NC	0.480 **
1 mean	Botrytis	Erysiphe-2	IN.5.	Botrytis-4	Erysiphe-5	Plasmopara-6	Botrytis	N.S.	Plasmopara
O Mato									
Deinfell	0.365 **	0.394 **	0.227 **	NC	-0.272 **	NC	0.224 **	0.231 **	NC
Kainiali	Botrytis-5	Erysiphe-6	Plasmopara-6	IN.5.	Erysiphe-2	<i>N.S.</i>	Botrytis-6	Erysiphe-7	N.5.
DLI	0.593 **	0.491**	0.377**	0.224 **	-0.354 **	0.163 *	0.373 **	0.359 **	NIC
КП	Botrytis-2	Erysiphe-6	Plamopara-2	Botrytis-3	Erysiphe-2	Plasmopara-4	B otrytis-7	Erysiphe-6	N.5.
Т	-0.386 **	-0.476 **	-0.309 **	NC	0.483 **	0.186 *	NG	-0.297 **	NC
	Botrytis-7	Erysiphe-7	Plasmopara-7	IN.3.	Erysiphe	Plasmopara	IN.3.	N.S. Erysiphe-6	
T	-0.153 *	-0.231 **	-0.177*	0.248 **	0.509 **	0.209 **	0.331 **	NC	NC
1 min	Botrytis-7	Erysiphe-6	Plasmopara-7	Botrytis	Erysiphe	Plasmopara	Botrytis	IN.5.	N.5.
Tanaaa	-0.367 **	-0.439 **	-0.332 **	0.224 **	0.582 **	0.218 **	0.169 *	-0.251 **	NC
T mean	Botrytis-7	Erysiphe-7	Plasmopara-7	Botrytis	Erysiphe	Plasmopara	Botrytis	Erysiphe-6	IN.5.

Table 4. Spearman's rank correlation of different parameters for the studied cultivars. Spore concentrations (*Botrytis, Erysiphe* and *Plasmopara*) and the main meteorological parameters. (plevel: * < 0.05; ** < 0.01; N.S.: not significative).

The Principal Component Analysis (PCA) resulted in the extraction of three Principal Components (PC) for both Cenlle and O Mato vineyards. The accumulated explained variance of the original data in Cenlle was 70.9% and 67.6% in O Mato (Table 5).

		CENLLE			O MATO	
Principal Components	PC1	PC2	PC3	PC1	PC2	PC3
Self-value	3.298	1.956	1.135	3.184	1.807	1.098
Variance (%)	36.640	21.729	12.610	35.376	20.087	12.207
Accumulated percentage	36.640	58.369	70.980	35.376	55.463	67.671
Botrytis	-0.003	0.227	0.846	0.196	0.414	0.624
Erysiphe	0.012	-0.001	0.504	-0.065	0.011	0.723
Plasmopara	-0.008	0.076	0.809	0.088	-0.109	0.650
Max T	-0.800	0.512	0.015	0.796	-0.518	-0.043
Min T	-0.158	0.826	0.232	0.848	0.047	0.137
Mean T	-0.643	0.719	0.135	0.901	-0.381	0.047
RH	0.937	-0.037	0.149	-0.185	0.881	0.186
Rainfall	0.805	0.152	-0.082	-0.037	0.817	-0.137
Wind speed	-0.132	-0.697	-0.018	-0.612	-0.075	-0.024

Table 5. Principal Component Analysis, principal component one, principal component two and principal component three. Confidence interval 95% (p < 0.05).

Principal Component Analysis (PCA), principal component one (PC1), principal component two (PC2) and principal component three (PC3).

In both analyzed areas, the Principal Components PC1 and PC2 included meteorological variables and the PC3 grouped the three phytopathogenic fungi spore concentrations. For Cenlle, PC1 includes the maximum and mean temperatures, humidity, and rainfall and PC2 includes the minimum temperature and wind speed. In the case of O Mato, PC1 include the temperatures and the wind speed, whereas PC2 includes the water-related parameters. Most of the variables had significant positive loads, except wind speed (PC2 in Cenlle and PC1 in O Mato). For graphic representation, we selected PC2 versus PC3 for the Cenlle vineyard, and PC1 versus PC3 for the O Mato vineyard (Figure 3). In the obtained charts we showed in detail the relation between fungal airborne spores and the main meteorological variables. In both vineyards we observed a high positive association degree between meteorology and fungal spores counts.



Figure 3. Graphical representation of the Principal Components Analysis (PCA). **a**: PC2 vs. PC3 in Cenlle, **b**: PC1 vs. PC3 in o Mato.

3.4. Predictive Models

The selected variables for the *Botrytis, Erysiphe* and *Plasmopara* spore concentration predictive models development carried out in this study were different for each fungal type, although mean temperature of the same day, relative humidity of the same day and three days before, and spore concentration of the previous day were the most used variables. The *Botrytis* model linear regression equations explained the spore concentration variability of 59.4% in O Mato and 70.9% in Cenlle. For the *Erysiphe* model, the regression equations explained the data variability of 57.6% in O Mato and 61% in Cenlle, and, for the *Plasmopara* model, they explained the data variability of 39.9% in Cenlle and 55.8% in O Mato (Table 6).

Table 6. Botrytis, Erysiphe, and Plasmopara spore concentrations regression models.

Cenlle	Beta	В	Std. Error B	t	<i>p</i> <
Botrytis	R= 0.843	$R^2 = 0.710$	Adjusted R ² = 0.709		
Intercepted		-197.800	42.778	-4.624	0.000
Botrytis-1	0.768	0.768	0.027	28.068	0.000
T mean	0.124	3.650	0.888	4.111	0.000
RH-3	0.144	2.247	0.468	4.800	0.000
Erysiphe	R= 0.782	$R^2 = 0.611$	Adjusted R ² = 0.610		
Intercepted		10.038	2.528	3.971	0.000
Erysiphe-1	0.782	0.782	0.028	28.199	0.000

Plasmopara	R= 0.633	$R^2 = 0.401$	Adjusted R ² = 0.399		
Intercepted		12.016	3.871	3.105	0.000
Plasmopara-1	0.620	0.620	0.035	17.851	0.000
Wind Speed-1	-0.071	-3.278	1.609	-2.037	0.042
O Mato					
Botrytis	R= 0.722	$R^2 = 0.596$	Adjusted R ² = 0.594		
Intercepted		-136.199	63.573	-2.142	0.033
Botrytis-1	0.734	0.827	0.035	23.587	0.009
RH-3	0.082	2.361	0.895	2.637	0.000
Erysiphe	R= 0.760	$R^2 = 0.578$	Adjusted R ² = 0.576		
Intercepted		-54.391	17.578	-3.094	0.002
Erysiphe-1	0.737	0.737	0.029	25.346	0.000
RH-3	0.107	0.885	0.240	3.680	0.000
Plasmopara	R= 0.480	$R^2 = 0.560$	Adjusted R ² = 0.558		
Intercepted		-24.085	12.483	-1.929	0.054
Plasmopara-1	0.742	0.742	0.029	25.282	0.000
RH-1	0.070	0.402	0.170	2.370	0.018

The Beta coefficient is the standardized regression coefficient (all variables standardized to a mean of 0 and a standard deviation of 10). The B coefficient is the unstandardized regression coefficient and S.E. B is its standard error, t value measures the relationship between the coefficient and its standard error, and p value is a probability that measures the evidence against the null hypothesis.

In order to select the meteorological variables for model development we took into account the statistical results. The relative humidity of three days before and the spore concentration of the previous day for each fungal type were the highest significant parameters. For most cases, we obtained a good fit between real and forecast values (Figure 4). The greatest predictive difficulty was focused on the maximum spore peaks, such as *Botrytis* in 2018 at both vineyards, *Erysiphe* in 2017 at Cenlle vineyard, and *Plasmopara* in 2016 at O Mato vineyard, since the models failed to reach the exact real values.



Figure 4. Observed and predicted *Botrytis, Erysiphe,* and *Plasmopara* spore concentrations during the 2016–2018 period, in both vineyards (Cenlle and O Mato).

4. Discussion

The chemical fungicide application following preset treatment schedules is a widely used control strategy among wine growers to reduce the fungal disease impact on the crop, which has several environmental repercussions [33]. Pathogens have a great impact on global agriculture, especially in viticulture as climate change can aggravate this situation [34]. Aerobiological models associated with climatic variables and the use of local cultivars, adapted to the specific area conditions, would enhance the control of grey mould, powdery mildew, and downy mildew [35,36]. Furthermore, these models are useful optimization tools for wine growers to achieve more effective pest management and crop protection as they can predict, in advance, the inoculum concentration of these pathogens [37].

For predictive model's development, we carried out atmospheric monitoring of the *Botrytis*, *Erysiphe*, and *Plasmopara* reproductive airborne structures in two NW Spain vineyards from 2016 to 2018. We simultaneously applied a phenological study on the Godello and Mencía cultivars grown there and we registered the main meteorological parameters values during the study period.

4.1. Relationship with Phenological Stage and Meteorological Influence

Botrytis spore counts in aerobiological samples were markedly higher than *Erysiphe* and *Plasmopara* ones, which coincide with previous studies carried out at nearby vineyards [38], although in this case the airborne *Plasmopara* sporangia atmospheric content (a SSIn of 747 sporangia) was higher than the *Erysiphe* records (a SSIn of 578 spores). In other Southern Spain viticulture areas, the *Erysiphe* spore concentrations exceeded the *Botrytis* concentrations [4], which is related to the much warmer and drier climate and the different cultivars grown there (Pedro Ximénez, Verdejo, Muscat Blanc à Petits Grains and Chardonnay).

Besides the meteorological factors, other aspects influence fungal development, such as the vegetative canopy structure that determines the leaf spatial distribution. Thus, shaded areas are favourable for pathogens development, while those exposed to solar radiation, which results in higher temperatures, inhibit them [39]. Other appropriate management techniques for fungal control for this type of crops include sprouting spacing and length reduction, reducing plant vigour techniques establishing other crops for resource competition, or nutrient intake control especially for nitrogen [14].

The highest *Botrytis* airborne spore abundance that was have found does not mean a higher grey mould incidence for our area's crops, as it depends on other factors such as the grapevine phenological stage at the moments of peak infection (the considered critical period for infection encompasses flowering–maturation) or the applied chemical treatments [14,40]. As we had confirmed, the phytopathological state of the vineyards in the studied wine regions indicates a higher prevalence and damage caused by downy mildew, as it is one of the worst vine diseases occurring alongside favourable climatic conditions [17]. The characteristic high relative humidity values and moderate temperatures of the considered areas favour the survival of *Plasmopara* sporangia [41]. The susceptibility of *Vitis vinifera* to cryptogamic diseases also depends on cultivar and even on the clones used in the crop [42,43].

Several authors had pointed out a constant *Botrytis* spore presence in the atmosphere of the vineyards [44–47]. However, total airborne concentrations vary depending on the vineyard location and the study period. Close to our study area, [26] found SSIn *Botrytis* levels between 1,700–37,299 spores depending on the year. In a Portuguese vineyard located at Amares, [48] found an SSIn level of 800–1,858 *Botrytis* fungal spores, also varying according to the considered study year. In any case, the concentrations were much lower than those recorded in the present study, mainly in the O Mato vineyard during 2018 with an SSIn of 49,620 *Botrytis*. For this year, we also found the highest *Plasmopara* incidence at both vineyards, with a mean value of 4,631 sporangia for 2018 against a mean value of 1,893 sporangia for the rest of the studied years. On the contrary, the *Erysiphe* airborne spore levels stood out in 2016, with a mean SSIn value of 15,108 for both vineyards regarding the other two analysed periods that had a mean SSIn of 3,445.

The vineyard soil management is another influential factor. The pruning remains and the surrounding herbaceous vegetation have an impact on the released spore amount, mainly for *Botrytis* [23,49,50]. This could be one of the explanatory causes for the high *Botrytis* spore concentrations at the O Mato vineyard in 2018, which had a dense vegetation cover throughout the active vine period in that season.

On the other hand, *Botrytis* sporulation has a wider temperature and humidity tolerance range than the other two studied pathogens. Moreover, this fungus can survive in the previous year's dried leaves and mummified bunch grapes [23], which would explain the early infections detected during the leaf development (S-1) phenological stage.

[49] pointed out that grey mould occurs at the beginning of spring and in a more variable way in harvest time (autumn) at different NW Spain wine making regions. Powdery mildew thrives during fruit set, principally during the June and July months, while downy mildew outbreaks are irregular during spring, coinciding with Flowering but also occurring in autumn, when the grape harvest takes place.

Our results placed the highest *Botrytis* airborne spore concentration since the beginning of inflorescence emergence (S-5) until flowering (S-6), which agrees with other studies conducted at

vineyards close by [26]. Despite this, the *Botrytis* incidence may endanger the fruit ripening as inoculum introduction can occur prior to these phenological stages [51]. Grey mould is able to infect vines at any stage [52], but flowering (S-6) may be more vulnerable since pollen release processes and sugary exudates present in flowers enhance the fungal colonization of vegetal tissues [53,54]. Some authors indicated that *Botrytis* conidia germination could occur at 13 °C [10], which explains the early infections that we detected during leaf development (S-1) at both vineyards in 2017. However, we detected that the highest *Botrytis* spore concentrations coincided with higher temperatures, around 20 °C and slightly rainy days, therefore within the optimal temperature range for infection development [45,55,56].

In our study area, the *Erysiphe* and *Plasmopara* infections were later than the *Botrytis*, between the end of flowering (S-6) and development of fruits (S-7). These results agree with findings by other authors [57-59]. Generally, powdery mildew requires relatively dry conditions and moderate temperatures, while downy mildew is greatly affected by wet conditions, either the presence of dew or rainfall [60]. According to [16], temperatures above 15 °C favour Erysiphe development and spread, with an optimal temperature of 20–27 °C. A certain humidity degree is enough for spore germination and excessive rainfall has an inhibitory effect [61–63]. This explains the higher *Erysiphe* incidence in dry seasons. We found lower *Erysiphe* airborne spore concentrations in the rainiest days, which is possibly related to the leaves surface free water detrimental effect on the fungus development [64]. Temperature also determines the Plasmopara maturation period duration, with an optimal temperature of 20-25 °C [17]. In our studied vineyards, we registered an important temperature increase that coincided with high relative humidity at the beginning of July in 2017 and 2018, which could explain the development of several primary and secondary Plasmopara infection cycles. For the analysed vineyards with the maximum downy mildew record levels (Cenlle and O Mato in 2018, and O Mato 2016) we observed that the Hydrothermal Index, which relates the mean temperature and rainfall (Tm.mm), ranged between 4151 Tm.mm and 4376 Tm.mm in Cenlle, approximated to high Plasmopara infection risk cited values [65].

4.2. Predictive Models

Models are powerful tools that can be applied to agricultural pest management and prevention by analyzing an organism's reply to environmental conditions and by improving epidemiological studies to achieve considerable crop protection optimization [37]. Several predictive models that consider the environmental conditions influence on the spore release process have been developed for many important viticulture regions, but they cannot be applied to areas with different climatic characteristics [66]. The obtained models for the prediction of *Botrytis, Erysiphe*, and *Plasmopara* spore concentration showed accurate results, explaining high spore concentration variability.

Knowledge advancement concerning phytopathological fungi epidemiology has led to vineyard disease control improvement [67]. From this point of view, the local monitoring of airborne spores is a valuable element for the predictive model development of fungal propagation. This allows the application of the phytosanitary treatments when real infection risk is detected, which increases environmental protection, and provides added value for wine growers and food safety for consumers.

5. Conclusions

The combination of meteorological and aerobiological parameters is a useful forecast tool for fungal spore concentrations, therefore, for the identification of infection risk moments, providing a strategy that can be integrated into the management of these fungal vine diseases. The total *Botrytis* spore concentration was much higher than the other fungi; however, this was not the most problematic fungus in the considered wine-growing areas. Downy mildew, caused by *Plasmopara*, had a higher prevalence and crop damages, since the high humidity and moderate temperatures in the studied viticulture areas favour its development and high incidence; it is considered as one of the worst vine diseases for the considered wine-growing regions.

Author Contributions: Investigation, Jose Antonio Cortiñas Rodríguez; Methodology, Estefanía González Fernández; Writing-original draft, Rosa A. Vázquez–Ruiz; Writing–review & editing, María Fernández-González and María Jesús Aira. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Acknowledgments: The authors thank to the Xunta de Galicia (Consellería de Educación, Universidade e Formación Profesional) the financial support for the recognition as Grupo de Referencia Competitivo de Investigación (GRC GI–1809 BIOAPLIC, ED431C 2019/07), the Agrupación Estratégica de Investigación BioReDes (ED431E 2018/09), the CITACA Strategic Partnership ED431E 2018/07 (Xunta de Galicia, Spain) and the grant number AGL2014-60412-R of the Economy and Competence Ministry of Spain Government. González–Fernández E. is supported by the FPU Formación de Profesorado Universitario grant from Ministerio de Ciencia, Innovación y Universidades.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Ciliberti, N.; Fermaud, M.; Roudet, J.; Languasco, L.; Rossi, V. Environmental effects on the production of *Botrytis cinerea* conidia on different media, grape bunch trash, and mature berries. *Aust. J. Grape Wine Res.* **2016**, 22(2), 262–270.
- Bois, B.; Zito, S.; Calonnec, A. Climate vs grapevine pests and diseases worldwide: the first results of a global survey. OENO One 2017, 51(3), 133–139.
- 3. Boso, S.; Gago, P.; Santiago, J.L.; de la Fuente, M.; Martínez, M.C. Factors Affecting the Vineyard Populational Diversity of *Plasmopara viticola*. *Plant Pathol*. *J* **2019**, 35(2), 125–136.
- 4. Martínez-Bracero, M.; Alcázar, P.; Velasco-Jiménez, M.J.; Galán, C. Fungal spores affecting vineyards in Montilla-Moriles southern Spain. *Eur. J. Plant Pathol.* 2019, 153(1), 1–13.
- 5. Hidalgo, L. Tratado de viticultura general. 3ª Edición. Ed. Mundi–Prensa. Madrid (España), 2002; pp. 983.
- Williamson, B.; Tudzynski, B.; Tudzynski, P.; Kan, J.A.L.V. *Botrytis cinerea*: the cause of grey mould disease. *Mol. Plant Pathol.* 2007, 8, 561–580.
- Kassemeyer, H.H.; Berkelmann-Löhnertz, B. Fungi of grapes, 61–87. In: Biology of microorganisms on grapes, in must and in wine. Eds, Berlin Heidelberg: Springer, 2009; pp. 61–87.
- Pszczolkowski, P.; Latorre, B.A.; Ceppi Di Lecco, C. Efectos de los mohos presentes en uvas cosechadas tardíamente sobre la calidad de los mostos y vinos Cabernet Sauvignon. *Cienc. Investig. Agrar.* 2001, 28(3), 157–163.
- Aleixandre, J.L.; Giner, J.F.; Aleixandre-Tudó, J.L. Evaluación del efecto terroir sobre la calidad de la uva y el vino (I). *Enovinicultura* 2013, 20, 1–11.
- 10. Coertze, S.; Holz, G.; Sadie, A. Germination and establishment of infection on grape berries by single airborne conidia of *Botrytis cinerea*. *Plant Dis.* **2001**, 85, 668–677.
- 11. Elmer, P.; Michailides, T. Epidemiology of *Botrytis cinerea* in Orchard and Vine Crops. Biology, Pathology and Control, **2007**; pp. 243–272.
- 12. Evans, K.J. Overview of R&D for managing *botrytis* bunch rot in Australia. In: Proceedings of the Australian Society of Viticulture and Oenology Seminar on Grapevine Pests and Disease "Breaking the mould—a pest and disease update", **2008**, pp. 4–15.
- Dufour, M.C.; Lambert, C.; Bouscaut, J.; Mérillon, J. M.; Corio–Costet, M.F. Benzothiadiazole–primed defence responses and enhanced differential expression of defence genes in *Vitis vinifera* infected with biotrophic pathogens *Erysiphe necator* and *Plasmopara viticola*. *Plant Pathol*. 2013, 62, 370–382.
- Grove, G.G.; Moyer, M. Podredumbre por *Botrytis* en la uva para producción comercial en Washington: Biología y manejo de la enfermedad. Washington State University FS046ES, 2015; pp. 1–5.
- Gadoury, D.; Seem, R.; Wilcox, W.; Henick-Kling, T.; Conterno, L.; Day, A.; Ficke, A. Effects of diffuse colonization of grape berries by *Uncinula necator* on bunch rots, berry microflora and juice and wine quality. *Phytopathology* 2007, 97, 1356–1365.
- Oriolani, E.J.A.; Moschini, R.C.; Salas, S.; Martínez, M.I.; Banchero, S. Weather-based models for predicting grape powdery mildew (*Uncinula necator* (Schwein) Burrill) epidemics. *Revista de la Facultad de Ciencias Agrarias, Universidad Nacional de Cuyo* 2015, 47 (2), 197–211.

- 17. Barrios, G.; Reyes, J. "Modelización del mildiu en la vid". Phytoma 2004, 164, 1-9.
- Cambra, M.; Bernal, I. Plasmopara viticola (Berk. y Curtis) Berl. & de Toni. Mildiu de la vid. Fichas de diagnóstico en laboratorio de organismos nocivos de los vegetales. Ficha 64. 2ª edición. MAPA. Madrid. 2006.
- Rossi, V.; Caffi, T. Effect of water on germination of *Plasmopara viticola* oospores. *Plant Pathol.* 2007, 56, 957–966.
- Díaz, T.; Riquelme, A. Control Integrado en el cultivo de uva de mesa en la Región de Murcia. *Phytoma* 2012, 239, 48–54.
- Lucas, A. Control integrado de plagas y enfermedades en el viñedo. Dossier gestión integrada. Vida Rural 2012, pp. 46–51.
- 22. Merlet, H.; Navarro, A.; Rosales, J. Manual técnico productivo y económico Vid. 2016, (Pub. CIREN N° 193) bibliotecadigital.ciren.cl
- González–Domínguez, E.; Caffi, T.; Ciliberti, N.; Rossi, V. A mechanistic model of *Botrytis cinerea* on grapevines that includes weather, vine growth stage, and the main infection pathways. *PLoS One* 2015, 10, 1–23.
- Porras, A. Mejora de la tecnología de la pulverización de productos fitosanitarios sobre plantaciones de vid en espaldera. Doctoral Thesis. Universidad de Córdoba. 2006.
- Rojas, V.; Dennis, M. Nuevas alternativas de control para el oídio de la vid (*Uncinula necator* Schw.). Universidad de Chile. Fac. de CC. Agronomicas. 2003, pp.58.
- Rodríguez-Rajo, F.J.; Jato, V.; Fernández-González, M.; Aira, M.J. The use of aerobiological methods for forecasting *Botrytis* spore concentrations in a vineyard. *Grana* 2010, 49, 56–65.
- Arafat, K.H. Application of statistical model for forecasting powdery mildew of grapes under Egyptian conditions based on meteorological data. *Int. J. Plant Pathol.* 2015, 6, 48–57.
- MAPAMA. Ministerio de Agricultura y Pesca, Alimentación y Medio Ambiente. Registro de variedades comerciales, 2017. www.mapama.gob.es
- Lorenz, D.H.; Eichorn, K.W.; Bleiholder, H.; Klose, R.; Meier, U.; Weber, E. Phänologische Entwicklungsstadien der Weinrebe (*Vitis vinifera* L. ssp. *vinifera*). Codierung und Beschreibung nach der erweiterten BBCH–Skala. *Vitic. Enol. Sci.* 1994, 49, 66–70.
- 30. Hirst, J.M. An automatic volumetric spore-trap. Ann. Appl. Biol. 1952, 36, 257-265.
- Galán, C.; Cariñanos, P.; Alcázar, P.; Domínguez E. Manual de Calidad y Gestión de la Red Española de Aerobiología. Servicio de Publicaciones. Universidad de Córdoba. 2007, pp. 61.
- Galán, C.; Ariatti, A.; Bonini, M.; Clot, B.; Crouzy, B.; Dahl, A.; Fernandez-González, D.; Frenguelli, G.; Gehrig, R.; Isard, S.; Levetin, E.; Li, D.W.; Mandrioli, P.; Rogers, C.A.; Thibaudon, M.; Sauliene, I.; Skjøth, C.; Smith, M.; Sofiev, M. Recommended terminology for aerobiological studies. *Aerobiologia* 2017, 33, 293– 295.
- Gargallo, P.; García-Casarejos, N. Impactos ambientales y medidas de mitigación en el sector vitivinícola español. E3S Web of Conferences 50, 01029. XII Congreso Internacional Terroir. 2018.
- Gautam, H.R.; Bhardwaj, M.L.; Robitash, K. Climate change and its impact on plant diseases. *Curr. Sci.* 2013, 105 (12), 1685–1691.
- Casanova, J. Situación actual de la Viticultura Ecológica: Técnicas de producción de la uva y productos autorizados. *Vida rural* 2003, 171, 41–45.
- Pérez-Sanz, R.; Manzano, Y.; Santiago, L.; de La Iglesia, G.; Campillo, C.; Alberte, L.; Miranda, J.S.; Juárez, I. Metodología para la Validación de Modelos de Desarrollo asociados al clima para el seguimiento del Mildiu, Oidio y Podredumbre Gris en Viñedos de Castilla y León. Revista ITA. 2008.
- Orlandini, S.; Magarey, R.D.; Park, E.W.; Sporleder, M.; Kroschel, J. Methods of agroclimatology: modeling approaches for pests and diseases. In: Hatfield, J.L., M.V.K. Sivakumar, and J.H. Prueger (eds.). Agroclimatology: linking agriculture to climate. Agron. Monogr. 60. ASA, CSSA, and SSSA, Madison, WI, USA. 2017.
- Fernández-González, M., Rodríguez-Rajo, F.J., Jato, V., Aira, M.J. Incidence of fungals in a vineyard of the denomination of origin Ribeiro (Ourense NW Spain). *Ann. Agr. Environ. Med.* 2009, 16, 263–271.
- Calonnec, A., Cartolaro, P., Naulin, J. M., Bailey, D., Langlais, M. A host-pathogen simulation model: powdery mildew of grapevine. *Plant Pathol.* 2008, 57, 493–508.

- Dugan, F.M., Lupien, S.L., Grove, G.G. Incidence, aggressiveness and in planta interactions of *Botrytis cinerea* and other filamentous fungi quiescent in grape berries and dormant buds in central Washington State. *Phytopathology* 2002, 150, 375–381.
- 41. Gessler, C., Pertot, I., Perazzolli, M. *Plasmopara viticola*: a review of knowledge on downy mildew of grapevine and effective disease management. *Phytopathol. Mediterr.* **2011**, 50, 3–44.
- 42. Galet, P., 2000. Précis de viticulture. 7 ª ed. Imp. JF, Montpellier.
- Boso, S., Santiago, J.L., Villaverde-Alonso, V., Gago, P., Martínez, M.C., Rodríguez, E. Evaluación de la incidencia a enfermedades fúngicas en diferentes clones del cv. Albariño (*Vitis vinifera* L.). *Phytoma* 2009, 210, 1–5.
- 44. Díaz, M.R., Iglesias, I., Jato, M.V. Airborne concentrations of *Botrytis*, *Uncinula* and *Plasmopara* spores in a vineyard in Leiro-Ourense (N.W. Spain). *Aerobiologia* **1997**, 13, 31–35.
- 45. Díaz, M.R., Iglesias, I., Jato M.V. Seasonal variation of airborne fungal spore concentrations in a vineyard of North-West Spain. *Aerobiologia* **1998**, 14, 221–227.
- 46. Oliveira, M., Guerner-Moreira, J., Mesquita, M.M., Abreu, I. Important phytopathogenic airborne fungal spores in a rural area: incidence of *Botrytis cinerea* and *Oidium* spp. *Ann. Agric. Environ. Med.* **2009**, 16, 197–204.
- Rodríguez-Rajo, F.J., Seijo Coello, M.C., Jato, V. Estudio de los niveles de los principales fitopatógenos para la optimización de cosechas de *Vitis vinifera* en Valdeorras Ourense (1998). *Botanica Complutensis* 2002, 26, 121–135.
- Fernández-González, M., Rodríguez Rajo, F.J, Jato, V., Aira, M.J., Ribeiro, H, Oliveira, M. Abreu, I. Forecasting ARIMA models for atmospheric vineyard pathogens in Galicia and Northern Portugal: *Botrytis cinerea* spores. *Ann. Agr. Environ. Med.* 2012, 19, 255–262.
- Martínez, C., Boso, S. Evaluación de la virulencia de distintas poblaciones de hongos responsables del Mildiu, Oídio y Botrytis en distintas denominaciones de origen gallegas. Grupo de Viticultura de la MBG–CSIC (Pontevedra). 2015.
- Cortiñas, J.A., Aira, M.J., Fernández–González, M., Rodríguez-Rajo, F.J., Vázquez–Ruiz, R.A. Potential sustainable wine–growing model in the Ribeira Sacra D.O. (NW Spain). *Cienc. Tec. Vitivinic.* 2018, 33(3), 114–145.
- 51. Latorre, B.A., Rioja, M.E. Efecto de la temperatura y humedad relativa sobre la germinación de conidias de *Botrytis cinerea. Cien. Inv. Agr.* **2001**, 29(2), 67–72.
- Molitor, D., Baus, O., Hoffmann, L., Beyer, M. Meteorological conditions determine the thermal-temporal position of the annual *Botrytis* bunch rot epidemic on *Vitis vinifera* L. cv. Riesling grapes. *OENO One* 2016, 50, 231–234.
- Keller, M., Viret, O., Cole, M. *Botrytis cinerea* infection in grape flowers: defense reaction, latency and disease expression. *Phytopathology* 2003, 93, 316–22.
- Viret, O., Keller, M., Jaudzems, V.G., Cole, F.M. *Botrytis cinerea* infection of grape flowers: Light and electron microscopical studies of infection sites. *Phytopathology* 2004, 94, 850–857.
- 55. Latorre, B.A., Rioja, M.E., Lillo, C. Efecto de la temperatura en el desarrollo de la infección producida por *Botrytis cinerea* en flores y bayas de uva de mesa. *Cien. Inv. Agr.* **2002**, 29(3), 145–151.
- Ciliberti, N., Fermaud, M., Languasco, L., Rossi, V. Influence of fungal strain, temperature, and wetness duration on infection of grapevine inflorescences and young berry clusters by *Botrytis cinerea*. *Phytopathology* 2015, 105(3), 325–333.
- 57. Calonnec, A., Cartolaro, P., Poupot, C., Dubourdieu, D., Darriet, P. Effects of *Uncinula necator* on the yield and quality of grapes (*Vitis vinifera*) and wine. *Plant Pathol.* **2004**, 53, 434–445.
- 58. Campbell, P., Bendek, C., Latorre, B.A. Risk of powdery mildew (*Erysiphe necator*) outbreaks on grapevines in relation to cluster development. *Cien. Inv. Agr.* **2007**, 34 (1), 5–11.
- Fernández-González, M., Piña-Rey, A., González-Fernández, E., Aira, M.J., Rodríguez-Rajo, F.J. First assessment of Goidanich Index and aerobiological data for *Plasmopara viticola* infection risk management in north-west Spain. *J. Agr. Sci.* 2019, 157, 129–139.
- Thind, T.S., Arora, J.K., Mohan, C., Raj, P. Epidemiology of powdery mildew, downy mildew and anthracnose diseases of grapevine. In Diseases of Fruits and Vegetables Volume I (pp. 621–638). Springer, Dordrecht. 2004.
- Chellemi, D.O., Marois, J.J. Effect of fungicides and water on sporulation of *Uncinula necator*. *Plant Dis*. 1991, 75, 455–457.

- 62. Sivapalan, A. Effects of impacting rain drops on the growth and development of powdery mildew fungi. *Plant Pathol.* **1993**, *42*, 256–263.
- Jarvis, W.R., Gubler, W.D., Grove, G.G. Epidemiology of Powdery Mildews in Agricultural Pathosystems. In: R. R. Bélanger, W. R. Bushnell, A. J. Dik, T. L. W. Carver (Eds.): The Powdery Mildews: a Comprehensive Treatise. American Phytopathological Society, APS Press. St. Paul, Minnesota, USA. 2002, pp. 169–199.
- 64. Campbell, P. Efecto de factores ambientales y métodos de control sobre la germinación y desarrollo de *Uncinula necator* en *Vitis vinifera*. Tesis de grado. Santiago, Chile. Universidad Católica de Chile, Facultad de Agronomía e Ingeniería Forestal. **2003**.
- 65. Almendro, J.P. Índices climáticos propios de la vid en el sector central de tierra de barros. Centro Universitario Santa Ana. Universidad de Extremadura, **2013**, pp. 121–131.
- Thiessen, L.D., Thiessen, L., Neill, T.M., Mahaffee, W.F. Assessment of *Erysiphe necator* ascospore release models for use in the Mediterranean climate of Western Oregon. *Plant Dis.* 2018, 102 (8), 1500–1508.
- 67. Magarey, P.A., Magarey, R.D., Emmett, R.W. Principles for managing the foliage diseases of grapevine with low input of pesticides. In: 6th International Congress on Organic Viticulture (p. 140). Print-Online. **2000**.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).