

Article

Floral and Pollen Traits of *Moringa oleifera* Lam. and *Moringa peregrina* (Forssk.) Fiori Provide Reproductive Adaptations for Arid Conditions

Yiftach Vaknin ^{1,*}, Dan Eisikowitch ² and Adina Mishal ³
¹ Department of Natural Resources, Institute of Plant Sciences, Agricultural Research Organization, The Volcani Center, 68 HaMaccabim Road, P.O. Box 15159, Rishon LeZion 7505101, Israel

² Department of Plant Sciences, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv 69978, Israel; dane@tauex.tau.ac.il

³ Faculty of Science, Kibbutzim College of Education, Tel Aviv University, 149 Derech Namir, Tel Aviv 6250769, Israel; adinamish@gmail.com

* Correspondence: yiftachv@volcani.agri.gov.il; Tel.: +972-3-968-3486



Citation: Vaknin, Y.; Eisikowitch, D.; Mishal, A. Floral and Pollen Traits of *Moringa oleifera* Lam. and *Moringa peregrina* (Forssk.) Fiori Provide Reproductive Adaptations for Arid Conditions. *Agronomy* **2021**, *11*, 1090. <https://doi.org/10.3390/agronomy11061090>

Academic Editor: Yehoshua (Shuki) Saranga

Received: 11 May 2021

Accepted: 25 May 2021

Published: 27 May 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Our study attempted to elucidate the significance of floral and pollen traits of the highly nutritious tropical trees *Moringa oleifera* and *Moringa peregrina* for their reproductive success under arid conditions. We found that the pollen grains of both species were immersed in a pollenkitt that constituted ~60% of the pollen. Successful pollination was achieved by large bees inserting the pollen into a narrow stylar tube. We found that, upon removal of the pollenkitt, approximately 65% fewer pollen grains penetrated the stylar tube for both species. The pollenkitt protected against heat and desiccation, while removing the pollenkitt resulted in significantly reduced levels of the viability of pollen grains, especially in *M. oleifera*, and significantly reduced levels of germinability in both species. The stylar tube provided high protection for pollen grains against heat and desiccation even when the pollenkitt was removed. Chemical analysis of pollenkitts of the two species revealed a waxy blend of 21 hydrocarbon compounds, in which *n*-alkanes constituted >90% of the compounds and their identity corresponded to known plant and animal hydrocarbons, associated with protection against heat and water stress. We concluded that, under arid conditions, the reproductive success of both *Moringa* species is potentially enhanced by their unique floral and pollen traits. This supports the prospect of cultivating *M. oleifera* and *Moringa peregrina* as food crops in arid regions across the globe.

Keywords: arid conditions; pollen adhesion; pollen germinability; pollenkitt; pollen viability; reproductive adaptation; water stress

1. Introduction

Moringaceae is a family consisting of 13 known perennial species from a dry tropical origin [1] and certain species, such as *M. oleifera* and *M. peregrina*, can potentially be grown under arid conditions. *Moringa oleifera*, also known as “the miracle tree,” is the most widely distributed and economically important tree species in this family. Its leaves and immature pods are highly nutritious. It is traditionally utilized as a food crop throughout the dry tropics in Asia, Africa, and America, with very little cultivation in arid regions [2]. It is also cultivated as a source of seed oil for food, bioenergy, and cosmetics [3–5]. Various plant parts are used as natural medicine [6,7], and seed powder is used as a coagulant for water purification [8]. *Moringa peregrina* is the second most utilized *Moringa* species [1]. It is a wild tree, highly tolerant to heat and drought [9], originating in the desert along the Red Sea throughout the Arabian Peninsula. *M. peregrina* is a highly nutritious tree grown primarily for human consumption, medicinal purposes, and animal fodder [10]. The seeds of *M. peregrina* provided oil for food and cosmetics in antiquity [11]. However,

its commercial value as a food and medicinal and industrial commodity has rarely been explored [12,13].

M. oleifera is cultivated in tropical climatic regions worldwide, while arid regions are being carefully explored for its cultivation. *M. peregrina*, on the other hand, is well-adapted to arid conditions with <200–300 mm rainfall and is, therefore, considered to be drought-tolerant. Both *Moringa* species require hot and dry conditions with sufficient water supply for growth and reproduction. When grown in arid regions such as the Negev and Arava Deserts in Israel, they have better performances, both vegetatively and reproductively, than when grown under wetter and colder conditions, as long as they are sufficiently irrigated and saturation of the soil with water is avoided by proper drainage [14].

In our previous study, we assessed both *M. oleifera* and *M. peregrina* as potential seed-oil and seed-protein sources under semi-arid, Mediterranean conditions [15]. We found that *M. oleifera* was better suited for oil and protein production than *M. peregrina* under semi-arid conditions. It produced more seeds more predictably than *M. peregrina*. *M. peregrina*, on the other hand, produced larger seeds with higher oil concentrations and was less susceptible to local diseases than *M. oleifera*.

The floral morphologic and chemical traits and the pollination services provided by pollen vectors of *M. oleifera* and *M. peregrina* are vital to their reproductive success. Both *M. oleifera* and *M. peregrina* have zygomorphic and horizontally oriented hermaphroditic flowers with five sepals, five petals, and five unequal stamens surrounding a single pistil (Figure 1) consisting of one open-type stigma and a hollow stylar tube. At maturity, the posterior petal remains erect, while the others are reflexed together with the sepals. The intermediate and front petals are curved forward, bearing short trichomes on their ventral side, and they function as a landing platform for visiting insects [16]. Flower anthesis of *M. oleifera* is associated with a temperature range of 27–29 °C and relative humidity of 68–78%. The flowers are protandrous and herkogamous. At anthesis, the anthers dehisce through longitudinal slits releasing spheroidal pollen grains (~35 µm) while the style remains shorter than the anthers [17]. After ~24 h, the style elongates beyond the length of the anthers, and the stigma becomes receptive for the next 48 h [18]. A similar floral structure and phenological pattern were recorded for *M. peregrina* by Vaknin and Mishal [15].



Figure 1. Open flowers of *M. oleifera* (A) and *M. peregrina* (B). Scale bar is 1 cm.

Pollination and fertilization of *M. oleifera* were studied in India by Bhattacharya and Mandal [19]. They reported that floral anthesis occurred before noon, and the anthers

opened shortly after the anthesis and produced ca. 23,000 pollen grains per flower. The average fruit set was approximately 10% and resulted from successful pollination by visiting insects, primarily carpenter bees (*Xylocopa* sp.). They also clarified that successful pollination requires the insertion of pollen grains into the stylar tube by the visiting insect. In Israel, *M. peregrina* blooms during spring and summer (March–July), and the flowers are visited by a wide range of invertebrates, mainly honeybees (*Apis mellifera*) and yellow carpenter bees (*Xylocopa pubescence*) (Vaknin, personal information). These two *Moringa* species produce pollen that is heavily immersed in a sticky fluid-like pollenkitt. However, comparative floral traits and pollination and fertilization biology of these species have rarely been studied before. Furthermore, the significance of the reproductive traits of the two *Moringa* species for their reproductive success under arid conditions has barely been tested.

Here, we attempted to elucidate the significance of the reproductive traits of *M. oleifera* and *M. peregrina*, associated with the interactions between pollen grains and pollen and floral style, for their reproductive success under arid conditions.

2. Materials and Methods

To elucidate the roles of the pollenkitt outside of the stylar tube in protecting from the hot and dry conditions associated with arid regions, we tested first whether the pollenkitt enabled pollen insertion into the stylar tube. We then tested the effects of the pollenkitt, inside and outside of the stylar tubes, on pollen viability and pollen germinability under extreme temperatures. We also tested the impact of the pollenkitt on pollen germinability under low relative humidity. To provide further insight on the functionality of the pollenkitt under arid conditions, we analyzed the chemical composition of the pollenkitts of *M. oleifera* and *M. peregrina*.

2.1. In Vitro Pollen Viability and Germinability Tests

Samples of flowers and pollen were acquired from a *M. oleifera* and *M. peregrina* plantation at Volcani Center research institute in central Israel (31°59′14.25″ N, 34°49′27.50″ E). Local climatic conditions are hot and dry in summer, and cool and wet in winter, with an average annual precipitation of 524 mm (Israel Meteorological Service). At anthesis, 3–5 fresh *Moringa* flowers were randomly sampled for pollen viability and germinability tests from a *Moringa* plantation at the ARO campus in central Israel. Pollen viability was tested using the 3,3′-diaminobenzidine (DAB) staining technique [20]. The percentage of viable pollen was calculated by observing 100 pollen grains under an Olympus Cx33 light microscope (Southend-on-Sea, UK). Pollen in vitro germinability was evaluated by incubation under a 10% (*w/w*) sucrose solution supplemented with 200 µg/mL H₃BO₃. Pollen was inoculated into micro-wells (*n* = 6) containing 50 µL of solution and incubated at 25 °C and 40% RH for 4 h. Due to the very high number of pollen grains being clumped together, we could not count individual germinating pollen grains, and germinability was evaluated on a scale of 1 to 5, with 1 representing very low germinability and 5 representing very high germinability (Figure 2).




Figure 2. Various levels of in vitro pollen germinability: zero germinability (A), low germinability (B), high germinability (C), and high germinability of clumped pollen (D). The scale bar is 200 μ m.

2.2. Removal of the Pollenkitt

To elucidate the significance of the pollenkitt for the pollen grains, we compared different pollen traits, with and without the pollenkitt. A method to remove the pollenkitt from the pollen grains without harming their viability [21] was followed with minor modification. The pollenkitt was removed by immersing fresh pollen grains in Eppendorf vials filled with 0.5 mL hexane, and the vials were vortexed for 5 min. Then the vials were centrifuged at 10,000 RPM, the liquid was decanted, and the process was repeated as a second wash with hexane. The remaining liquid, covering the pollen grains, was fully evaporated for 1 h. The remaining pollen grains, with their pollenkitt being removed, were then used for the tests of pollen viability, pollen germinability, and pollen insertion compared to the intact ones.

2.3. Test of Pollenkitt for Pollen Insertion into the Styler Tube

First, fresh pollen grains with and without pollenkitt were spread onto a microscope slide in a thin layer. Then the tip of the style of the unpollinated fresh *Moringa* flowers was repeatedly brushed against the pollen grains at a 45° angle until a clump of pollen grains appeared at the tip of the style. Excess pollen grains were removed from the tip of the style. The style was laid on a microscope slide, and the pollen was squeezed out from the styler tube onto the slide with a dissecting needle. The pollen grains were stained using a single drop of Basic Fuchsin, and the numbers of them were counted under the microscope.

2.4. Effects of Temperature on Pollen Viability and Germinability, with or without Pollenkitt, inside the Styler Tube

To test the effects of temperature on in vitro pollen viability and germinability, fresh pollen from the anthers and those after removal of the pollenkitt from the pollen grains were placed in open Eppendorf vials and exposed to various temperatures (-20°C , 4°C , 25°C , and 40°C) for 24 h. These temperatures were arbitrarily selected to cover a broad spectrum of temperatures, from extremely low, through low and moderate, to high ambient temperatures during bloom, these temperatures being -20°C , 4°C , 25°C , and 40°C , respectively. Although not present in real life, -20°C was also added to test for extreme temperatures on the lower part of the spectrum. Following exposure to different ambient temperatures, the viability and germinability of the pollen grains were tested. As a control, fresh pollen, with or without pollenkitt, was also tested for germinability at 25°C .

Similarly, we took fresh pollen from the anthers. After removing the pollenkitt from a portion of the pollen grains, we inserted the pollen, with or without the pollenkitt, into the styler tubes as described above. We placed the styles in open Petri dishes and exposed them to various temperatures (-20°C , 4°C , 25°C , and 40°C) for 24 h. Following exposure to different ambient temperatures, we squeezed out the pollen grains using a dissecting needle and tested them for viability only, due to the relatively low number of pollen grains inserted into the styler tube.

2.5. Test for Germinability under Low RH

To test the effects of relative humidity on in vitro pollen germinability, fresh *M. oleifera* pollen was taken from the anthers, and a portion of it was tested for germinability. Then, the pollenkitt was removed, and they were divided into two groups; one was tested immediately for germinability and the other was placed in a closed chamber with 100% RH for 1 h and was subsequently tested for germinability.

2.6. Observation of Pollen on Anthers, Insects, and inside the Styler Tube

At anthesis, SEM imaging of pollen grains taken from newly opened anthers and gold-coated in a vacuum was carried out using a scanning electron microscope (SEM) (JEOL 840A, Tokyo, Japan). External observations of deposited pollen grains on visiting insects, i.e., honeybees (*Apis mellifera*) and carpenter bees (*Xylocopa pubescens*), were carried out using a Leica DM LB stereomicroscope. Images were acquired using a Leica DC-200 digital camera (Leica Microsystems, Wetzlar, Germany). A confocal microscopy study was carried out on the styles of open-pollinated flowers to view the pollen grains inside the styler tube. The styles were prepared and stained using a modification of the Aniline blue epifluorescence method [22]. Stained styles were placed on a microscope slide and observed by an Olympus 1X-81 (Olympus Co., Tokyo, Japan) inverted laser scanning confocal microscope (Fluoview 500) equipped with a diode 405 nm laser and a 10×0.4 numerical aperture or 20×0.7 numerical aperture UPlanApo objective, at an excitation laser line of 405 nm and emission filter barrier of 430–460.

2.7. Estimation of Weight Ratio of Pollenkitt to Pollen Grain

Fresh pollen from several flowers was placed in an Eppendorf vial and weighed using a scale (Pionner™, Ohaus Corp. Parsippany, NJ, USA) to an accuracy of 0.0001 g. The pollenkitt was removed, as described above, and the pollen grains were reweighed to estimate the percentage of the weight of the pollenkitt.

2.8. GC-MS Chemical Analysis of the Pollenkitt

Hexane extracts of pollen grains (see above, pollenkitt removal) or head-space samples ($n = 3$ for both), collected after exposure for 30 min at 60°C , were analyzed on an Agilent GC-MSD apparatus (Agilent 6890/5977A GC-MS system, Agilent, CO, USA) equipped with an Rtx-5SIL MS (“Restek”) ($30\text{ m} \times 0.25\text{ mm i.d.}$, $0.25\text{ }\mu\text{m}$ film thickness) fused-silica capillary column. Helium at constant pressure (13 p.s.i.) was used as a carrier gas with

retention time locking. Samples of the extracts were injected in splitless injection modes. The injection temperature was 250 °C, and the detector temperature was 300 °C. Column conditions were 50 °C for 2 min, followed by an increase of 15 °C/min to 300 °C, then kept for 15 min at 300 °C. For MS detection, we used an electron ionization system with ionization energy of 70 eV. MS information, covering a mass range from 50–500 atomic mass units, was collected with the full-scan mode. Molecular identification was undertaken via the NIST MS Search v2.0 software, Gaithersburg, MD, USA.

2.9. Statistical Analyses

Statistical analyses were performed with JMP 14.0.0 software (SAS Institute Inc. Cary, NC, USA). A two-tailed *t*-test was applied to compare pollen insertion into the stylar tube, with and without the pollenkitt, for *M. oleifera* and *M. peregrina*, and to compare the pollen grain to pollenkitt weight ratios of *M. oleifera* or *M. peregrina*. Comparisons of the effects of various temperatures on pollen viability, with and without pollenkitt, on the anthers and inside the stylar tube were carried out by two-way ANOVA followed by one-way ANOVA and post hoc Tukey HSD tests, for either *M. oleifera* or *M. peregrina*. Comparisons of the effects of various temperatures on pollen germinability, with or without pollenkitt, were carried out by Kruskal–Wallis tests and post hoc Dunn’s tests for *M. oleifera* and *M. peregrina*. All data are represented as the mean \pm standard error. Values are reported as significantly different if $p < 0.05$.

3. Results

3.1. The Significance of the Pollenkitt for Pollen Insertion into the Stylar Tube

Pollen grains of both *Moringa* species penetrated the stylar tube at relatively large numbers of 337 ± 31.5 and 252 ± 30.5 on average for *M. oleifera* and *M. peregrina*, respectively. When the pollenkitt was removed, less than 65% of pollen grains penetrated the stylar tube for *M. oleifera* (118 ± 15.6 ; $t = 6.24$, $p < 0.0001$) and *M. peregrina* (88 ± 17.6 ; $t = 4.66$, $p < 0.0001$). It was also observed that, without pollenkitt, pollen grains repelled each other and scattered to all directions at the entrance to the stylar tube.

3.2. Effects of Temperatures on Pollen Viability and Pollen Germinability, with and without Pollenkitt

Analysis of two-way ANOVA for *M. oleifera* revealed a significant difference in pollen viability between pollen grains exposed to different temperatures ($p < 0.0001$) and between pollen grains with and without pollenkitt ($p < 0.0001$). A significant interaction was also revealed between the temperatures and the presence and absence of pollenkitt ($p < 0.0001$). The intact pollen grains retained their relatively high viability under the temperatures tested, with the highest level of 93% at 25 °C and the significantly lowest level of 72% at 40 °C (Figure 3A; Tukey HSD $p = 0.0001$). When the pollenkitt was removed, all levels of viability were reduced by 9–15% at -20 °C, 4 °C, and 25 °C and by a drastic 79% at 40 °C, from a viability of 73% to a viability of 15% (Figure 3A). Pollen germinability showed a similar trend, with a more drastic and statistically significant reduction, from nearly maximal levels, under all temperatures for intact pollen grains, to nearly 50% reduction in germinability once the pollenkitt was removed (Figure 4A; Kruskal–Wallis $p < 0.0001$). The most significant reductions in germinability, by almost 60%, were detected for pollen grains with their pollenkitt removed and exposed to 25 °C and 40 °C (Dunn’s tests $p < 0.05$).




Figure 3. Effects of exposure to various temperatures (values in parentheses) on pollen viability of *M. oleifera* (A) and *M. peregrina* (B), with (+) and without (−) pollenkitt. Values are means \pm SE. Different letters indicate significant differences at $p < 0.05$.

Analysis of two-way ANOVA for *M. peregrina* revealed a significant difference in pollen viability between pollen grains exposed to different temperatures ($p = 0.0018$) and between pollen grains with and without the pollenkitt ($p = 0.0002$). However, no significant interaction was revealed between the temperatures and the presence and absence of pollenkitt ($p = 0.1746$). The intact pollen grains retained their relatively high viability under all temperatures tested, with the highest levels of 93%, 95%, and 94% measured at -20 °C, 4 °C, and 40 °C, respectively, and the lowest level of 77% measured at 25 °C (Figure 3B). When the pollenkitt was removed, all levels of viability were reduced. However, more drastic reductions were measured under the temperatures of -20 °C and 25 °C, and much smaller reductions were measured at 4 °C and 40 °C (Figure 3B). Pollen germinability of intact pollen grains was nearly maximal under temperatures of -20 °C and 4 °C and was non-significantly reduced by 27–37% at 25 °C and 40 °C (Figure 4A). When the pollenkitt

was removed, the levels of germinability at $-20\text{ }^{\circ}\text{C}$ and $4\text{ }^{\circ}\text{C}$ were also reduced by 34–37%, while at $40\text{ }^{\circ}\text{C}$ the pollen grains completely lost their ability to germinate (Figure 4A; Dunn's tests $p < 0.05$).




Figure 4. Effects of exposure to various temperatures (values in parentheses) on pollen germinability of *M. oleifera* (A) and *M. peregrina* (B), with (+) and without (−) pollenkitt. Values are means \pm SE. Different letters indicate significant differences at $p < 0.05$.

Thus, while *M. oleifera* pollen grains retained their viability under high temperatures and lost some of their viability when the pollenkitt was removed, the *M. peregrina* pollen grains retained most of their viability at $40\text{ }^{\circ}\text{C}$. However, under this condition, they completely lost their ability to germinate.

3.3. Effects of Extreme Temperatures on Pollen Germinability inside the Styler Tube

Under extreme temperatures, both *M. oleifera* and *M. peregrina* pollen retained most of their viability within the styler tube (Figure 5). At the temperatures of -20°C and 4°C , both *M. oleifera* and *M. peregrina* intact pollen grains had 72–76% viability, while at 25°C and 40°C , they showed higher values of viability of 88% and 93%, respectively. Without the pollenkitt, the viability of the pollen grains within the styler tube was relatively high for both *M. oleifera* and *M. peregrina* under all measured temperatures, ranging from 86–95% for *M. oleifera* (Figure 5A) and 88–100% for *M. peregrina* (Figure 5B).




Figure 5. Effects of exposure to various temperatures (values in parentheses), inside the styler tube, on pollen viability of *M. oleifera* (A) or *M. peregrina* (B), with (+) or without (−) pollenkitt. Values are means \pm SE. Different letters indicate significant differences at $p < 0.05$.

Two-way ANOVA for *M. oleifera* did not reveal significant differences in pollen viability between pollen grains inside the stylar tube and exposed to different temperatures ($p = 0.0806$). Removal of the pollenkitt, however, revealed significant effects on pollen viability inside the stylar tube ($p < 0.0001$), with a slight increase in viability for pollen grains without pollenkitt and exposed at $-20\text{ }^{\circ}\text{C}$ and $4\text{ }^{\circ}\text{C}$ (Figure 5A). A significant interaction was also revealed between temperature and presence of pollenkitt ($p = 0.0159$), suggesting a different effect of temperature depending on whether the pollen grains inside the stylar tube were coated with pollenkitt or not. Analysis of two-way ANOVA for *M. peregrina* did not show significant differences in pollen viability between pollen grains inside the stylar tube and those exposed to different temperatures ($p = 0.6873$). Removal of the pollenkitt significantly affected pollen viability inside the stylar tube ($p < 0.0001$), showing a slight increase in viability for pollen grains exposed to $-20\text{ }^{\circ}\text{C}$, $4\text{ }^{\circ}\text{C}$, and $25\text{ }^{\circ}\text{C}$, and a slight decrease for pollen grains exposed to $40\text{ }^{\circ}\text{C}$ (Figure 5B).

3.4. The Significance of the Pollenkitt for Germinability under Low RH

We found that within several hours of pollenkitt removal from the pollen grains, their germinability was significantly reduced from the highest level of 5.0 to the lowest level of ~ 1.0 (Figure 6; Kruskal–Wallis $p = 0.0004$, post hoc Dunn’s test $p = 0.0003$). However, after 1 h of exposure to high RH, the pollen grains restored most of their germinability to level 4.0 (Figure 6).




Figure 6. Effects of removal of pollenkitt (–) and exposure to high relative humidity for 1 h on pollen germination of *M. oleifera*, with (+) or without (–) pollenkitt. Values are means \pm SE. Different letters indicate significant differences at $p < 0.05$.

3.5. Morphological Analyses on Flowers and Insects and within the Stylar Tube

SEM analysis of *M. oleifera* pollen grains in the anthers revealed that the pollen grains were immersed in a thick grainy substance that was coating them and was also filling the space between them (Figure 7A). Thus, the pollen grains in the anthers were glued together, forming large clumps (Figure 7A). SEM analysis of a pollinated floral style showed the entrance to the stylar tube filled with pollen (Figure 7B). Confocal microscopy of the pollinated style revealed that the stylar tube was filled with pollen grains and only a few of

them protruded beyond the tip of the style (Figure 7C). Microscopic analysis of the heads of the visiting carpenter bees revealed that their eyes and antennae were coated with sticky pollen grains and straight grooves, forming an X shape that appeared between the eyes and antennae of the bees (Figure 7D).




Figure 7. Morphological imaging of *M. oleifera* pollen grains: in a large clump coated with pollenkitt (A—SEM), at the entrance to the styler tube (B—SEM), inside the styler tube (C—confocal microscope), and on the head of a carpenter bee between the antennae (D—stereomicroscope).

3.6. Estimation of Weight Ratio of the Pollenkitt to the Pollen Grains

Weight analysis of the pollenkitt, compared to the weight of the pollen grains, revealed that the majority of the pollen, collected by the visiting insects, was the pollenkitt, constituting approximately 60% of the weight of the pollen ($58.8\% \pm 2.4$ and $59.3\% \pm 1.9$ for *M. oleifera* and *M. peregrina*, respectively). No significant difference was detected between the two *Moringa* species (*t*-test, $p = 0.437929$).

3.7. GC-MS Chemical Analysis of the Pollenkitt

Chemical analysis of the pollenkitt of both *M. oleifera* and *M. peregrina* revealed that the hexane extract was a waxy blend of 21 compounds. The blend included 11 *n*-alkanes, 2 fatty acids (oleic and/or palmitic acid), 4 long-chain aldehydes, 1 sesquiterpene (transnerolidol), 1 sterol (γ -sitosterol), and 2 tocopherols (both α and γ) (Table 1). Although in tiny quantities, the pollenkitt of the two *Moringa* species differed primarily in the absence of long-chain aldehydes in the *M. oleifera* pollenkitt and the near absence of tocopherols in the *M. peregrina* pollenkitt. Of the 11 alkanes identified, four constituted approximately 80% of the pollenkitt extract in both species. In *M. oleifera*, the alkanes constituted about 94% of the extracted pollenkitt. Pentacosane (*n*-C25), heptacosane (*n*-C27), nonacosane (*n*-C29), and hentriacontane (*n*-C31) comprised about 28%, 20%, 17%, and 15% of the hexane extract, respectively, amounting to 80.2% in total of the extracted pollenkitt (Table 1). In *M. peregrina*, the alkanes constituted approximately 92% of the extracted pollenkitt. The same alkanes, pentacosane (*n*-C25), heptacosane (*n*-C27), nonacosane (*n*-C29), and hentriacontane (*n*-C31), constituted approximately 24%, 24%, 19%, and 16% of the hexane extract, respectively, amounting to 82.5% in total of the extracted pollenkitt (Table 1).

Table 1. Comparative chemical analysis of the pollenkitt of *M. oleifera* and *M. peregrina* extracted by hexane.

Chemical Name	Molecular Formula	Retention Time	<i>M. oleifera</i>	<i>M. peregrina</i>
Trans-nerolidol	C ₁₅ H ₂₆ O	12.321	1.22 ± 0.31	---
9-methyl-nonadecane	C ₂₀ H ₃₂	14.859	---	0.54 ± 0.29
Palmitic acid	C ₁₆ H ₃₂ O ₂	15.138	1.32 ± 0.38	1.51 ± 0.75
Oleic acid	C ₁₈ H ₃₄ O ₂	16.268	---	0.42
Tricosane	C ₂₃ H ₄₈	17.228	3.48 ± 0.12	5.54 ± 16.
Tetracosane	C ₂₄ H ₅₀	17.78	1.22 ± 0.21	1.8 ± 0.18
N.I, long chain	Aldehyde	18.129	---	0.65 ± 0.15
Pentacosane *	C ₂₅ H ₅₂	18.327	23.74 ± 0.05	27.98 ± 3.87
N.I, long chain	Aldehyde	18.432	---	0.45
Hexacosane	C ₂₆ H ₅₄	18.828	2.31 ± 0.31	2.27 ± 0.13
N.I, long chain	Aldehyde	19.134	---	0.58 ± 0.18
Heptacosane *	C ₂₇ H ₅₆	19.392	24.34 ± 0.08	20.22 ± 1.39
Octacosane	C ₂₈ H ₅₈	19.975	1.23 ± 1.26	0.68
N.I* long chain	Aldehyde	20.085	---	3.97 ± 1.48
Nonacosane *	C ₂₉ H ₆₀	20.678	18.50 ± 0.02	17.08 ± 0.61
Triacotane	C ₃₀ H ₆₂	21.455	1.33 ± 0.88	0.94 ± 0.12
γ-Tocopherol	C ₂₈ H ₄₈ O ₂	21.995	1.41 ± 0.06	0.36
Hentriacontane *	C ₃₁ H ₆₄	22.414	15.95 ± 0.1	14.92 ± 2.15
α-Tocopherol	C ₂₉ H ₅₀ O ₂	22.788	1.37 ± 0.68	---
Tritriacontane	C ₃₃ H ₆₈	24.865	1.91 ± 0.13	---
γ-Sitosterol	C ₂₉ H ₅₀ O	25.297	0.68 ± 0.26	---
Total			100 (82.53 *)	100 (80.2 *)

* Four most abundant chemicals in the pollenkitt. The total amount of the four chemicals in parentheses. MF—molecular formula, R.T.—retention time.

Head-space analysis of the volatiles emitted by the pollen grains revealed that they constituted 33 compounds, including 13 *n*-alkanes, 2 fatty acids (palmitic and stearic), 2 aldehydes, and a list of 16 aromatic constituents (Table 2). However, while most chemical constituents occurred in *M. oleifera* (*n* = 32), *M. peregrina* had significantly fewer volatile constituents (*n* = 24) (Table 2).

Table 2. Comparative head-space chemical analysis of the volatiles emitted by the pollen grains of *M. oleifera* and *M. peregrina*.

Chemical Name	Molecular Formula	Retention Time	<i>M. oleifera</i>	<i>M. peregrina</i>
Nonanal	C ₉ H ₁₈ O	8.791	✓	✓
Trans-2-nonenal	C ₉ H ₁₆ O	9.377	✓	-----
3'-Methylacetophenone	C ₉ H ₁₀ O	9.707	✓	-----
α-Terpineol	C ₁₀ H ₁₈ O	9.791	-----	✓
Decanal	C ₁₀ H ₂₀ O	9.831	✓	✓
α-Ionol	C ₁₃ H ₂₂ O	10.256	✓	-----
3-Heptylacrolein	C ₁₀ H ₁₈ O	10.386	✓	✓
Undecanal	C ₁₁ H ₂₂ O	10.8	✓	✓
2,4-Decadienal	C ₁₀ H ₁₆ O	10.928	✓	✓
2-Undecenal	C ₁₁ H ₂₀ O	11.324	✓	✓
Myrcene	C ₁₀ H ₁₆	11.402	✓	-----
Methyl cinnamate	C ₁₀ H ₁₀ O ₂	11.561	✓	✓
Dodecyl aldehyde	C ₁₂ H ₂₄ O	11.71	✓	✓
β-Bisabolene	C ₁₅ H ₂₄	12.987	✓	✓
Tetra decyl aldehyde	C ₁₄ H ₂₈ O	13.379	✓	✓
2-undecanol	C ₁₄ H ₂₈ O	13.969	✓	✓
Pentadecanal	C ₁₅ H ₃₀ O	14.148	✓	✓
9-methylnonadecane	C ₂₀ H ₄₂	15.691	✓	✓

Table 2. Cont.

Chemical Name	Molecular Formula	Retention Time	<i>M. oleifera</i>	<i>M. peregrina</i>
Palmitic acid	C ₁₆ H ₃₂ O ₂	15.797	✓	✓
δ-tetradecalactone	C ₁₄ H ₂₆ O ₂	16.99	✓	-----
Stearic acid	C ₁₈ H ₃₆ O ₂	17.06	✓	✓
Bisphenol A	C ₁₅ H ₁₆ O ₂	17.239	✓	✓
Tricosane	C ₂₃ H ₄₈	18.028	✓	✓
δ-Stearolactone	C ₁₈ H ₃₄ O ₂	18.223	✓	-----
Tetracosane	C ₂₄ H ₅₀	18.441	✓	✓
Pentacosane	C ₂₅ H ₅₂	18.971	✓	✓
Hexacosane	C ₂₆ H ₅₄	19.485	✓	✓
Heptacosane	C ₂₇ H ₅₆	20.029	✓	✓
Octacosane	C ₂₈ H ₅₈	20.63	✓	✓
Nonacosane	C ₂₉ H ₆₀	21.312	✓	✓
triacontane	C ₃₀ H ₆₂	22.106	✓	-----
Hentriacontane	C ₃₁ H ₆₄	23.04	✓	-----
Dotriacontane	C ₃₂ H ₆₆	24.156	✓	-----

(✓) chemical present, (-----) chemical not present.

4. Discussion

Drylands cover about 50% of the global land area and provide sustenance and livelihood for about 20% of the world's population. In these lands, precipitation provides plants with less than 65% of the potential evaporation and transpiration ($ET_p < 0.65$) [23]. The most prevailing conditions in drylands are high levels of temperature and solar irradiation and low levels of relative humidity and water availability, which pose a severe threat to the successful growth and reproduction of many staple crops [24]. Here, we attempted to elucidate adaptations in the sexual reproduction of *M. oleifera* and *M. peregrina* that allow them to grow successfully and produce leaves, seeds, and oil under these conditions.

Our emphasis on the Moringa pollenkitt was due to its uniquely high relative content of pollenkitt in both *Moringa* species. Pollenkitt is an adhesive material present around pollen grains of animal pollinated angiosperms. Pacini and Hesse [25] reported more than 20 functional stages, namely in the anthers, from pollen presentation in the anthers, in relation to pollinators, during pollen dispersal, and when the pollen reaches the stigma. Among those functions, some may play significant roles in protecting pollen grains against adverse conditions in arid regions, including protection from water loss, protection from UV radiation, and facilitation of pollen rehydration [25]. However, other potential functions, such as the ones tested here, including protection against high temperatures and improving the penetration of the pollen grains into the stylar tube, were not considered.

4.1. Weight Ratio of Pollenkitt to the Pollen Grains

The amount of pollenkitt produced by the flower depends on its pollination mechanism and the specific function of the pollenkitt in the pollination process [26]. In zoophilous species, for example, the pollenkitt provides adhesion to animal bodies for dispersal as well as to the stigmatic surface and may contain volatiles that are attractive to pollinators. In contrast, the pollenkitt in anemophilous species has a minimal function in pollen dispersal and advertisement and is almost absent in some cases [27]. However, since pollen dispersal in clumps may lead to high pollination efficiency in angiosperms, pollenkitt may be found in many types of pollination mechanisms but in different amounts [26]. Haisheng et al. [28] studied the content of pollenkitt in some wind- and insect-pollinated species and reported a range from 8% in olive pollen up to 60% in dandelion pollen. They showed that pollen adhesion was directly and positively affected by pollenkitt content. Our data fall within the upper range of this pollenkitt content. Thus, additional functions other than enhanced adhesion of pollenkitt might be considered in *Moringa* species.

4.2. The Significance of the Pollenkitt for Pollen Insertion into the Styler Tube

Our results suggested that the significance of the pollenkitt for pollen insertion into the styler tube was at least threefold. First, as the insects visited the newly opened flowers, the pollen grains were evenly spread on their heads, between their antennae, in a thin and very sticky layer, thus preparing the pollen to be shuffled into the styler tube. Then, once a large insect visited a non-pollinated flower, the open style was brushed against the thin layer of pollen and scooped up the pollen grains into the hollow tube. Second, the pollenkitt provided lubrication to allow many pollen grains to penetrate the tube and reach the receptive inner tissue. Third, since the outermost exine layer of pollen grains is electrically charged, it may cause mutual repulsion [29]. The insulating properties of the thick layer of pollenkitt probably reduced the electrostatic charges on the pollen grains. Thus, very few pollen grains repulsed each other as they penetrated the styler tube. Due to their electrical insulating properties, natural plant waxes have been used by the industry for insulating coatings for the past 70 years or more [30]. Furthermore, the relatively large flowers of *M. oleifera* and *M. peregrina*, and the need to insert pollen grains into a narrow styler tube, require the pollinating services of large and powerful insects. The most frequent visitor of *Moringa* flowers, the carpenter bee (*Xylocopa* sp.), is probably a perfect match to these requirements. As it pushes itself into the flower to gather nectar, sticky pollen smears on its head and, when it visits an un-pollinated flower, the open style is brushed against the pollen and gets inserted into the style. Similarly, in *Cassia fistula* Linn., this process also requires large and heavy-bodied insects such as carpenter bees and honey bees. Out of 650,000 pollen grains produced per flower, only 250–350 are eventually pushed into the style [31].

4.3. The Significance of the Pollenkitt for Protection against Heat and Water Stress

Although they originated from two completely different regions of the world, the pollen grains of the two *Moringa* species studied showed relatively similar responses to extreme temperatures, prevalent under arid conditions. When exposed to high temperatures, slight reductions in pollen viability and germinability were found compared to those at room temperature, probably due to the desiccation of the pollen grains. The significance of the pollenkitt was revealed when the pollen grains were exposed to high temperatures, with and without the pollenkitt. Removal of the pollenkitt from *M. oleifera* pollen resulted in almost a complete loss of pollen viability under high temperatures and drastic reductions in germinability under all temperatures, probably due to desiccation. The statistically significant interaction between extreme temperatures and pollenkitt suggests a different effect of temperature depending on whether the pollen grains are coated with pollenkitt or not. Furthermore, exposure to high RH restored most of the germinability of the pollen grains, which supports our assertion that the pollenkitt plays a pivotal role in protecting the pollen grains against desiccation. Removal of the pollenkitt from *M. peregrina* pollen, however, resulted in a significantly smaller reduction of pollen viability under high temperatures, suggesting a better adaptation to high temperatures. Pollen germinability, however, was less reduced at cold to moderate temperatures of -20°C to 25°C , but was completely lost at high temperatures, suggesting that the viable pollen of *M. peregrina* require the pollenkitt to retain their germinability. However, germinability on the stigma occurs inside the styler tube with conditions that are probably less desiccating than in our experiments. Thus, we have to assume that the detrimental effects of exposure to high temperatures were primarily on over-heating and desiccation of the pollen grains. The presence of the waxy pollenkitt provided some protection against these stressful conditions. Our results are supported by the work of Pacini and Hesse [25], who reported that the function of pollenkitt in protection against water loss is often weak. In most cases, it does not provide complete protection but only slows down the desiccating process.

Chemical analysis of the pollenkitt revealed a lipidic, waxy blend of hydrocarbons, primarily *n*-alkanes (>90%), with some fatty acids, aldehydes, and aromatic volatiles. The differences between the two *Moringa* species were relatively small, suggesting a similar

role for the pollenkitt for both species. The fact that the *M. oleifera* pollenkitt constituted more volatile components may have significance related to the attractiveness of the flowers to pollinating insects and should be further explored in future research. Hydrocarbons are known to play a critical role in protection against arid conditions in both the plant and animal kingdoms. In terrestrial plants, their aerial parts, and especially the leaves, are covered by a hydrophobic cuticular layer that serves as a waterproof barrier, protecting the plants against desiccation. This layer consists of a cutin matrix as well as cuticular waxes in which very long-chain alkanes are the major components, representing up to 70% of the total wax content in the leaves. The biosynthesis of these alkanes was found to be highly linked to responses to stresses, both biotic and abiotic [32]. Leaf waxes have also been associated with protection against heat stress and cooler canopies under water limiting conditions [33,34]. In further support of this claim, Mondal et al. [35] suggested that, in wheat, production of leaf cuticular wax may contribute to lower leaf temperatures under heat stress. The epicuticular hydrocarbons of scorpions also play an important role in protecting them against desiccation and water loss under hot and arid conditions [36]. Those hydrocarbons act as an efficient barrier that limits transpiration. The most abundant *n*-alkanes were *n*-heptacosane (C_{27} ; $19 \pm 2\%$ of total HCs), *n*-nonacosane (C_{29} ; $16 \pm 1\%$), and *n*-hentriacontane (C_{31} ; $11 \pm 1\%$) [36]. Interestingly, these three *n*-alkanes are also three of the four most abundant *n*-alkanes in the *Moringa* pollenkitt and their relative abundance corresponds almost identically to their abundance in the pollenkitt. This type of desiccation resistance has been determined as an adaptive life-history trait dependent upon the presence of hydrocarbons known to conserve internal water stores by preventing the loss of water due to transpiration [37].

4.4. The Significance of the Styler Tube for Protection against Heat and Water Stress

Our results revealed that the styler tube of both *M. oleifera* and *M. peregrina* flowers protected the pollen grains against the stressful conditions imposed by the extreme temperatures, both high and low. Once inserted into the styler tube, the pollen grains retained their relatively high levels of viability even when the pollenkitt was removed. Thus, the styler tube provided maximal protection against heat and water stresses and offered optimal conditions for pollen germination on the inner stigmatic tissue. Only very few similar germinating conditions have been reported in the plant kingdom, as in *Cyclamen persicum* (Mill.) flowers, where it has been reported that *Cyclamen* pollen grains must also enter a hollow style to germinate [38]. The authors suggested no functional advantage, but since *Cyclamen* flowers bloom in the Mediterranean winter and the styles point downwards, we can assume that the hollow style provides the germinating pollen grains with some protection against the falling rain.

5. Conclusions

Our study attempted to elucidate the significance of floral and pollen traits of two *Moringa* species, one originating from the wet tropics (*M. oleifera*) and the other from the dry tropics (*M. peregrina*), for their reproductive success under arid conditions. Our results provide evidence that the pollenkitt coating the pollen grains is unique in quantity and quality. Making up the majority of the pollen weight, the pollenkitt protects against extreme temperatures and desiccation for both species. The unusual hollow style, with its narrow styler tube, requires the pollination services of large and powerful insects and provides the pollen grains with protection against stressful arid conditions. Additional functionality for the pollenkitt was discovered in providing lubrication for the pollen grains in their passage into the styler tube and in reducing undesirable electrostatic repulsion. We conclude that the reproductive success of *M. oleifera* and *M. peregrina* under arid conditions is enhanced mainly by their unique floral and pollen traits.

Author Contributions: Y.V., A.M., and D.E. conceived the project. Y.V., A.M., and D.E. collected data and conducted the analyses. Y.V. and A.M. wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This research was partially supported by the Fuel Choices Initiative at the Israeli Prime Minister's office.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data will be provided by the authors upon request.

Acknowledgments: The authors wish to thank Hanita Zemach and Eduard Belausov in their assistance with the SEM and confocal microscopes and Hiba Haj-Yehya for her assistance in the lab and fieldworks.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Olson, M.E.; Carlquist, S. Stem and root anatomical correlations with life form diversity, ecology, and systematics in *Moringa* (Moringaceae). *Bot. J. Linn. Soc.* **2001**, *135*, 315–348. [\[CrossRef\]](#)
- Ramachandran, C.; Peter, K.V.; Gopalakrishnan, P.K. Drumstick (*Moringa oleifera*): A multipurpose Indian vegetable. *Econ. Bot.* **1980**, *34*, 276–283. [\[CrossRef\]](#)
- Sengupta, A.; Gupta, M.P. Studies on the seed fat composition of Moringaceae family. *Fett. Wiss. Technol.* **1970**, *72*, 6–10. [\[CrossRef\]](#)
- Kleiman, R.; Ashley, D.A.; Brown, J.H. Comparison of two seed oils used in cosmetics, *Moringa* and *Marula*. *Ind. Crop. Prod.* **2008**, *28*, 361–364. [\[CrossRef\]](#)
- Rashid, U.; Anwar, F.; Moser, B.R.; Knothe, G. *Moringa oleifera* oil: A possible source of biodiesel. *Bioresour. Technol.* **2008**, *99*, 8175–8179. [\[CrossRef\]](#)
- Caceres, A.; Cabrera, O.; Morales, O.; Mollinedo, P.; Mendia, P. Pharmacological properties of *Moringa oleifera*. 1: Preliminary screening for antimicrobial activity. *J. Ethnopharmacol.* **1991**, *33*, 213–216. [\[CrossRef\]](#)
- Holst, S. *Moringa: Nature's Medicine Cabinet*; Sierra Sunrise Books: Sherman Oaks, CA, USA, 2000; Volume 14622.
- Jahn, S.A.A.; Mussad, H.A.; Burgstaller, H. The tree that purifies water: Cultivating multipurpose Moringaceae in the Sudan. *Unasylva* **1986**, *152*, 23–28.
- Hegazy, A.K.; Hammouda, O.; Lovett-Doust, J.; Gomaa, N.H. Population dynamics of *Moringa peregrina* along altitudinal gradient in the northwestern sector of the Red Sea. *J. Arid Environ.* **2008**, *72*, 1537–1551. [\[CrossRef\]](#)
- Alaklbi, A. Genetic diversity of *Moringa peregrina* species in Saudi Arabia with ITS sequences. *Saudi J. Biol. Sci.* **2015**, *22*, 186–190. [\[CrossRef\]](#)
- Wilkinson, A. *The Garden in Ancient Egypt*; Rubicon Press: London, UK, 1998.
- Somali, M.A.; Bajneid, A.; Fhainian, S.S. Chemical composition and characteristic of *Moringa peregrina* seeds and seed oil. *J. Am. Oil Chem. Soc.* **1984**, *16*, 85–86. [\[CrossRef\]](#)
- Mridha, M.A.U. Prospects of *Moringa* cultivation in Saudi Arabia. *J. Appl. Environ. Biol. Sci.* **2015**, *5*, 39–46.
- Abud-Archila, M.; Espinosa-Arriola, A.K.; González-Soto, T.; Gutiérrez-Oliva, V.F.; Ruiz-Valdiviezo, V.; González-Mendoza, D.; Rodríguez-Hernández, L.; Gutiérrez-Miceli, F.A. Growth and biochemical responses of *Moringa* (*Moringa oleifera* L.) to vermicompost and phosphate rock under water stress conditions. *Phyton* **2018**, *87*, 209–215.
- Vaknin, Y.; Mishal, A. The potential of the tropical “miracle tree” *Moringa oleifera* and its desert relative *Moringa peregrina* as edible seed-oil and protein crops under Mediterranean conditions. *Sci. Hort.* **2017**, *225*, 431–437. [\[CrossRef\]](#)
- Decraene, L.R.; De Laet, J.; Smets, E.F. Floral development and anatomy of *Moringa oleifera* (Moringaceae): What is the evidence for a capparalean or sapindalean affinity? *Ann. Bot.* **1998**, *82*, 273–284. [\[CrossRef\]](#)
- Jyothi, P.V.; Atluri, J.B.; Reddi, C.S. Pollination ecology of *Moringa oleifera* (Moringaceae). *Proc. Plant Sci.* **1990**, *100*, 33–42.
- Krieg, J.; Goetze, D.; Porembski, S.; Arnold, P.; Linsenmair, K.E.; Stein, K. Floral and reproductive biology of *Moringa oleifera* (Moringaceae) in Burkina Faso, West Africa. *Acta Hort.* **2017**, *1158*, 63–69. [\[CrossRef\]](#)
- Bhattacharya, A.; Mandal, S. Pollination, pollen germination and stigma receptivity in *Moringa oleifera* Lamk. *Grana* **2004**, *43*, 48–56. [\[CrossRef\]](#)
- Dafni, A.; Neppi, M.; Pacini, E. Pollen and stigma biology. In *Practical Pollination Biology*; Dafni, A., Kevan, P.G., Husband, B., Eds.; Enviroquest: Cambridge, ON, Canada, 2005; pp. 83–147.
- Evans, D.E.; Taylor, P.E.; Singh, M.B.; Knox, R.B. Quantitative analysis of lipids and protein from the pollen of *Brassica napus* L. *Plant Sci.* **1991**, *73*, 117–126. [\[CrossRef\]](#)
- Martin, F.W. Staining and observing pollen tubes in the style by means of fluorescence. *Stain Technol.* **1959**, *34*, 125–128. [\[CrossRef\]](#) [\[PubMed\]](#)
- Middleton, N.; Thomas, D. (Eds.) *World Atlas of Desertification*, 2nd ed.; Hodder Headline Group: London, UK, 1997.
- Beaumont, P. *Drylands: Environmental Management and Development*; Routledge: London, UK; New York, NY, USA, 2014.
- Pacini, E.; Hesse, M. Pollenkitt—Its composition, forms and functions. *Flora-Morphol. Distrib. Funct. Ecol. Plants* **2005**, *200*, 399–415. [\[CrossRef\]](#)

26. Pacini, E.; Franchi, G.G. Some cytological, ecological and evolutionary aspects of pollination. *Acta Soc. Bot. Pol.* **1996**, *65*, 11–16. [[CrossRef](#)]
27. Heslop-Harrison, J. An interpretation of the hydrodynamics of pollen. *Am. J. Bot.* **1979**, *1*, 737–743. [[CrossRef](#)]
28. Haisheng, L.; Gomez, I.; Meredith, J.C. Pollenkitt wetting mechanism enables species-specific tunable pollen adhesion. *Langmuir* **2013**, *29*, 3012–3023.
29. Galati, B.G.; Rosenfeld, S. The pollen development in *Ceiba insignis* (Kunth) Gibbs & Semir ex *Chorisia speciosa* St. Hil. (Bombacaceae). *Phytomorphology* **1998**, *48*, 121–130.
30. Hodge, W.H.; Sineath, H.H. The Mexican candelilla plant and its wax. *Econ. Bot.* **1956**, *10*, 134–154. [[CrossRef](#)]
31. Saradhi, P.P.; Ram, H.M. Some Aspects of floral biology of *Cassia fistula* Linn. (The Indian laburnum)—Part I. *Curr. Sci.* **1981**, *50*, 802–805.
32. Bourdenx, B.; Bernard, A.; Domergue, F.; Pascal, S.; Léger, A.; Roby, D.; Pervent, M.; Vile, D.; Haslam, R.P.; Napier, J.A.; et al. Overexpression of *Arabidopsis* ECERIFERUM1 promotes wax very-long-chain alkane biosynthesis and influences plant response to biotic and abiotic stresses. *Plant Physiol.* **2011**, *156*, 29–45. [[CrossRef](#)]
33. Richards, R.A.; Rawson, H.M.; Johnson, D.A. Glaucousness in wheat: Its development and effect on water—Use efficiency, gas exchange and photosynthetic tissue temperatures. *Aust. J. Plant Physiol.* **1986**, *13*, 465–473. [[CrossRef](#)]
34. Bennett, D.; Izanloo, A.; Edwards, J.; Kuchel, H.; Chalmers, K.; Tester, M.; Reynolds, M.P.; Schnurbusch, T.; Langridge, P. Identification of novel quantitative trait loci for days to ear emergence and flag leaf glaucousness in bread wheat (*Triticum aestivum* L.) population adapted to southern Australian conditions. *Theor. Appl. Genet.* **2012**, *124*, 697–711. [[CrossRef](#)] [[PubMed](#)]
35. Mondal, S.; Mason, R.E.; Huggins, T.; Hays, D.B. QTL on wheat (*Triticum aestivum* L.) chromosomes 1B, 3D and 5A are associated with constitutive production of leaf cuticular wax and may contribute to lower leaf temperatures under heat stress. *Euphytica* **2015**, *201*, 123–130. [[CrossRef](#)]
36. Gefen, E.; Talal, S.; Brendzel, O.; Dror, A.; Fishman, A. Variation in quantity and composition of cuticular hydrocarbons in the scorpion *Buthus occitanus* (Buthidae) in response to acute exposure to desiccation stress. *Comp. Biochem. Phys. A* **2015**, *182*, 58–63. [[CrossRef](#)] [[PubMed](#)]
37. Krupp, J.J.; Nayal, K.; Wong, A.; Millar, J.G.; Levine, J.D. Desiccation resistance is an adaptive life-history trait dependent upon cuticular hydrocarbons, and influenced by mating status and temperature in *D. melanogaster*. *J. Insect Physiol.* **2020**, *121*, 103990. [[CrossRef](#)] [[PubMed](#)]
38. Reinhardt, S.; Ewalda, A.; Hellwig, F. The Anatomy of the stigma and style from *Cyclamen persicum* (Mill.) cv. “pure white” and its relation to pollination success. *Plant Biol.* **2007**, *9*, 158–162. [[CrossRef](#)] [[PubMed](#)]