



Table S1. Minimum, mean and maximum number of *C. chalcites* alive larvae observed on banana bunches at 1 day prior to application and 1, 3 and 7 days post application of biorational products.

Treatment	Day -1			Day 1			Day 3			Day 7		
	Min	Mean	Max	Min	Mean	Max	Min	Mean	Max	Min	Mean	Max
Prevam®	17	26.75	35	4	8	13	1	1.5	2	0	0	0
Cinamite®	23	25.75	28	8	9	11	4	7.75	10	3	4.25	7
Intruder®	5	8	12	2	2.75	4	1	2	3	0	1.5	3
Avenger®	14	17	22	6	10	15	4	7.75	10	3	6.5	10
Indasol®	17	21.75	34	12	16.5	20	3	6.25	8	0	3.25	5
Ripels®	18	25	32	12	16	21	10	13	16	4	6.75	11
Bioknock®	13	15.5	19	6	9.5	13	4	5.75	6	4	4.75	6
Garlitol Forte®	12	17.5	23	7	10.75	15	4	5.75	8	0	2	4
Control	12	22.5	32	10	19	27	9	16	23	7	14	20


a.




b.




c.




d.




e.



f.



g.



h.




Figure S1. Survival analysis of *C. chalcites* larvae 1 day before, 1, 3, 7 days after the application of (a) Bioknock®, (b) Intruder®, (c) Avenger®, (d) Cinamite®, (e) Garlitrol Forte®, (f) Ripelser®, (g) Prevam® and (h) Indasol® (neem oil).