

# The physiological role of inulin in wild cardoon (*Cynara cardunculus* L. var. *sylvestris* Lam.)

Ferdinando Branca<sup>1</sup>, Sergio Argento<sup>2</sup>, Anna Maria Paoletti<sup>3</sup> and Maria Grazia Melilli<sup>2\*</sup>

Supplementary data

**Table S1.** Pearson product moment correlation coefficient ( $r^2$ ) among sugars and enzymes in roots of wild cardoon. \*\*\*;  
\*\*and \* indicate significant at  $p < 0.001$ , 0.05 and 0.01, respectively, and ns not significant

Glucose	Fructose	Sucrose	Inulin	1-SST	1-FFT	1-FEH	Invertase
Vegetative phase							
Glucose	ns <sup>1</sup>	ns	0.624*	ns	ns	ns	0.760**
Fructose		ns	ns	ns	ns	ns	ns
Sucrose			0.727**	0.920***	-0.843***	0.731**	ns
Inulin				0.727**	0.785**	ns	ns
1-SST					-0.888***	0.717***	ns
1-FFT						-0.653*	ns
1-FEH							ns
Invertase							
Reproductive phase							
Glucose	Fructose	Sucrose	Inulin	1-SST	1-FFT	1-FEH	Invertase
Glucose	ns	-0.764*	ns	0.989***	ns	0.986***	0.897**
Fructose		0.900**	ns	ns	ns	ns	ns
Sucrose			0.800*	ns	ns	-0.708*	ns
Inulin				-0.706*	ns	-0.733*	ns
1-SST					ns	0.998***	0.927***
1-FFT						-0.726*	ns
1-FEH							0.909**
Invertase							