

Supplementary Material for

Inoculum Location in The Soil Influences Damage of *Globisporangium ultimum*, *Fusarium culmorum* and *Rhizoctonia solani* and Effectiveness of Control Agents in Maize Seedlings

Eckhard Koch ¹, Petra Zink ¹, Tanja Bernhardt ¹, Tim Birr ² and Ada Linkies ^{1,*}

¹ Julius Kühn Institute, Federal Research Centre for Cultivated Plants, Institute for Biological Control, 69221 Dossenheim, Germany

² Department of Plant Diseases and Crop Protection, Institute of Phytopathology, Faculty of Agricultural and Nutritional Science, Christian-Albrechts-University of Kiel, 24118 Kiel, Germany

* Author for correspondence: Ada Linkies (Ada.Linkies@Julius-Kuehn.de)




Figure S1. Habitus of maize seedlings (A) eight and (B) 14 days after sowing in pots with inoculum of *Rhizoctonia solani* located in different positions. h.c.: healthy control, potting substrate not inoculated; (1) inoculum mixed homogenously in entire pot volume; (2) inoculum present only in upper half of pots; (3) inoculum present only in lower half of pots.



Figure S2. Maize seedlings germinated in non-inoculated potting substrate and placed five days after planting on non-inoculated potting substrate (control; **left**) or on potting substrate inoculated with *Rhizoctonia solani* (**right**).