



Correction

Correction: Beelman et al. Soil Disturbance Impact on Crop Ergothioneine Content Connects Soil and Human Health. *Agronomy* 2021, 11, 2278

Robert B. Beelman ^{1,*}, John P. Richie, Jr. ², Allen T. Phillips ³, Michael D. Kalaras ¹, Dongxiao Sun ⁴ ⁶ and Sjoerd W. Duiker ⁵

- Department of Food Science, College of Agricultural Sciences, Penn State University, 202 Rodney A. Erickson Food Science Building, University Park, State College, PA 16802, USA; mdk219@psu.edu
- Department of Public Health Sciences, College of Medicine, Penn State University, 500 University Dr., Hershey, PA 17033, USA; jrichie@pennstatehealth.psu.edu
- Department of Biochemistry and Molecular Biology, Eberly College of Science, Penn State University, 203A South Frear Building, University Park, State College, PA 16802, USA; atp@psu.edu
- Department of Pharmacology, Mass Spectrometry Core Facilities, College of Medicine, Penn State University, 500 University Dr., Hershey, PA 17033, USA; dsun@pennstatehealth.psu.edu
- Department of Plant Science, College of Agricultural Sciences, Penn State University, 408 ASI Building, University Park, State College, PA 16802, USA; sduiker@psu.edu
- * Correspondence: rbb6@psu.edu

1. Text Correction

There was an error in the original publication. On the fourth line of Section 3: Results, the value of 32% needs to be changed to 31%.

A correction has been made to Section 3. Results, "The effect of soil tillage on the ERGO content of maize, soybeans, and oats is presented in Figure 1. A trend was observed where ERGO concentrations generally declined as tillage intensity increased from NT to CD to MB. ERGO content was decreased from NT to MB by 32%, 33%, and 28% for maize, soybeans, and oats, respectively".

Corrected sentence: "The effect of soil tillage on the ERGO content of maize, soybeans, and oats is presented in Figure 1. A trend was observed where ERGO concentrations generally declined as tillage intensity increased from NT to CD to MB. ERGO content was decreased from NT to MB by 31%, 33%, and 28% for maize, soybeans, and oats, respectively".

2. Error in Figures

In the original publication [1], there was a mistake in Figures 1 and 3, as published. In the calculations for ergothioneine content, the authors failed to account for a dilution factor of the samples, as mentioned in the methodology, prior to injecting them into the LC/MS system. Therefore, these values need to be multiplied by a factor of 2. The corrected Figures 1 and 3 appear below.

The authors apologize for any inconvenience caused and state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.



Citation: Beelman, R.B.; Richie, J.P., Jr.; Phillips, A.T.; Kalaras, M.D.; Sun, D.; Duiker, S.W. Correction: Beelman et al. Soil Disturbance Impact on Crop Ergothioneine Content Connects Soil and Human Health. *Agronomy*, 2021, 11, 2278. *Agronomy* 2022, 12, 1817. https://doi.org/10.3390/agronomy12081817

Received: 19 June 2022 Accepted: 24 June 2022 Published: 31 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

Agronomy **2022**, 12, 1817 2 of 2




Figure 1. ERGO content of crops (**a**) maize (2018), (**b**) soybean (2019), and (**c**) oats (2020) grown using different tillage methods; different letters above the bars indicate significant difference (p > 0.05) within each crop type, error bars represent the standard deviation of each tillage type.




Figure 3. Ergothioneine production per hectare of crops (a) maize (2018), (b) soybean (2019), and (c) oats (2020) grown using different tillage methods; different letters above the bars indicate significant difference (p > 0.05) within each crop type, error bars represent the standard deviation of each tillage type.

Reference

1. Beelman, R.B.; Richie, J.P., Jr.; Phillips, A.T.; Kalaras, M.D.; Sun, D.; Duiker, S.W. Soil Disturbance Impact on Crop Ergothioneine Content Connects Soil and Human Health. *Agronomy* **2021**, *11*, 2278. [CrossRef]