

**Table S1. Average body sizes (from head to wing edge) of the adult insects tested**

Group of test insects	Test insects	Scientific name	Abbreviation	Body size (mm) <sup>a</sup>
Small size	Whiteflies	<i>Bemisia tabaci</i> (Gennadius) (Hemiptera: Aleyrodidae)	WH	0.82 ± 0.17
	Western flower thrips	<i>Frankliniella occidentalis</i> (Pergande) (Thysanoptera: Thripidae)	WFT	1.41 ± 0.44
	Tomato leaf miners (syn. vegetable leaf miner)	<i>Liriomyza sativae</i> Blanchard (Diptera: Agromyzidae)	TLM	1.94 ± 0.56
Middle size	Shore flies	<i>Scatella stagnalis</i> (Fallén) (Diptera: Ephydriidae)	SF	3.12 ± 0.27
	Vinegar flies	<i>Drosophila melanogaster</i> Meigen (Diptera, Drosophilidae)	VF	3.29 ± 0.43
	Winged green peach aphids	<i>Myzus persicae</i> (Sulzer) (Hemiptera: Aphididae)	GPA-w	3.65 ± 0.29
Large size	Houseflies (male)	<i>Musca domestica</i> (Linnaeus) (Diptera: Muscidae)	HF-m	6.05 ± 0.73
	Greenbottle flies (male)	<i>Lucilia sericata</i> (Meigen) (Diptera: Calliphoridae)	GBF-m	6.31 ± 0.31
	Houseflies (female)		HF-f	8.15 ± 0.68
	Greenbottle flies (female)		GBF-f	8.81 ± 0.41

<sup>a</sup> Means and SD calculated from 20 adults of each species.