

Supplementary material

Table S3. Morphological parameters of the wheat and Fusarium head blight severity.

Treatment ¹	Plant height (cm)	Ear length (cm)	Fusarium head blight severity (%)
T0	65 ^a	6 ^{ab}	10 ^a
T1	63 ^c	4 ^a	43 ^b
T2	67 ^{ab}	7 ^{bc}	7 ^a
T3	63 ^c	6 ^{ab}	28 ^c
T4	68 ^b	8 ^c	7 ^a
T5	65 ^a	8 ^c	25 ^c
T6	68 ^b	9 ^c	5 ^a
T7	66 ^a	8 ^c	24 ^c

¹ T0 – control; T1 – *F. culmorum*; T2 – herbicide + fungicide 1 + fungicide 2; T3 – herbicide + fungicide 1 + fungicide 2 + *F. culmorum*; T4 – herbicide + fungicide 1 + fungicide 2 + humic acids; T5 - herbicide + fungicide 1 + fungicide 2 + humic acids + *F. culmorum*; T6 – herbicide + fungicide 1 + fungicide 2 + nitrophenols; and T7 – herbicide + fungicide 1 + fungicide 2 + nitrophenols + *F. culmorum*. The values of each parameter are averaged. The same letter indicates no statistically significant difference between treatments ($P \geq 0.05$), determined with ANOVA followed by Fisher's test.