Modeling Soil Organic Carbon Dynamics Under Two Cropping Modes in Salinized Paddy Fields in the Yellow River Delta

Minghui Li [1,2], Jia Dong [1], Sijia Guo [1,2], Deyong Zhao [1,3], Chunhong Wu [1,3], Jikun Xu [1,2,3,4], Liping Zhao [1,3], Jun Wang [1,3], Haiyang Wang [4], Jianlin Wang [5] and Shuaipeng Zhao [1,2,3,4,*]

[1] College of Biological and Pharmaceutical Engineering, Shandong University of Aeronautics, Binzhou 256603, China; liminghui@stu.sdua.edu.cn (M.L.); 18766142198@163.com (J.D.); guosijia1222@163.com (S.G.); dyzhao@sdua.edu.cn (D.Z.); wuchunhong.123@163.com (C.W.); xujikun@sdua.edu.cn (J.X.); zhaoliping_bz@163.com (L.Z.); ivywangjun@163.com (J.W.)
[2] Shandong Key Laboratory of Eco-Environmental Science for the Yellow River Delta, Binzhou 256603, China
[3] Shandong Province Engineering and Technology Research Center for Wild Plant Resources Development and Application of Yellow River Delta, Shandong University of Aeronautics, Binzhou 256603, China
[4] The Yellow River Delta Sustainable Development Institute of Shandong Province, Dongying 257100, China; wwhhyy20000@163.com
[5] Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China; jianlinwang@genetics.ac.cn
* Correspondence: zhaoshuaipeng@sdua.edu.cn

Supplementary Figure S1. Some field experimental plots



Supplementary Figure S2. Schematic plan of diagonal sampling method and field soil sample collection





