

Article

# Chronic Chemogenetic Activation of the Superior Colliculus in Glaucomatous Mice: Local and Retrograde Molecular Signature

Marie Claes <sup>1</sup>, Emiel Geeraerts <sup>1</sup>, Stéphane Plaisance <sup>2</sup>, Stephanie Mentens <sup>1,3,4</sup>, Chris Van den Haute <sup>5,6</sup>, Lies De Groef <sup>3</sup>, Lut Arckens <sup>4</sup> and Lieve Moons <sup>1,\*</sup>

- <sup>1</sup> Neural Circuit Development and Regeneration Research Group, Department of Biology, KU Leuven, Leuven Brain Institute, 3000 Leuven, Belgium; marie.claes@kuleuven.be (M.C.); emiel\_geeraerts@hotmail.com (E.G.); stephanie.mentens@kuleuven.be (S.M.)
- <sup>2</sup> VIB Nucleomics core, 3000 Leuven, Belgium; stephane.plaisance@vib.be
- <sup>3</sup> Cellular Communication and Neurodegeneration Research Group, Department of Biology, KU Leuven, Leuven Brain Institute, 3000 Leuven, Belgium; lies.degroef@kuleuven.be
- <sup>4</sup> Neuroplasticity and Neuroproteomics Research Group, Department of Biology, KU Leuven, Leuven Brain Institute, 3000 Leuven, Belgium; lut.arckens@kuleuven.be
- <sup>5</sup> Neurobiology and Gene Therapy Research Group, Department of Neurosciences, KU Leuven, Leuven Brain Institute, 3000 Leuven, Belgium; chris.vandenhaute@kuleuven.be
- <sup>6</sup> KU Leuven Viral Vector Core, 3000 Leuven, Belgium
- \* Correspondence: lieve.moons@kuleuven.be

## Supplementary information

**Table S1.** Mean values  $\pm$  SEM for the different techniques used to probe glaucomatous induction (i.e. IOP and anterior chamber depth) and neuroprotection (i.e. pSTR recordings), corresponding to Figure 9a-c.

Experimental group	Mean $\pm$ SEM
IOP (area under the curve)	
SHAM	26.84 $\pm$ 0.53
DREADD	27.36 $\pm$ 0.45
OHT	34.97 $\pm$ 0.93
OHT + DREADD	33.68 $\pm$ 1.00
Anterior chamber depth ( $\mu$ m)	
SHAM	388.87 $\pm$ 2.48
DREADD	385.11 $\pm$ 3.28
OHT	472.22 $\pm$ 9.91
OHT + DREADD	472.04 $\pm$ 10.62
pSTR ( $\mu$ V)	
SHAM	160.79 $\pm$ 5.61
DREADD	170.74 $\pm$ 5.22
OHT	129.90 $\pm$ 5.94
OHT + DREADD	121.83 $\pm$ 4.63

**Table S2.** Estimation statistics (Hedges' *g* values with corresponding confidence intervals) for the different techniques used to probe glaucomatous induction (i.e. IOP and anterior chamber depth) and neuroprotection (i.e. pSTR recordings), corresponding to Figure 9a-c.

Comparison	Unpaired Hedges' <i>g</i>	95% Confidence interval
IOP		
SHAM VS. DREADD	0.18	[-0.31, 0.67]
SHAM VS. OHT	1.75	[1.24, 2.23]
SHAM VS. OHT + DREADD	1.42	[0.92, 1.87]
ANTERIOR CHAMBER DEPTH		
SHAM VS. DREADD	-0.22	[-0.74, 0.26]
SHAM VS. OHT	1.87	[1.45, 2.28]
SHAM VS. OHT + DREADD	1.78	[1.28, 2.37]
pSTR		
SHAM VS. DREADD	0.32	[-0.16, 0.83]
SHAM VS. OHT	-0.92	[-1.4, -0.43]
SHAM VS. OHT + DREADD	-1.32	[-1.8, -0.81]